This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

PLEASE DO NOT REMOVE THIS BAND

REMOTE STORAGE

Please return at the circulation desk.
To renew your material call:
(650) 723-6691 ext. 3

Date due in Lane Library:

Ba2

LANE
MEDICAL LIBRARY

SEAL OF THE LELAND STANFORD JUNIOR UNIVERSITY
SEMPER VIRENS
1885

GIFT
SHRINERS' HOSPITAL

AMERICAN BANK NOTE CO LITHO

2140.

William Major Collins

Nov. 1902.

N.Y. Homeopathic M.C.

Ba

MEDICAL

William Major Collin

Nov. 1902.

N. Y. Homeopathic M. C.

Copyright 1888, 1889, 1890, 1891,
By W. B. SAUNDERS.

PRESS OF
W. B. SAUNDERS, PHILADA.

FOURTEENTH THOUSAND

ESSENTIALS OF ANATOMY

AND

MANUAL OF PRACTICAL DISSECTION,

TOGETHER WITH THE

ANATOMY OF THE VISCERA.

PREPARED ESPECIALLY FOR

STUDENTS OF MEDICINE.

BY

CHARLES B. NANCREDE, M.D.,

PROFESSOR OF SURGERY AND OF CLINICAL SURGERY IN THE UNIVERSITY OF MICHIGAN, ANN ARBOR; CORRESPONDING MEMBER OF THE ROYAL ACADEMY OF MEDICINE, ROME, ITALY; LATE SENIOR SURGEON TO EPISCOPAL HOSPITAL; LATE SURGEON TO JEFFERSON MEDICAL COLLEGE HOSPITAL; FORMERLY LECTURER ON OSTEOLOGY, ETC., IN MEDICAL DEPARTMENT, UNIVERSITY OF PENNSYLVANIA; LATE PROFESSOR OF GENERAL AND ORTHOPÆDIC SURGERY IN PHILADELPHIA POLYCLINIC, ETC.

FOURTH EDITION REVISED AND ENLARGED BY AN APPENDIX CONTAINING

HINTS ON DISSECTION,

BY J. CHALMERS DA COSTA, M.D.,

BASED UPON THE LAST EDITION OF GRAY'S ANATOMY.

30 HANDSOME FULL-PAGE LITOGRAPHIC PLATES, IN COLORS,
AND **188** FINE WOODCUTS.

PHILADELPHIA:
W. B. SAUNDERS,
925 WALNUT STREET.
1894.

PREFACE TO FIRST EDITION.

THE author has endeavored in this little book to embody only those facts which have appeared to him to be really the "essentials of anatomy;" not that he considers it likely that the student will master every minute detail therein contained, but he believes that the knowledge gained by a study of this work will enable the future practitioner, during the remainder of his professional life, to recall such general impressions as will render intelligible current medical literature, or even the more elaborate monographs, and will at once suggest *where* to consult his anatomical text-books for such terms or facts as may have become indistinct through lapse of time.

While this book cannot replace the larger anatomical works, sufficient descriptive matter has been introduced to enable the student to refresh his memory of the more numerous facts learnt in the lecture and dissecting room, or from his "Gray" or other text-book, differing in this respect from most of the works of its class, which are little more than a list of names, without any distinctive facts connected with them to aid the student in the difficult task of acquiring a knowledge of a branch of medical study almost solely dependent upon the unassisted powers of the memory.

Conciseness, rather than elegance of diction, has been the aim, so that all words such as the articles "a," "an," "the," have been omitted, except where absolutely necessary.

Recognizing that a work of this kind should, as far as possible, conform to that text-book most commonly used, the last edition of Gray's *Anatomy* has been chosen as the chief authority, although free use has been made of the works of Quain, Leidy, Bock, Allen, Morris *On the Joints*, Starr *On Diseases of Children* (dentition), Tomes *Dental Anatomy*, Potter, Frey, Holden, Politzer, H. Thompson, Astley Cooper *On the Breast*, and original work of one of the author's former students.

The author would here acknowledge his obligations and return his thanks to Prof. Joseph Leidy, of the University of Pennsylvania, for the kind permission to reproduce numerous cuts from the first edition of his *Anatomy*, and to Dr. F. M. Varrell for much valuable assistance in the correction of proof.

CHARLES B. NANCREDE.

PHILADELPHIA, September, 1888.

PUBLISHER'S PREFACE TO THIRD EDITION.

THE favorable acceptance of this little volume, by both Students and Practitioners, since its first publication in the Fall of 1888, together with a still ever increasing demand, (it being now in its tenth thousand,) has suggested to the Publisher the addition of a series of Osteological plates, which, by following the teachings and arrangement of " Gray's Anatomy," should render the work of even greater service than before to the novice in Anatomical Science.

CONTENTS.

PAGE
Introduction 17
Bones of the head 23
    The frontal bone 23
    Parietal bones 25
    Occipital bone 27
    The sphenoid 30
    The temporal bones 34
    Ethmoid bone 38
    Nasal bones 40
    The superior maxillary bones 40
    Lachrymal bones 44
    Malar bones 44
    The palate bones 45
    The inferior turbinated bones 47
    Vomer 48
    The inferior maxillary bone 49
    The sutures and fontanelles 51
    The cerebral fossæ 53
    The orbital cavities 55
    The nasal fossæ 58
Bones of the trunk 61
    The vertebral column 61
    The thorax 64
    Hyoid bone 65
    The sternum 65
    Ribs 66
    The pelvic bones 68
    The pelvis 73
    Bones of the upper extremity 74

PAGE
- The arm 78
- The forearm 80
- The hand 82

Lower extremity 85
- The thigh 85
- The patella 87
- The leg 88
- The foot 90

The articulations 92
- Muscles and fasciæ 114
- Muscles of the head 116
- Muscles of the neck 120
- Palatal region 122
- Vertebral region 122
- Muscles of the back 123
- Muscles of the abdomen 129
- Muscles of thorax 132
- Muscles of the forearm 136
- Fasciæ of the hand 139
- Muscles of the hand 140
- Muscles of lower extremity 141
- Muscles of the leg 147
- Fasciæ of foot 149
- Muscles of the foot 150
- Vascular system 152
- The arteries 157
- The veins 177
- The lymphatics 184
- Nervous system 185
- The spinal cord 198
- The cranial nerves 200
- The spinal nerves 205
- Dorsal nerves 208
- The sympathetic nerve 211

Visceral anatomy 216
- The digestive organs 216
- The teeth 217

PAGE
The mouth 220
The pharynx 224
The œsophagus 226
The stomach 227
The small intestine 229
The large intestine 231
The liver 233
The pancreas 238
The ductless glands 239
The spleen 239
The thyroid gland 241
The thymus gland 242
The suprarenal capsules 242
The abdominal cavity 243
The peritoneum 246
Organs of voice and respiration 249
The larynx 249
The trachea and bronchi 254
The lungs 256
The pleuræ 259
The urinary organs 260
The ureters 265
The bladder 266
The male urethra 269
Female urethra 271
The male generative organs 271
The penis 272
The testes and their coverings 274
Female organs of generation 279
The vagina 281
The uterus and its appendages 281
The mammary glands 286
The organs of sense 287
The skin and its appendages 287
Organs of taste 289
Organs of smell 290
The eye 292

PAGE
Sclerotic and cornea 293
The humors 300
Appendages of eye 306
Tutamina oculi 306
The ear 309
The external ear 310
The middle ear or tympanum 311
The tympanum 313
The internal ear or labyrinth 318
Anatomy of inguinal hernia 325
Femoral hernia 328
Perineum and ischio-rectal regions 330
Introduction to osteological plates 336
Series of osteological plates from Gray's Anatomy 337

PLATE 1.

SUPERFICIAL DISSECTION OF THE THORAX AND ABDOMEN.

A.—Upper bone of the sternum.

B.B.*—Two first ribs.

C.C.*—Second pair of ribs.

D.D.*—Right and left lungs.

E.—Pericardium, enveloping the heart—the right ventricle.

F.—Lower end of sternum.

G.G.*—Lobes of the liver.

H.H.*—Right and left halves of the diaphragm, in section; the right half separating the right lung from the liver, the left half separating the left lung from the broad cardiac end of the stomach.

I.I.*—Eighth pair of ribs.

K.K.*—Ninth pair of ribs.

L.L.*—Tenth pair of ribs.

M.M.*—The stomach; m, its cardiac bulge; m,* its pyloric extremity.

N.—The umbilicus.

O.O.*—The transverse colon.

P.P.*—The omentum, covering the transverse colon and small intestines.

Q.—The gall bladder.

R.R.*—The right and left pectoral prominences.

S.S.*—The small intestines.

PLATE I.



PLATE II.



PLATE 2.

RELATIVE POSITION OF THE DEEPER ORGANS OF THORAX AND THOSE OF THE ABDOMEN.

A.—Upper end of the sternum.

B.B.*—First pair of ribs.

C.C.*—Second pair of ribs.

D.—Aorta, with left vagus and phrenic nerves crossing its transverse arch.

E.—Root of pulmonary artery.

F.—Right ventricle.

G.—Right auricle.

H.—Vena cava superior, with right phrenic nerve on its outer border.

I.I.*—Right and left lungs collapsed, and turned outward to show the heart's outline.

K.K.*—Seventh pair of ribs.

L.L.*—The diaphragm, in section.

N.—The gall bladder, with its duct to form the common bile duct. The hepatic artery is seen superficial to the common duct; the vena portæ is seen beneath it. The patent orifices of the hepatic veins are seen on the cut surface of the liver.

O.—The stomach.

P.—Inferior vena cava.

Q.—The cœliac axis, dividing into the gastric, splenic and hepatic arteries.

R.—The spleen.

S.S.* S.**—The transverse colon, between which and the lower border of the stomach is seen the gastro-epiploic artery, formed by the splenic and hepatic arteries.

S.***—Ascending colon, in the right iliac region.

T.—Convolutions of the small intestines, distended with air.

PLATE 3.

DEEP DISSECTION OF THROAT AND ABDOMEN.

A.—The thyroid body.
B.—The trachea.
C.C.*—The first ribs.
D.D.*—The clavicles, cut at middle.
E.—Humeral part of great pectoral muscle, cut.
F.—Coracoid process of the scapula.
G.—Arch of the aorta; **G,*** descending aorta in throat.
H.—Right bronchus; **H,*** left bronchus.
I.—Œsophagus.
K.—Vena azygos, receiving the intercostal veins.
L.—Thoracic duct.
M.M.*—Seventh ribs.
N.N.—Diaphragm in section.
O.—Cardiac orifice of stomach.
P.—Liver in section, showing patent orifices of hepatic veins.
Q.—Cœliac axis, sending off branches to liver, stomach and spleen. The stomach has been removed to show the looping anastomosis of these vessels around the superior and inferior borders of the stomach.
R.—Inferior vena cava, about to enter its notch in the posterior thick part of the liver, to receive the hepatic veins.
S.—Gall-bladder, communicating by its duct with the hepatic duct, which is lying upon the vena portæ, and by the side of the hepatic artery.
T.—The pyloric end of stomach, joining **T,*** duodenum.
U.—The spleen.
V.V.—The pancreas.
W.—The sigmoid flexure of colon.
X.—The caput coli.
Y.—The mesentery, supporting the numerous looping branches of the superior mesenteric artery.
Z.—Some coils of the small intestine.

2.—Innominate artery.
3.—Right subclavian artery.
4.—Right common carotid artery.
5.—Left subclavian artery.
6.—Left common carotid artery.
7.—Left axillary artery.
8.—Coracoid attachment of the smaller pectoral muscle.
9.—Subscapular muscle.
10.—Coracoid head of the biceps muscle.
11.—Tendon of the latissimus dorsi muscle.
12.—Superior mesenteric artery, with its accompanying vein.
13.—Left kidney.

PLATE III.



PLATE IV.



PLATE 4.

RELATION OF THE PRINCIPAL BLOOD-VESSELS OF THE THORAX AND ABDOMEN TO THE OSSEOUS SKELETON, ETC.

A.—Arch of the aorta.

B.B.—Descending thoracic, part of the aorta giving off *b*, *b*, the intercostal arteries.

C.—Abdominal part of aorta.

D.D.—First pair of ribs.

E.—The xyphoid cartilage.

F.—Right crus of diaphragm.

G.G.*—Right and left kidneys.

H.—Brachio-cephalic artery.

I.—Left common carotid artery.

K.—Left subclavian artery.

L.—Right common iliac artery at its place of division.

M.—Left common iliac artery, seen through the meso-rectum.

N.—Inferior vena cava.

O.O.—Sigmoid flexure of the colon.

P.—The rectum.

Q.—Urinary bladder.

R.—Right iliac fossa.

S.S.—Right and left ureters.

T.—Left common iliac vein, joining the right under the right common iliac artery to form the inferior vena cava.

U.—Fifth lumbar vertebra.

V.—External iliac artery of right side.

W.—The symphysis pubis.

X.—An incision made over the locality of the femoral artery.

b.b.—The dorsal intercostal arteries.

c.—The cœliac axis.

d.—Superior mesenteric artery.

f.f.—Renal arteries.

g.—Inferior mesenteric artery.

h.—The vas deferens, bending over the epigastric artery and the os pubis, after having passed through the internal abdominal ring.

PLATE 5.

SUPERFICIAL DISSECTION OF THE HEAD AND NECK.

A.—Third portion of the subclavian artery.
B.—Sterno-mastoid muscle.
C.—Common carotid artery.
D.—External " "
E.—Internal " "
F.—Continuation of the external carotid, through the parotid gland.
G.—Temporal artery.
H.—External jugular vein.
I.—Brachial plexus.
K.—Posterior belly of omo-hyoid.
L.—Suprascapular artery.
M.—Transverse cervical artery.
N.—Scalenus anticus.
O.—Glandulæ concatenatæ.
P.—Superficial descending cervical nerves.
Q.—Great auricular nerve.
R.—Occipital artery and nerve.
S.—Facial nerve.
T.—Duct of Stenson.
U.—Facial vein.
V.—Facial artery.
W.—Submaxillary gland.
X.—Digastric muscle.
Y.—Lymphatic gland.
Z.—Hyoid bone.
1.—Thyroid cartilage.
2.—Superior thyroid artery.
3.—Anterior jugular vein.
4.—Anterior belly of omo-hyoid.
5.—Sterno-hyoid (right).
6.—Inter-clavicular notch.
7.—Clavicle.
8.—Trapezius muscle.
9.—Splenius capitis.
10.—Posterior belly of occipito-frontalis.
11.—Attollens aurem.
12.—Anterior belly of occipito-frontalis.
13.—Orbicularis palpebrarum.
14.—Zygomaticus major.
15.—Buccinator.
16.—Depressor anguli oris.

PLATE V.



PLATE VI.



PLATE 6.

DISSECTION OF THE DEEP CERVICAL AND FACIAL REGIONS.

A.—Innominate artery, at its point of bifurcation.
B.—Subclavian artery, crossed by the vagus nerve.
C.—Common carotid artery, with the vagus nerve at its outer side, and the descendens noni nerve lying on it.
D.—External carotid artery.
E.—Internal carotid artery, with the descendens noni nerve lying on it.
F.—Lingual artery passing under the fibres of the hyoglossus muscle.
G.—Tortuous facial artery.
H.—Temporo-maxillary artery.
I.—Occipital artery, crossing the internal carotid artery and jugular vein.
K.—Internal jugular vein, crossed by some branches of the cervical plexus, which join the descendens noni nerve.
L.—Spinal accessory nerve, which pierces the sterno-mastoid muscle, to be distributed to it and the trapezius.
M.—Cervical plexus of nerves giving off the phrenic nerve to descend the neck on the outer side of the internal jugular vein, and over the scalenus muscle.
N.—Vagus nerve, between the carotid artery and internal jugular vein.
O.—Ninth, or hypoglossal nerve, distributed to the muscles of the tongue.
P.P.—Branches of the brachial plexus of nerves.
Q.—Subclavian artery in connection with the brachial plexus of nerves.
R.R.—Post-scapular artery passing through the brachial plexus.
S.—Transversalis humeri artery.
T.—Transversalis colli artery.
U.—Union of the post-scapular and external jugular veins, which enter the subclavian vein by a common trunk.
V.—Posterior half of the omo-hyoid muscle.
W.—Part of the subclavian vein, seen above the clavicle.
X.—Scalenus muscle, separating the subclavian artery from vein.
Y.—Clavicle.
Z.—Trapezius muscle.

1.—Sternal origin of sterno-mastoid muscle of left side.
2.—Clavicular origin of sterno-mastoid muscle of right side, turned down.
3.—Scalenus posticus muscle.
4.—Splenius muscle.
5.—Mastoid insertion of sterno-mastoid muscle.
6.—Internal maxillary artery, passing behind the neck of lower jaw-bone.
7.—Parotid duct.
8.—Genio-hyoid muscle.
9.—Mylo-hyoid muscle, cut and turned aside.
10.—Superior thyroid artery.
11.—Anterior half of omo-hyoid muscle.
12.—Sterno-hyoid muscle, cut.
13.—Sterno-thyroid muscle, cut.

PLATE 7.

DEEP DISSECTION OF THE HEAD AND NECK.

A.—Root of common carotid artery.
B.—Subclavian artery at its origin.
C.—Trachea.
D.—Thyroid axis of subclavian artery.
E.—Vagus nerve, crossing the origin of subclavian artery.
F.—Subclavian artery at third division of its arch.
G.—Post-scapular branch of subclavian artery.
H.—Transversalis humeri branch of subclavian artery.
I.—Transversalis colli branch of subclavian artery.
K.—Posterior belly of omo-hyoid muscle, cut.
L.—Median nerve branch of brachial plexus.
M.—Musculo-spiral branch of same plexus.
N.—Anterior scalenus muscle.
O.—Cervical plexus giving off the phrenic nerve, which takes tributary branches from brachial plexus of nerves.
P.—Upper part of internal jugular vein.
Q.—Upper part of internal carotid artery.
R.—Superior cervical ganglion of sympathetic nerve.
S.—Vagus nerve lying external to sympathetic nerve, and giving off *t*, its laryngeal branch.
T.—Superior thyroid artery.
U.—Lingual artery, separated by hyoglossus muscle from
V.—Hypoglossal or ninth cerebral nerve.
W.—Sublingual salivary gland.
X.—Genio-hyoid muscle.
Y.—Mylo-hyoid muscle, cut and turned aside.
Z.—Thyroid cartilage.

1.—Upper part of sterno-hyoid muscle.
2.—Upper part of omo-hyoid muscle.
3.—Inferior constrictor of pharynx.
4.—Cricoid cartilage.
5.—Crico-thyroid muscle.
6.—Thyroid body.
7.—Inferior thyroid artery of thyroid axis.
8.—Sternal tendon of sterno-mastoid muscle, turned down.
9.—Clavicular portion of sterno-mastoid muscle, turned down.
10.—Clavicle.
11.—Trapezius muscle.
12.—Scalenus posticus muscle.
13.—Rectus capitis anticus major muscle.
14.—Stylo-hyoid muscle, turned aside.
15.—Temporal artery.

PLATE VII.



PLATE 7.—CONTINUED.

16.—Internal maxillary artery.
17.—Inferior dental branch of fifth pair of cerebral nerves.
18.—Gustatory branch of fifth pair of nerves.
19.—External pterygoid muscle.
20.—Internal pterygoid muscle.
21.—Temporal muscle, cut to show the deep temporal branches of fifth pair of nerves.
22.—Zygomatic arch.
23.—Buccinator muscle, with buccal nerve and parotid duct.
24.—Masseter muscle, cut on the lower maxilla.
25.—Middle constrictor of pharynx.

PLATE 8.

DEEP DISSECTION OF THE LATERAL REGION OF THE NECK.

1.—Auricular cartilage.
2.—Parotid gland.
3.—Digastric.
4.—Sterno-cleido mastoid.
5.—Trapezius.
6.—Splenius capitis.
7.—Mastoid process.
8.—Trachelo-mastoid muscle.
9.—Complexus.
10.—Obliquus capitis superior.
11.— " " inferior.
12.—Splenius colli.
13.—Scalenus medius.
14.—Common carotid artery.
15.—Internal carotid artery.
16.—External carotid artery.
17.—Hypoglossal nerve.
18.—Middle sterno-mastoid artery.
19.—Princeps cervicis artery.
20.—Occipital artery.
21.—Posterior auricular artery.

PLATE VIII.



PLATE IX.



PLATE 9.

DISSECTION OF THE POSTERIOR ASPECT OF THE NECK AND HEAD.

1.—Trapezius.

2.
3. } Splenius capitis et colli.

4.—Trachelo-mastoid.

5.—Biventer cervicis and complexus.

6.—Semispinalis colli.

7.—Spinous process of axis.

8.—Obliquus capitis inferior.

9.—Rectus capitis posticus major.

10.— " " " minor.

11.—Spinous process of atlas (first).

12.—Sterno-cleido-mastoid.

13.—Digastric.

14.—Posterior arch of atlas.

15.—Vertebral artery.

16.—Occipital artery.

17.—Sigmoid vein (split), passing between the arches of the first and second vertebra. It has many connections with the veins both without and within the spinal canal.

18.—Section of occipital bone.

19.—Right transverse sinus.

20.—Torcular Herophili.

21.—Longitudinal sinus.

22.—Dura mater.

23.—Occipitalis major.

24.—Posterior branch of the third cervical nerve.

25.—Occipitalis minor.

26.—Superficial occipital and mastoid glands.

27.—Section of the skin.

28.—Subcutaneous fat, including the large blood-vessels in its deepest layer.

29.—Section of the aponeurosis occipito-frontalis.

30.—Section of the parietal bone, showing diploe, inner and outer tables.

[illegible] striae & optic thr. goes off at rt. ∠s, [illegible] in
[illegible], underneath → Ant. Perf. space.
[illegible]
[illegible] → F. Rolando. [illegible] = P. parietal in F. [illegible]
[illegible] temporal

PLATE 10.

ARTERIAL SUPPLY TO THE BASE OF THE BRAIN.

A.A.—Vertebral arteries.

B.—Anterior spinal branches, forming median anterior trunk.

D.—Posterior inferior cerebellar artery.

E.—Basilar trunk, from convergence of two vertebrals.

F.F.—Transverse arteries.

G.—Anterior inferior cerebellar artery.

H.—Superior cerebellar artery.

I.—Posterior cerebral artery, terminal branching of basilic.

J.—Choroidean artery, posterior.

K.—Posterior communicating artery.

L.—Carotid artery, internal.

M.—Anterior choroidean artery. Comm. w. Post. Choroid = 2° Circle

N.—Middle cerebral artery.

O.—Anterior cerebral artery.

P.—Point of reflection of anterior cerebral over the corpus callosum.

R.—Anterior communicating artery.

PLATE X.



PLATE XI.



PLATE 11.

DISSECTION OF THE AXILLA.

A.—Axillary vein, drawn apart from the artery to show the nerves lying between both vessels. On the bicipital border of the vein is seen the internal cutaneous nerve; on the tricipital border is the nerve of Wrisberg, communicating with some of the intercosto-humeral nerves; *a*, the common trunk of the venæ comites, entering the axillary vein.

B.—Axillary artery crossed by one root of the median nerve; *b*, basilic vein forming with *a*, the axillary vein, **A.**

C.—Coraco-brachialis muscle.

D.—Coracoid head of biceps muscle.

E.—Pectoralis major muscle.

F.—Pectoralis minor muscle.

G.—Serratus magnus muscle, covered by *g*, the axillary fascia, and perforated at regular intervals by nervous branches called intercosto-humeral.

H.—Conglobate gland, crossed by nerve called "external respiratory" of Bell, distributed to the serratus magnus muscle. This nerve descends from the cervical plexus.

I.—Subscapular artery.

K.—Tendon of latissimus dorsi muscle.

L.—Teres major muscle.

PLATE 12.

DISSECTION OF THE AXILLARY AND BRACHIAL REGIONS.

A.—Axillary vein cut and tied; *a*, the basilic vein cut.

B.—Axillary artery; *b*, brachial artery, in the upper part of its course, having *h*, the median nerve, lying rather to its outer side; *b*,* the artery on the lower part of its course, with the median nerve to its inner side.

C.—Subclavius muscle.

C.*—Clavicle.

D.—Axillary plexus of nerves, of which *d* is a branch on the coracoid border of the axillary artery; *e*, the musculo-cutaneous nerve, piercing the coraco-brachialis muscle; *f*, the ulnar nerve; *g*, musculo-spiral nerve; *h*, the median nerve; *i*, the circumflex nerve.

E.—Humeral part of the great pectoral muscle.

F.—Biceps muscle.

G.—Coraco-brachialis muscle.

H.—Thoracic half of the lesser pectoral muscle.

I.—Thoracic half of the greater pectoral muscle.

K.—Coracoid attachment of the lesser pectoral muscle.

K.*—Coracoid process of the scapula.

L.—Lymphatic glands.

M.—Serratus magnus muscle.

N.—Latissimus dorsi muscle.

O.—Teres major muscle.

P.—Long head of triceps muscle.

Q.—Inner condyle of humerus.

PLATE XII.



PLATE XIII.

Fig 1.



Fig. 2.



PLATE 13.

SUPERFICIAL AND DEEP DISSECTION OF THE BEND OF THE ELBOW AND FOREARM.

FIG. 1.

A.—Fascia, covering the biceps muscle.

B.—Basilic vein, with internal cutaneous nerve.

C.—Brachial artery, with the venæ comites.

D.—Cephalic vein, with the external cutaneous nerve; *d*, the median nerve.

E.—A communicating vein, joining the venæ comites.

F.—Median basilic vein.

G.—Lymphatic gland.

H.—Radial artery at its middle.

I.—Radial artery of the pulse.

K.—Ulnar artery, with ulnar nerve.

L.—Palmaris brevis muscle.

FIG. 2.

A.—Biceps muscle.

B.—Basilic vein, cut.

C.—Brachial artery.

D.—Median nerve; *d*, ulnar nerve.

E.—Brachialis anticus muscle; *e*, the internal condyle.

F.—Origin of radial artery.

G.—Supinator radii longus muscle.

H.—Aponeurosis of the tendon of the biceps muscle.

I.—Pronator radii teres muscle.

K.—Flexor carpi ulnaris muscle.

L.—Flexor carpi radialis muscle.

M.—Palmaris longus muscle.

N.—Radial artery at middle, with radial nerve on outer side.

O.—Flexor digitorum sublimis.

P.—Flexor pollicis longus.

Q.—Median nerve.

R.—Lower end of radial artery.

S.—Lower end of ulnar artery, in company with ulnar nerve.

T.—Pisiform bone.

U.—Extensor metacarpi pollicis.

PLATE 14.

DISSECTION OF THE WRIST AND HAND.

FIG. 1.

A.—Radial artery.
B.—Median nerve; *b*,*b*,*b*, its branches.
C.—Ulnar artery, forming **F**, the superficial palmar arch.
D.—Ulnar nerve; **E**,*e*,*e*, its continuation, branching to the little and ring fingers etc.
G.—Pisiform bone.
H.—Abductor muscle of the little finger.
I.—Tendon of flexor carpi radialis muscle.
K.—Opponens pollicis muscle.
L.—Flexor brevis muscle of the little finger.
M.—Flexor brevis pollicis muscle.
N.—Abductor pollicis muscle.
O.O.O.O.—Lumbricale muscles.
P.P.P.P.—Tendons of the flexor digitorum sublimis muscle.
Q.—Tendon of the flexor longus pollicis muscle.
R.—Tendon of extensor metacarpi pollicis.
S.—Tendons of extensor digitorum sublimis; **P**,**P**,**P**, their digital prolongations.
T.—Tendon of flexor carpi ulnaris.
U.—Union of the digital arteries at the tip of the fingers.

FIG. 2.

A.—Radial artery.
B.—Tendons of the extensors of the thumb.
C.—Tendon of extensor carpi radialis.
D.—Annular ligament.
E.—Deep palmar arch, formed by radial artery giving off *e*, the artery of the thumb.
F.—Pisiform bone.
G.—Ulnar artery, giving off the branch **I**, to join the deep palmar arch **E**, of the radial artery.
H.—Ulnar nerve; *h*, superficial branches given to the fingers. Its deep palmar branch is seen lying on the interosseous muscles, **M**,**M**.
K.—Abductor minimi digiti.
L.—Flexor brevis minimi digiti.
M.—Palmar interossei muscles.
N.—Tendon of flexor digitorum sublimis and profundus, and the lumbricales muscles, cut and turned down.
O.—Tendon of flexor pollicis longus.
P.—Carpal end of the metacarpal bone of the thumb.

PLATE XIV.



Fig. 3.



Fig. 2.



Fig. 1.

PLATE 14.—Continued.

FIG. 3.

A.A.A.—Tendons of extensor digitorum communis; **A***, tendon overlaying that of indicator muscle.

B.—Dorsal part of the annular ligament

C.—End of the radial nerve, distributed over the back of the hand, to two of the fingers and the thumb.

D.—Dorsal branch of the ulnar nerve, supplying the back of the hand and the three outer fingers.

E.—Radial artery, turning round the carpal end of the metacarpal bone of the thumb.

F.—Tendon of extensor carpi radialis brevior.

G.—Tendon of extensor carpi radialis longior.

H.—Tendon of third extensor of the thumb.

I.—Tendon of second extensor of the thumb.

K.—Tendon of extensor minimi digiti, joining a tendon of extensor communis.

PLATE 15.

SUPERFICIAL DISSECTION OF THE BACK.

1.—Acromion.
2.—Trapezius.
3.—Latissimus dorsi.
4.—Rhomboideus major.
5.—Infraspinatus.
6.—Teres minor.
7.—Teres major.
8 —Deltoid.
9.—Triceps—long head.
10. { Dorsalis scapulæ artery.
11. { " " "
12.—Intercostal nerves.
13 —Rhomboideus minor.
14.— " major.
15.—Dorsal fascia.
16.—Splenius capitis.
17.—Intercostal arteries.
18.—Transverse scapular artery.
19.—Dorsal nerves.
20 —Fifth to ninth ribs.
21.—Inferior angle of scapula.
22.—Spine of scapula.
23.—Acromion.
24.—Clavicle.
25.—Transverse ligament.
26.—Capsular ligament.
27.—Insertion of rhomboideus major.
28.— " " minor.
29.—Levator anguli scapulæ.
30.—Clavicular insertion of trapezius.
31 —Supraspinatus.
32.—Infraspinatus.
33.—Deltoid (cut across).
34.—Teres minor.
35.—Serratus magnus.
36.—External intercostals.
37.—Dorsalis scapulæ artery, nerve and vein.
38.—Transverse scapular artery.
39.— " " vein and nerve.
40.—Supra-scapular artery, nerve and vein.
41.—Recurrent or circumflex, scapular artery and vein.
42.—External oblique.
43.—Intercostal nerves.
44.—Serratus posticus inferior.
45.—Three lower ribs.

PLATE XV.


37
39
25
33
16
30
23
24
29
12
40
14
41
2
1
18
4
5
10
15
21
17
19
13
3
20
36
45
43
44
46
45
42
49
52
53
48
57
51
64
55
57
60
50
67
68
58
62
61
66
63
59
56
54

PLATE 15.—Continued.

46.—Tendinous origin of latissimus dorsi.
47.—External oblique, back part.
48.—Crest of ileum.
49.—Internal oblique.
50.—Gluteus maximus.
51.— " medius.
52.—Lowest intercostal nerve.
53.—Superior cutaneous nerves.
54.—Inferior " "
55.—Sacrum.
56.—Coccyx.
57.—Origin of gluteus maximus.
58.—Greater trochanter.
59.—Great sacro-sciatic ligament.
60.—Pyriformis.
61.—Obturator internus.
62.—Gemelli muscles.
63.—Quadratus femoris.
64.—Superior gluteal artery and nerve.
55.—Great sciatic nerve.
66.—Sciatic artery and vein.
67.—Cutaneous nerve to posterior femoral region.
68.—Inferior gluteal nerve.

PLATE 16.

DEEP DISSECTION OF THE BACK.

1.—Subscapular muscle, cross section.
2.—Serratus magnus " "
3.—Teres major.
4.—Triceps, long head.
5.—Teres minor, origin.
6.—Acromion.
7.—Humerus, anatomical neck.
8.—Splenius capitis.
9.—Splenius colli.
10.—Transversalis colli muscle.
11.—Cervicalis ascendens muscle.
12.—Trachelo-mastoid.
13.—Complexus.
14.—Semi-spinalis dorsi.
15.—Spinalis dorsi.
16.—Longissimus dorsi.
17.—Sacro-lumbalis; tendinous extension inserted into the nine upper ribs.
18.—Cervicalis ascendens, costal origin from the six upper ribs.
19.—Intercostal muscle (external).
20.—Intercostal artery and vein (passing between the inner and outer layer of the internal intercostal muscle).
21.—Supra-spinatus.
22.—Capsule of shoulder joint.
23.—Deltoid.
24.—Triceps muscle, external head.
25.—Triceps " long head.
26.—Teres minor.
27.—Teres major.
28.—Latissimus dorsi.
29.—Sacro-lumbalis.
30.—Venous plexus of the spinal canal.
31.—Lumbar aponeurosis (fascia).
32.—Union of the leaflets of the lumbar fascia.
33.—Levatores costarum breviores.
34.— " " longiores.
35.—Roots of the lumbar and sacral nerves.
36.—Internal oblique muscle.
37.—Transversalis "
38.—Intercostal artery, vein and nerve (twelfth).
39.—Ileo-hypogastric nerve.
40.—Great sacro-sciatic foramen.
41.—Neck of femur.
42.—Trochanter major.
43.—Trochanter minor.

PLATE XVI.



9
11
10
12
21
22
13
5
14
23
24
25
16
27
28
17
19
35
29
36
37
39
38
56
52
51
53
40
48
47
42
57
54
41
49
50
58
46
44
43
55
45

PLATE 16.—Continued.

44.—Insertion of psoas and iliacus into trochanter minor.
45.—Tuberosity of the ischium.
46.—External sphincter ani.
47.—Coccyx.
48.—Great sacro-sciatic ligament.
49.—Obturator internus, cut across immediately after passing under great sacro-sciatic ligament.
50.—Levator ani.
51.—Pyriformis, cross section immediately after its exit from the foramen magnum.
52.—Gluteal artery, veins and nerve.
53.—Sciatic nerve.
54.—Ischiatic artery.
55.—Internal pudic artery.
56.—Gluteus medius muscle.
57.—Capsule of hip joint.
58.—Tendinous insertion of iliacus and psoas.

PLATE 17.

DISSECTION OF PRINCIPAL BLOOD-VESSELS AND NERVES OF THE ILIAC AND FEMORAL REGIONS.

A.—The aorta at its point of bifurcation.

B.—Anterior superior iliac spine.

C.—The symphysis pubis.

D.—Poupart's ligament, immediately above which are seen the circumflex iliac and epigastric arteries, with the vas deferens and spermatic vessels.

E.E.*—Right and left iliac muscles, covered by the peritonæum; the external cutaneous nerve is seen through the membrane.

F.—The vena cava.

G.G.*—The common iliac arteries giving off the internal iliac branches on the sacro-iliac symphyses; *g, g*, the right and left ureters.

H.H.*—Right and left common iliac veins.

I.I.—Right and left external iliac arteries; each is crossed by the circumflex iliac vein.

K.K.—Right and left external iliac veins.

L.—Urinary bladder, covered by peritonæum.

M.—The rectum intestinum.

N.—The profundus branch of femoral artery.

O.—Femoral vein; *o*, saphena vein.

P.—Anterior crural nerve.

Q.—Sartorius muscle, cut.

S.—Pectinæus muscle.

T.—Adductor longus muscle.

U.—The gracilis muscle.

V.—The tendinous sheath given off from the long adductor muscle, crossing the vessels, and becoming adherent to the vastus internus muscle (forming Hunter's Canal).

W.—Femoral artery. The letter is on the part where the vessel becomes first covered by the sartorius muscle.

PLATE XVII.



PLATE XVIII.



PLATE 18.

DISSECTION SHOWING THE SURGICAL RELATIONS OF THE EXTERNAL ILIAC ARTERY AND THE INTERNAL ABDOMINAL RING.

A.—That part of the ilium which abuts against the sacrum.

B.—The spine of the ischium.

C.—The tuberosity of the ischium.

D.—The symphysis pubis.

E.—Situation of the anterior superior iliac spine.

F.—Crest of the ilium.

G.—Iliacus muscle.

H.—Psoas magnus muscle, supporting the spermatic vessels.

I.—Transversalis muscle.

K.—Termination of the sheath of the rectus muscle.

L^1. L^2. L^3.—The iliac, transverse and pelvic portions of the transversalis fascia.

M.—The peritoneum lining groin.

N.—The epigastric vessels lying between the peritoneum, **M**, and the transversalis fascia, **L^2**.

O.—The umbilical ligament.

P.—The neck of the sac of an internal inguinal hernia, formed before the spermatic vessels.

Q.—An interval which occasionally occurs between the umbilical ligament and the epigastric artery.

R. and **Q.**—Situations where the direct inguinal hernia occurs when, as in this case, the umbilical ligament crosses the space named the internal fossa—the triangle of Hesselbach.

S.—Lower part of the spermatic cord.

T.—The bulb of the urethra.

U.—External iliac vein, covered by the peritoneum.

V.—External iliac artery, covered by the peritoneum.

W.—Internal iliac artery.

X.—Common iliac artery.

PLATE 19.

DISSECTION OF THE MALE PELVIC ORGANS.

A.—The part of the sacrum which joins the ilium.

B.—The external iliac artery, cut across.

C.—The upper part of the rectum.

D.—The ascending pubic ramus.

E.—The spine of the ischium, cut.

F.—The horizontal pubic ramus, cut.

G.—Summit of the bladder, covered by the peritoneum; **G***, its side not covered by the peritoneal membrane.

H.H.—Recto-vesical peritoneal pouch.

I.—The vas deferens.

K.—The ureter.

L.—Vesicula seminalis.

M.—The spermatic cord.

N.—Corpus cavernosum penis; *n*, its artery.

O.—Urethra; *o*, the bulbous urethra.

P.—Sphincter ani muscle.

Q.—The coccyx.

R.—The sacro-sciatic ligament.

S.—Pudic artery and nerve.

T.—The sacral nerves.

U.—Pyriformis muscle, cut.

V.—The prostate.

W.—Lower part of the rectum.

X.—Deep perineal fascia.


C
A
G
H
I
U
T
F
G*
H
M
D
E
K
R
L
N
V
O
Q
W
o
P



GEO. S. HARRIS & SONS, LITH. PH

PLATE 20.

SIDE VIEW OF THE FEMALE PELVIS.

B.—Bladder (turned down).
R.—Rectum.
L.—Round ligament.
U.—Uterus.
O.—Ovary.
V.—Vagina.
S.—Sacro-iliac synchondrosis.
K.—Kidney.
T.—Fallopian tube.
P.—Pubic symphysis.
a.—Pyriformis muscle (cut).
b —Gluteal muscles.
c.—Coccygeus muscle.
d.—Obturator internus.
e.—Psoas magnus.
f.—Linea alba.
g.g.—Ureters.
h.—Obturator nerve.
i.—Internal abdominal ring.
k.—Great sacro-sciatic ligament.
1.—Abdominal aorta.
2.—Inferior mesenteric artery.
3, 3.—Common iliac arteries.
4.—Left external iliac artery.
5.—Vena cava inferior.
6, 6.—Renal veins.
7, 7.—Common iliac veins.
8.—External iliac vein.
9.—Internal iliac artery (cut).
10.—Gluteal vein.
11.—Ilio-lumbar vein.
12.—Lateral sacral vein.
13.—Sciatic vein.
14.—Pudic vein.
15.—Obturator vein.
16.—Epigastric vein.
17.—Uterine veins.
18.—Vesico-vaginal veins.
19.—Ovarian veins.
20.—Bulb of the ovary.
21.—Vein to round ligament.
22.—Fallopian veins.

PLATE 21.

SHOWING THE RELATIONS OF THE INTERNAL ILIAC ARTERY AND THE POSITION OF AN ISCHIATIC HERNIA.

1.—Cross section of two last lumbar vertebræ.
2.— " " " sacrum.
3.— " " " coccyx.
4.—Vertebral canal.
5.—Cross section of the pubic bone.
6.—Right pubic arch.
7.—Tuberosity of the ischium.
8.—Horizontal ramus of the pubis.
9.—Anterior superior spinous process of ilium.
10.—Crest of the ilium.
11.—Gluteus maximus.
12.—Greater sacro-sciatic ligament.
13.—Lesser sacro-sciatic ligament (covered by coccygeus muscle).
14.—Pyriformis muscle.
15.—Obturator internus.
16.—Obturator canal.
17.—Iliacus internus.
18.—Psoas major.
19.—Rectus femoris.
20.—Sartorius.
21.—Adductor longus.
22.—Gracilis.
23.—Adductor magnus.
24.—Iliac artery.
25.—External iliac artery.
26.—Deep epigastric artery.
27.—Circumflex iliac artery.
28.—Femoral artery.
29.—External iliac vein.
30.—Superior obturator vein.
31.—Internal iliac artery.
32.—Ileo-lumbar artery.
33.—Superior gluteal artery.
34.—Lateral sacral artery.
35.—Vesical artery.
36.—Anterior branch of internal iliac artery.
37.—Obturator artery.
38.—Middle hemorrhoidal artery.
39.—Internal pudic artery.
40.—Inferior gluteal or ischiatic artery.
41.—Internal pudic vein.


4
10
2
2
46
18
17
32
40
34
9
28
46
52
8
37
27
38
26
30
50
39
51
11
16
13
15
5
31
12
20
19
6
11
11
21
7
22
23

PLATE 21.—CONTINUED.

42.—Ischiatic vein.
43.—Union of two last mentioned veins into one vessel.
44.—Inferior obturator vein.
45.—Superior gluteal vein.
46.—Internal iliac vein.
47.—Fifth and portion of fourth lumbar nerves, forming with the sacral nerves the sacral plexus.
48.—Obturator nerve.
49.—Anterior branches of the sacral nerves.
50.—Sciatic nerve (ischiatic), formed of two lower lumbar and two upper sacral nerves.
51.—Pudic nerve.
52.—Mouth of the sack of an ischiatic hernia, above the pyriformis and anterior to the superior gluteal artery, with the obturator artery above and the inferior obturator vein below.

PLATE 22.

SUPERFICIAL DISSECTION OF THE INGUINO-FEMORAL REGION.

A.—The umbilicus.

B.—The upper margin of the pubic symphysis.

C.—The anterior superior spine of the left iliac bone.

D.—Position of internal abdominal ring.

E.—The saphenous opening in the fascia lata, receiving *e*, the saphenous vein.

F.—The lax and pendulous cord, overlying upper part of saphenous opening.

G.—Lymphatic glands, lying on fascia lata, in the neighborhood of the saphenous opening.

H.—The fleshy part of the external oblique muscle.

a.—Superficial fascia of the abdomen.

b.—Same fascia, forming an envelope for the spermatic cord and scrotum.

c.—Inguinal glands, lying near Poupart's ligament.

d.—A common venous trunk, formed by branches from thigh and abdomen, and joining—

e.e.—The saphenous vein.

f.—The middle cutaneous nerve, derived from the anterior crural nerve.

g.—Femoral lymphatic glands.

h.—Superficial external iliac vein.

i.—Superficial epigastric vein.

k.—External cutaneous branches of nerves from the lumbar plexus.

Plate XXII.



Geo. S. Harris & Sons, Lith. Phila.

Plate XXIII.





PLATE 23.

DISSECTION OF THE FIFTH, SIXTH, SEVENTH AND EIGHTH LAYERS OF THE INGUINAL REGION, AND THEIR CONNECTION WITH THOSE OF THE THIGH.

FIG. 1.

A.—Anterior superior iliac spine.
B.—The umbilicus.
C.—Spine of the pubis.
D.—External oblique muscle ;* *d*, its tendon.
E.—Internal oblique muscle; *e*, its tendon.
F.—Transverse muscle ; *f*, its tendon, forming, with *e*, the conjoined tendon.
G.—The rectus muscle enclosed in its sheath.
H.—The fascia spermatica interna, covering the cord; *h*, its funnel-shaped extremity.
L.—Epigastric artery.
N.—Femoral artery ; *n*, its profunda branch.
O.—Femoral vein.
P.—Saphena vein.
Q.—Sartorius muscle.
R.—Sheath of the femoral vessels.
S.—The falciform margin of the saphenous opening.
T.—Anterior crural nerve.
U.—Pubic portion of the fascia lata.
V.—Iliac portion attached to Poupart's ligament.
W.—Lower part of the iliacus muscle.

FIG. 2.

A.B.C, refer to same parts as in Fig. 1.
D.—External oblique muscle; *d*, its tendon ; *d*,* the external ring.
E.—Internal oblique muscle.
F.N.O.P.Q.R.S.T.U. and **W**, refer to same parts as in Fig. 1.
G.—Rectus muscle laid bare.
H.h.—Fascia spermatica interna laid open, above and below *d*,* the external ring.
I.—The peritonæum, closing the internal ring.
K.—Fascia transversalis; *k*, its pubic part.
L.—Epigastric artery and veins.
M.—Spermatic artery, veins and vas deferens, bending round the epigastric artery at the internal ring; *m*, the same vessels below the external ring.
V.—Iliac part of the fascia lata.

PLATE 24.

DISSECTION OF THE ANTERIOR CRURAL REGION, THE ANKLE AND THE FOOT.

FIG. 1.

A.—Tendon of the tibialis anticus muscle.

B.—Long saphena vein.

C.C.—Tendon of the tibialis posticus muscle.

D.—The tibia; *d*, the inner malleolus.

E.E.—Tendon of the flexor longus digitorum muscle.

F.—Gastrocnemius muscle; *b*, tendo-Achillis.

G.—Soleus muscle.

H.—Tendon of the plantaris muscle.

I.I.—The venæ comites.

K.K.—Posterior tibial artery.

L.L.—Posterior tibial nerve.

DISSECTION OF INNER AND POSTERIOR ASPECT OF ANKLE AND LOWER THIRD OF LEG.

FIG. 2.

A.—Tibialis anticus muscle; *a*, its tendon.

B.—Extensor longus digitorum muscle; *b*, *b*, *b*, *b*, its four tendons.

C.—Extensor longus pollicis muscle.

D.D.—The tibia.

E.—The fibula; *e*, the outer malleolus.

F.F.—Tendon of the peronæus longus muscle.

G.G.—Peronæus brevis muscle; *i*, the peronæus tertius.

H.H.—The fascia.

K.—Extensor brevis digitorum muscle; *k*, *k*, its tendons.

L.L.—Anterior tibial artery, and nerve descending to the dorsum of the foot.

Plate XXIV.



Plate XXV,

Fig. 1.

Fig. 2.



GEO S HARRIS & SONS. LITH PHIL.

PLATE 25.

DISSECTION OF THE POPLITEAL SPACE AND THE POSTERIOR CRURAL REGION.

FIG. 1.

A.—Tendon of the gracilis muscle.
B.B. —The fascia lata.
C.C.—Tendon of the semimembranosus.
D.—Tendon of the semitendinosus muscle.
E.E.—The two heads of the gastrocnemius muscle.
F.—Popliteal artery.
G.—Popliteal vein, joined by the short saphena vein.
H.—Middle branch of the sciatic nerve.
K.—Posterior tibial nerve, continued from the middle branch of the sciatic, and extending to **K***, behind the inner ankle.
L.—Posterior (short) saphena vein.
M.M.—The fascia, covering the gastrocnemius muscle.
N.—Short (posterior) saphena nerve, formed by the union of branches from peronæal and posterior tibial nerves.
O.—Posterior tibial artery, appearing from beneath the soleus muscle in lower part of the leg.
P.—Soleus muscle, joining the tendo-achillis.
Q.—Tendon of the flexor communis digitorum muscle.
R.—Tendon of flexor longus pollicis muscle.
S.—Tendon of the peronæus longus muscle.
T.—Peronæus brevis muscle.
U.U.—Internal annular ligament, binding down the vessels, nerves and tendons in the hollow, behind the inner ankle.
V.V.—Tendo-achillis.
W.—Tendon of the tibialis posticus muscle.
X.—The venæ comites of the posterior tibial artery.

FIG. 2.

A.C.D.E.F.G.H.I, refer to same parts as in Fig. 1.
B.—Inner condyle of the femur.
K.—The plantaris muscle, lying upon the popliteal artery.
L.—Popliteus muscle.
M.M.M.—The tibia.
N.N.—The fibula.
O.O.—Posterior tibial artery.
P.—Peronæal artery.
Q.R.S.T.U.V.W, refer to same parts as in Fig. 1.
X.—The astragalus.

PLATE 26.

DISSECTION OF SOLE OF FOOT.

FIG. 1.

A.—The calcaneum.

B.—Plantar fascia, and flexor brevis digitorum muscle, cut; *b*, *b*, *b*, its tendons.

C.—Abductor minimi digiti muscle.

D.—Abductor pollicis muscle.

E.—Flexor accessorius muscle.

F.—Tendon of the flexor longus digitorum muscle, subdividing into *f*, *f*, *f*, *f*, tendons for the four outer toes.

G.—Tendon of flexor pollicis longus muscle.

H.—Flexor pollicis brevis muscle; *i*, *i*, *i*, *i*, the four lumbricales muscles.

K.—External plantar nerve.

L.—External plantar artery.

M.—Internal plantar nerve and artery.

FIG. 2.

A.—The heel covered by the integument.

B.—Plantar fascia and flexor brevis digitorum muscle, cut; *b*, *b*, *b*, the tendons of the muscles.

C.—Abductor minimi digiti.

D.—Abductor pollicis.

E.—Flexor accessorius, cut.

F.—Tendon of the flexor digitorum longus, cut; *b*, *b*, *b*, its digital ends.

G.—Tendon of the flexor pollicis.

H.—Head of the first metatarsal bone.

I.—Tendon of the tibialis posticus.

K.—External plantar nerve.

L.L.—The arch of the external plantar artery.

M.M.M.M.—The four interosseous muscles.

N.—External plantar nerve and artery, cut.

Fig.1.



Fig.2





GEO. S. HARRIS & SONS, LITH.

PLATE 27.

DISSECTION OF THE SUPERFICIAL STRUCTURES OF THE MALE PERINÆUM.

A.—Superficial fascia, covering the urethral space

B.—The sphincter ani.

C.—The coccyx.

D.D.*—The right and left ischiatic tuberosities.

E.—The anus.

F.F.—The glutei muscles.

PLATE 28.

DISSECTION OF THE DEEP STRUCTURES OF THE MALE PERINÆUM.

FIG. I.

A.—The urethra.

B.—Accelerator urinæ muscle.

C.—Central perinæal tendon.

D.D.—Right and left erector penis muscle.

E.E.—The transverse muscles.

F.—The anus.

G.G.—The ischiatic tuberosities.

H.—The coccyx.

I.I.—The glutei muscles.

K.K.—Levator ani muscle.

L.—Left artery of the bulb.

FIG. 2.

A. D. F. G. H. I. K. L, refer to same parts as in Fig. 1.

B.—The urethra.

C.—Cowper's glands, between the two layers of—

E.E.—The deep perinæal fascia.

M.—The bulb of the urethra.

Plate XXVIII.

Fig. 2.



Fig. 1.



Plate XXIX.



GEO. E. HARRIS & SONS, LITH. PHILA.

PLATE 29.

DISSECTION OF THE DEEP STRUCTURES OF THE MALE PERINÆUM.

A.B.—The urethra.

C.C.—The two lobes of the prostate.

D.D.—The two crura penis.

E.—The rectum turned down.

F.—The ischiatic tuberosities.

G.—The coccyx.

H.H.—The glutei muscles.

I.I.—The levator ani muscle.

K.—The left artery of the bulb.

L.—The vesiculæ seminales.

M.—The membranous part of the urethra.

N.—The base of the bladder.

O.—The two vasa deferentia.

P.—The sacro-sciatic ligament.

PLATE 30.

FEMALE PERINEUM AND ISCHIO-RECTAL REGION.

1.—Coccyx.
2.—Gluteus maximus.
3.—Fascia lata, inserted into pubic arch.
4.—Tuberosity of ischium.
5.—Internal sphincter ani.
6.—External " "
7.—Attachment of sphincter ani to coccyx.
8.—Levator ani, forming floor of ischio-rectal fossa.
9.—Perineum.
10.—Transversus peronei muscle.
11.—Erector clitoridis.
12.—Constrictor vaginæ.
13.—Glands of Bartholini.
14.—Urethral opening.
15.—Labia majora.
16.— " minora.
17.—Clitoris.
18.—Mons veneris.
19.—Internal pudic artery.
20.—External hemorrhoidal artery. (The three arteries of this name are shown, the middle one only being marked.)
21.—Superficial perineal artery. (Supplying anus, perineum, vaginal lips and erector clitoridis.)
22.—Transversus peronei artery.
23.—Deep branch of internal pudic artery.
24.—Artery of the bulb (arteria bulbosa).
25.—Internal pudic vein (common).
26.—External hemorrhoidal vein. (Other branches of the same vessel not marked).


19
25
23
22
21
4
24
3
11
10
2
20
18
15
15
13
12
9
5
6
1
16
14
17
15
8
26
10
3
11
2
25

PRACTICAL

HINTS ON DISSECTION.

BY

JOHN CHALMERS DA COSTA, M.D.,
ASSISTANT DEMONSTRATOR OF SURGERY, JEFFERSON MEDICAL COLLEGE,
PHILADELPHIA;
SURGICAL REGISTRAR OF THE PHILADELPHIA HOSPITAL ETC.

COPYRIGHTED, 1891.

PREFACE.

THIS addition to Prof. NANCREDE'S excellent Handbook makes no pretence to originality, but is taken largely from the very valuable practical work of Weisse, of New York, whose methods of procedure are by far the best of which I have any knowledge. I have also employed "Clarke and Lockwood's Manual," "Wilson's Anatomy," and Gray's classical work, by Keen.

The end sought in these few pages is not the study of anatomy, but the way to study it, anatomy proper being taught in the body of the book. An attempt is made to set forth a method of dissection which enables the student to explore, and prevents him from destroying before the proper time. Far too often the absence of this technical knowledge leads even a good student to dissect irregularly and without precision, mutilating structures and destroying relations.

I have not fully gone into the dissection of every organ and tissue, but have displayed as much of the subject as seems requisite to the *student* of practical anatomy.

JOHN CHALMERS DA COSTA, M.D.

2050 LOCUST STREET.

PRACTICAL HINTS ON DISSECTION.

Dissection of the Arm.

In this dissection we may include with the arm proper the superficial muscles of the back, the axilla, and the structures of the anterior thoracic region.

Fig. 1.



Anterior Portion:

Position: The body is placed on its back with several blocks under it, the arm is extended at a right angle, and placed upon a board with the flexor surface upwards.

Lines of Incision: 1. Outwards from superior border of the sternum to the top of the shoulder.

2. Downwards in middle of sternum for its entire length, starting at origin of No. 1.

3. Outwards at a right angle to the termination of No. 2, to an imaginary line dropped from the middle of the axilla.

4. From end of No. 1 down front of arm to elbow.

5. A transverse line crossing the lowest part of No. 4.

Dissection: Remove the skin from the subcutaneous tissues of the thorax and arm, and so expose the mammary gland. Be careful to leave the nipple. Observe the perforating branch of the internal mammary artery.

Remove the subcutaneous tissues of arm and chest, and the gland with them, and so expose the origin of the platysma myoides muscle, the vessels and nerves which emerge from the intercostal spaces, nerves from the cervical plexus, the cutaneous arm nerves, and the superficial veins of the elbow and arm.

Remove the fascia over the muscles of the thorax and that over the insertion of the teres major and latissimus dorsi muscles, clear the fat from the axilla and expose the vessels and nerves on the anterior surfaces of the two last-named muscles.

Cut the cephalic vein, remove the anterior portion of the deltoid fascia, note the continuation of the cephalic vein and the humeral branch of the acromio-thoracic artery.

Remove the fascia from the front of the arm, exposing important muscles, nerves, and vessels.

Cut through the great pectoral muscle in its upper third, reflect out the upper end and expose the upper portion of the coraco-brachialis and biceps muscles.

Cut the intercosto-humeral and the small internal cutaneous nerves. Reflect the other portion of the cut pectoralis major muscle, noting its blood supply and the anterior thoracic nerves.

Cut the anterior thoracic nerves. Observe the costo-coracoid membrane and the fascia of the pectoralis minor.

Remove the coverings of the subclavius and pectoralis minor muscles.

Remove the fat between the small pectoral muscle and the latissimus dorsi and teres major. Find and follow the long thoracic subscapular and anterior circumflex arteries, the subscapular and posterior thoracic nerves, and observe the subscapularis and the serratus magnus muscles, both being in part visible.

Expose at the inner side of the arm a portion of the pronator radii teres, the two heads of the biceps: the axillary, median and ulnar veins, the internal cutaneous and other nerves.

Cut the biceps transversely about its middle, expose the brachialis anticus muscle, and follow the musculo-cutaneous nerve with its branches.

Sew the biceps together. Draw the median nerve out and the basilic vein in. This exposes the brachial artery and the third portion of the axillary.

Cut the pectoralis minor muscle a couple of inches from its insertion and reflect each portion.

Expose the intercostal muscles.

Remove the basilic vein, the venæ comites of the brachial artery, and after cutting the axillary vein remove the distal part. Note the outer and inner cords of the brachial plexus, and the commencement of the median nerve. Clean the axillary artery and the outer and inner cords of the axillary portion of the brachial plexus.

Draw the brachial artery and median nerve outwards with a hook and find the branches of the brachial.

Cut the outer and inner ends of the brachial plexus; the musculo-cutaneous, the internal cutaneous and the small internal cutaneous nerves, and the axillary artery and its branches.

Reflect these structures and expose the posterior end of the brachial plexus and the posterior circumflex artery.

Tie together the arteries, veins, and nerves, and proceed to study the forearm and palm.

INCISION: 1. From termination of No. 4, in previous dissection to wrist, down front of forearm.

2. Transverse at radio-carpal articulation.

3. Transverse across palm at metacarpo-phalangeal articulation.

4. From termination of No. 1 to No. 3.

5. From No. 3 down middle of each finger to its tip.

6. Transverse incision over the head of the metacarpal bone of the thumb.

7. Join No. 2 to No. 6, and extend this incision to the end of the thumb.

DISSECTION: Remove the skin of the forearm and expose the subcutaneous veins and nerves.

Remove the subcutaneous tissue, leaving the veins and nerves.

Remove the skin of the palm and expose the palmaris brevis muscle and the palmar cutaneous nerves.

Clean the palmar fascia and expose the digital arteries and nerves.

Cut the vessels and nerves of the forearm and reflect them up and down.

Incise the forearm fascia and reflect it off. This exposes important muscles, nerves, and vessels.

Cut the bicipital fascia near the biceps tendon and remove it from the surface of the pronator radii teres muscle.

Hook aside this muscle and expose the median nerve; the brachial, radial and ulnar arteries, and other important structures.

Cut the supinator longus muscle, about its middle and reflect the proximal end. Find the radial and radial recurrent arteries, the musculo-spiral, radial and other nerves.

Replace in position the pronator radii teres and supinator longus, and clean between the distal forearm tendons.

Hook the tendons of the palmaris longus and flexor carpi ulnaris to the ulnar side of the forearm. Note the position and relation of the radial artery, radial nerve, the ulnar artery, and the median and ulnar nerves. Demonstrate portions of flexor sublimis digitorum, flexor longus pollicis and pronator quadratus muscles.

Remove the palmar fascia from over the thenar eminence, and note the vessels and muscles.

Remove the palmaris brevis muscle and the fascia over the hypothenar eminence, and note the structures exposed.

Remove the palmar fascia from the wrist towards the fingers, and clean the tendons of the flexor sublimis digitorum. Study these tendons, the anterior annular ligament and the superficial palmar arch.

Expose the vessels, nerves, and muscles in the spaces between the tendons of the flexor sublimis digitorum muscle.

Dissect the skin from the fingers, note the nerves and arteries, and clean the flexor sheaths.

Return to the forearm. Cut the flexor carpi radialis and palmaris longus muscles and median nerve, and reflect the distal portions to radial side.

Cut the pronator radii teres and reflect it with the proximal portions of the two previously incised muscles to the ulnar side. Note the origin of the flexor muscles from the inner condyle.

Cut the posterior portion of the inner part of the pronator radii teres, withdraw it from under the median nerve, reflect it inwards, and turn the distal portion of the muscle out.

Find the long origin of the flexor sublimis digitorum, and note the median nerve, the ulnar artery, and their branches.

Cut the anterior annular ligament longitudinally, note the muscular attachments to it, and cut it away.

Cut the ulnar artery, the superficialis volæ artery, the digital nerves, and the arterial branch, joining the 2d digital artery and the superficial palmar arch.

Dissect up the superficial arch and the digital nerves towards the periphery, and expose the flexor tendon sheaths and the anterior surface of the flexor sublimis digitorum muscle.

Cut the tendons of the flexor sublimis in the wrist, cut the radial origin of the muscle, and reflect the proximal part of the muscles towards the ulnar side of forearm, and its tendons towards the fingers, after we have cut off its nerve from the median.

Cut the radial artery and nerve below the elbow and above the wrist, and remove the portion between.

Cut the abductor pollicis in the middle and reflect its cut portion. Study the opponens pollicis and flexor brevis pollicis muscles, and the median nerve with its branches.

Cut the flexor brevis minimi digiti from its origin and reflect it towards the little finger, and expose the opponens minimi digiti and branches of the deep palmar nerve.

Follow the ulnar nerve and artery along the forearm and into the palm.

Clean and expose the flexor carpi ulnaris, the flexor profundus digitorum, and the flexor longus pollicis, and observe the anterior interosseous nerve.

In the palm expose the tendons of the deep flexor and the tendon of the flexor longus pollicis, and the surfaces of the four lumbricales.

Cut the lumbricales and deep flexor tendons and reflect them, leaving in place the nerves of the 3d and 4th lumbricales.

Cut the flexor longus pollicis tendon near its digital attachment.

Cut the median nerve above its digital branches and remove the branches. Expose the outer and inner heads of the flexor brevis, the adductor, and the opponens pollicis.

Cut the abductor minimi digiti and remove it with the flexor brevis minimi digiti.

Cut and remove the opponens, flexor brevis, and adductor pollicis muscles.

Show the deep palmar arch and nerve and their branches, and the attachment of the flexor carpi radialis.

Clean the palmar surface of the interosseous muscles, and finding the transverse ligament of the metacarpus, cut it away.

Cut the ulnar artery, the ulnar nerve and its branches, the median nerve and its branches, the flexor profundus digitorum longitudinally, and remove its outer part. Cut the flexor longus pollicis and pronator quadratus from the radius. Remove the proximal portions of the palmaris longus, flexor sublimis pronator radii teres, and flexor carpi radialis muscles.

Posterior Portion:

POSITION: Extend arm. If back and anterior scapular muscles have been dissected remove the arm from the trunk. Flex the forearm on the arm.

INCISION: 1. Around the arm summit of shoulder.

2. Along the spine of the scapula to the acromion.

3. Down the back of the arm to the olecranon.

4. Around the elbow.

DISSECTION OF SHOULDER AND ARM.—Remove the skin and subcutaneous tissue, observing the cutaneous nervous branches, which we cut.

Remove the fascia and expose the deltoid and triceps.

Clean the supra-spinatous muscle.

Cut the deltoid near its origin, and reflect the distal portion out. Expose the infra-spinatous and teres minor muscles. Study the triceps, and expose the teres major and a portion of the latissimus dorsi. Note the musculo-spinal nerve and the superior profunda artery.

Fig. 2.



Draw inwards the long head of the triceps.

Follow the musculo-spiral nerve through the triceps to the musculo-spiral grove.

Cut away the triceps, observe the posterior circumflex artery and the circumflex nerve.

Cut the supra-spinatous muscle, expose its proximal portion, and find the supra-scapular nerve and artery.

Section the infra-spinatous and teres minor. Cut away their scapular portions and reflect their distal portions towards the humerus. See the supra-scapular and dorsalis scapulæ arteries.

Posterior Portion of Elbow, Forearm, and Hand:

Position: In extension. Forearm supinated. Dorsum of forearm and hand looks upwards.

Incision: Corresponds exactly with the anterior plan.

Dissection: Remove the skin. Note the subcutaneous veins and nerves, and the bursa upon the olecranon.

Incise the nerves of the subcutaneous tissue, and incise the fascia in lines which correspond to the skin incisions. Remove the fascia, but retain the posterior annular ligament. Turn the radial side of the hand and arm up, and place the hand midway between pronation and supination. Study the muscles, arteries, and tendons at its outer side.

Cut the extensor communis digitorum about the middle of the arm, and the extensor radialis longior and extensor minimi digiti lower down, and reflect them, removing the distal portions from the posterior annular ligament, and in freeing their prominal portions, cutting a branch of the posterior interosseous nerve. We can now study the posterior interosseous nerve and artery, and the extensor carpi radialis brevior muscle.

Cut away the branches of the posterior interosseous nerve and artery, and expose the extensor indicis and extensor secundi internodii pollicis muscles, opening the latter's space in the annular ligament.

Cut the extensor indicis above the wrist, the extensor secundi a little higher up, and the extensor carpi radialis brevior near its middle, and reflect them. Study the posterior interosseous nerve, the perforating branches of the anterior interosseous artery, and expose the supinator brevis, the extensor ossi metacarpi pollicis, and extensor primi internodii pollicis muscles. Clean the deep fascia upon the carpus and metacarpus.

Cut the posterior interosseous nerve above and below, and remove it.

Cut the middle perforating branch of the anterior interosseous artery.

Cut the extensor ossi metacarpi pollicis and the extensor primi internodii pollicis muscles, and reflect them.

Cut the ulnar attachments of the extensor indicis and extensor secundi internodii pollicis, the tendons of the extensor carpi radialis longior and brevior, and the extensor carpi ulnaris. Expose the radial attachment of the pronator radii teres, and remove the portion of the supinator brevis which is attached to the ulna.

Remove the deep fascia from the carpus and metacarpus, and from over the lower ends of the radius and ulna.

Study the posterior carpal branches of the radial and ulnar arteries, the dorsal arteries of the fingers, the perforating arteries, the dorsal interossei muscles and other structures.

Remove the skin of the fingers but not the vessels and nerves. Clean the tendons of the extensor communis digitorum, extensor indicis, and extensor minimi digiti. Note the attachments of the lumbricales, the palmar and dorsal interossei.

Note upon the dorsum of the fingers, from the 2d to the 5th inclusive, the aponeurosis formed by the extensors, the lumbricales and the palmar and dorsal interossei.

Anterior and Lateral Wall of Abdomen.

Position: Body on its back, with a block placed under the shoulders and another under the pelvis. The penis is hooked forward.

Fig. 3.



Incisions: 1. Transverse between the two anterior superior iliac spines.

2. From centre of No. 1 to root of penis, and along the dorsum of the penis to its head.

3. Continue the sternal incision (see anterior thoracic region) to the middle of No. 1.

Dissection: Reflect the skin outwards and downwards from junction of 1 and 2, and laterally from the penis.

Note the two layers of subcutaneous tissue and the vessels between them, and observe the superficial nerves.

Remove the superficial layer to beyond the saphenous opening.

Cut the superficial vessels and nerves of the dorsum of the penis and dissect them from the dartos fascia.

Expose the suspensory ligament of the penis.

Incise the dartos sheath and reflect it externally.

Incise and remove the elastic sheath of the penis, exposing the deep vessels and nerves.

Incise the deep layer of the superficial abdominal fascia, and, with the vessels, reflect it out.

Remove the superficial fascia from the external oblique muscle, preserving that portion at the pillars of the ring which goes to invest the spermatic cord.

Incise this fascia and reflect it from the cord.

Cut the aponeurosis of the external oblique by incisions corresponding to No. 1 and lower abdominal portion of 2. Reflect the flap upon the thigh.

Incise the lateral portion of the external oblique from the 11th rib to the ilium, and run the incision forward over the iliac crest. Dissect the flap loose at its superior lateral origin and reflect the muscle downwards and inwards to beyond the linea semilunaris. Note the septum between the two oblique muscles, and the ilio-hypogastric and ilio-inguinal nerves.

Remove this intermuscular septum.

Dissect the two above-mentioned nerves and reflect them externally.

Make a transverse incision through the internal oblique from the anterior superior spine of the ilium to the linea semilunaris.

Cut the fibres attached to Poupart's ligament. Reflect the flap to the semilunar line.

Cut the muscle from the ilium to the 11th rib, along the crest, and below the 11th rib and its cartilage. Reflect the flap inwards. Note the vessels and nerves between the internal oblique and transversalis muscles.

Remove the internal oblique up to the conjoined tendon and cut away the cremaster from the cord.

Remove the fibrous tissue from the transversalis. Study this muscle, the deep vessels and nerves, and the conjoined tendon.

Replace in position the external and internal oblique.

Make an incision between the linea alba and the semilunar line. Reflect the flaps and expose the rectus and pyramidalis muscles. Remove the pyramidalis muscle. Note the nerves running into the rectus, the deep epigastric artery, and the branch from the internal mammary artery.

Retain these arteries by removing the muscle between them.

Note the spermatic cord (in the female, the round ligament).

Remove the external and internal oblique up to the semilunar line.

Cut away the lateral sheath of the rectus between the semilunar line and the linea alba.

Cut the circumflex iliac arteries at the points of muscular perforation and the intercostal nerves and vessels at their points of emergence.

Cut through the transversalis muscle along the lower borders of the cartilages of the three lower ribs, and the lower border of the 11th rib, from the 11th rib to ilium, along the crest of the ilium, and through the fibres of origin from Poupart's ligament. Cut away at the linea semilunaris.

Note the transversalis fascia and internal abdominal ring.

Incise this fascia from the superior iliac spine to the semilunar line. Reflect a flap and expose the interior of the internal abdominal ring. Note the infundibuliform process of the transversalis fascia (internal spermatic fascia), and the deep epigastric artery.

Incise the infundibuliform fascia and expose the cord. Study the cord and trace it through the ring.

Anterior of Thigh.

POSITION: Body supine, legs extended, a block under the pelvis.

INCISIONS: 1. From middle of Poupart's ligament down front of thigh to below the lower margin of the patella.

2. Transverse across the knee at the termination of 1.

DISSECTION: Remove the skin.

Remove the superficial layer of the subcutaneous tissue and note the veins and nerves.

Remove the deep layer. Notice the lymphatic glands, prepatellar bursa, and saphenous vein.

After removing the lymphatic glands cut the branches of the internal saphenous vein. Trace the cutaneous nerves.

Clean the fascia lata, keeping its vessels and nerves intact.

Study the saphenous opening, cribriform fascia, internal saphenous vein, and the crural branch of the genito-crural nerve.

Cut the saphenous vein a little below the opening and the nerves close to the points at which they perforate the fascia lata. Remove the nerves and vessels and incise the fascia, the incisions corresponding to those in the skin. This exposes Scarpa's triangle and important muscles.

Clean the pectineus muscle. Find the femoral vein by means of the saphenous. Open the sheath of the vessel and nerve in the triangle. Clean the iliacus muscle. Find the anterior crural nerve by following up the remains of the internal and middle cutaneous.

Fig. 4.



Cut the sartorius across at its middle, reflect each portion, find the roof of Hunter's canal and the long saphenous nerve.

Cut the tensor vaginæ femoris muscle where it is attached to the fascia and reflect it, noting the nervous branch.

Study the rectus, gluteus medius and minimus muscles.

Cut the rectus through the middle and reflect each half. In its proximal part, note the branch from the anterior crural nerve and the external circumflex artery.

Open Hunter's canal and study its contents.

Expose the deep femoral artery and vein at the superior border of the adductor longus. Clean the vasti muscles and the crureus, and remove the venæ comites.

Cut through the quadriceps extensor by a curved incision below the patella (don't open the joint).

Make a long incision down to the bone through the vastus externus.

Abduct the thigh and partly flex the knee.

Reflect the aponeurosis from the inner side of the knee-joint from below upwards and outwards, cutting the ligamentum patella. Reflect the vastus internus, inner part of vastus externus and crureus. Remove them from the femur. Cut the branches of the anterior crural nerve and of the external circumflex artery. Remove the vastus internus from the linea aspera.

Cut the femoral artery and vein below the origin of their deep branches, and also at the opening in the adductor magnus.

Cut the long adductor high up, reflect its proximal portion and note the branches of the obturator nerve beneath it.

Reflect the distal portion and cut off close to the bone.

Study the deep femoral artery and other branches. Remove the venæ comites. Turn the femoral vein over Poupart's ligament. Make a long incision through the pectineus, and remove the proximal portion from its origin. Bring back the femoral vein. Find the obturator nerve and artery. Clean the long and great adductors. Note the femoral opening in the great adductor, and find the anastomotica magna artery. Cut the last portion of the deep femoral artery and its perforating branches, and the anterior crural nerve, femoral artery, and femoral vein close to Poupart's ligament. Remove the deep femoral artery and the remaining portions of the femoral artery and anterior crural nerve. Cut the internal circumflex artery near its origin, and cut the obturator nerve and its branches and remove them. Cut and reflect the short adductor, and cut it distally from the linea aspera.

Reflect the distal portion of the pectineus and cut it from the femur. Cut the obturator artery. Clean anteriorly the great adductor, portions of the external obturator, quadratus femoris, semimembranosus, iliacus internus, psoas magnus, and part of the hip-joint capsule.

Cut the iliacus internus and psoas magnus near their insertions, reflect them upwards and remove and observe the bursa under them.

Clean the external obturator muscle and note the proximal portions of the obturator nerve and vessels.

Cut the origin of the rectus and clean more of the capsule of the hip-joint.

Anterior of Leg and Upper Surface of the Foot.

POSITION : Legs extended. Blocks under the ankles. Great toes tied together.

INCISIONS : 1. From centre of incision made below the patella in the last dissection down the middle of the front of leg and ankle and along the dorsum of the foot, terminating at the root of the 2d toe.

2. Transverse along the metatarso-phalangeal articulations.
3. From 2 draw a line along the dorsum of each toe to its end.
4. From each malleolus to the heel.

DISSECTION : Reflect the skin from the dorsal and lateral surfaces of the foot and toes and from the leg. Note subcutaneous veins and nerves.

Dissect out the saphenous arch and its veins, and expose the superficial nerves. Clean the fascia of the toes and foot, retaining the veins and nerves.

Remove these veins and nerves from the foot. Cut the fascia in the long axes of the tendons, note the tendon synovial sheaths, remove the fascia, and find the annular ligament.

Remove the nerves and veins from the leg fascia, cut the fascia longitudinally a little external to the tibia, and reflect externally to the fibula. Retain the anterior annular ligament, and the fascia over the upper part of the tibialis anticus muscle. Study the exposed muscles and the anterior annular ligament.

Expose a tendon of the extensor brevis digitorum, the dorsalis pedis artery and veins, and the anterior tibial nerve.

Cut the fascia of the peroneus longus and brevis muscles, and remove it, and trace the tendons of these muscles.

Cut the tendons of the extensor longus digitorum and peroneus tertius muscles below the annular ligament, and reflect their distal portions. Clean the extensor brevis digitorum muscle.

Cut the tendon of the extensor proprius pollicis below the lower edge of the anterior annular ligament. Reflect the proximal portions of the extensor longus digitorum, peroneus tertius, extensor proprius pollicis, and tibialis anticus muscles. Draw the first three outwards and the other inwards. Note the anterior tibial vessels and nerve.

Cut the extensor brevis digitorum muscle above its tendons, reflect its proximal part and remove it. Reflect its tendons to the beginning of the toes and cut them off. Trace the dorsalis pedis and other arteries.

Remove the origins upon the fibula of the extensor longus digitorum, peroneus tertius, and extensor proprius pollicis muscles. Preserve the intermuscular septum and the fascial attachment. Trace the musculo-cutaneous nerve. Cut the peroneus longus muscle at the head of the fibula. Trace the anterior tibial nerve to its origin. Trace the anterior tibial artery upward, and trace the anterior peroneal artery.

Sole of the Foot.

POSITION: Raise the foot by several blocks and steady it by hooks.

INCISIONS: 1. Transverse across the metatarso-phalangeal articulation.

2. From No. 1 down the middle of each toe to the end.

3. From the middle of No. 1 backwards to the posterior portion of the os calcis.

DISSECTION: Remove the skin and cut it away.

Remove the subcutaneous tissue, retaining the vessels and nerves. Note the plantar fascia, plantar collateral digital vessels and nerves and digital tendon sheaths, and the terminal portion of the internal plantar artery.

Make a transverse incision through the plantar fascia at its middle, reflect its distal portion, and with it the superficial transverse ligament. Preserve the nerves and arteries between the fascial slips. Reflect its proximal portion, but do not remove it from the flexor brevis digitorum muscle. Note the branches of the internal plantar nerve and artery.

Open the flexor tendon sheaths, and clean the flexor brevis digitorum muscle.

Remove the inner portion of the plantar fascia, trace the internal plantar artery, and clean the abductor pollicis muscle.

Remove the outer portion of the plantar fascia, trace the first digital artery and 5th and 6th digital nerves. Clean the abductor minimi digiti muscle.

Remove the flexor tendon sheaths, expose the tendon of the flexor longus pollicis muscle and a portion of the tendons of the flexor longus digitorum muscle. Clean the accessible portions of the flexor brevis pollicis, flexor brevis minimi digiti, and lumbricales muscles. Note the digital arteries.

Cut proximally the 1st, 5th, and 6th digital nerves and the internal plantar nerve, the nerves to the flexor brevis pollicis muscle and to two of the lumbricales. Reflect the digital nerves from the toes. Cut the anastomotic, internal plantar, and first digital arteries, and reflect the two last over the toes. Reflect the branches of the anastomotic artery, and cut them away. Cut the flexor brevis digitorum, abductor pollicis and abductor minimi digiti muscles, and remove their peripheral portions. Reflect and remove the proximal portions of these muscles. Trace the internal and external plantar arteries and the internal and external plantar nerves.

Clear the tendons and expose the lumbricales, the accessorius, and the flexor brevis minimi digiti muscles, and portions of other muscles.

Cut the internal plantar artery and nerve, and remove their distal portions. Cut the 1st tendon of the flexor longus digitorum muscle, the flexor accessorius muscle, and the tendon of the flexor longus pollicis muscle. Reflect the distal portion of the tendons and the lumbricales muscles over the digits. Note nerves, arteries, and a tendon of junction.

Find the tarsal grooves for flexor tendons, and observe the tendon of the tibialis posticus muscle.

Clean the transversus pedis muscle and expose the abductor pollicis and the flexor brevis pollicis muscles. Observe the digital arteries.

Cut the nerve of the flexor brevis minimi digiti muscle, and cut this muscle from its origin, reflect it to its insertion and remove it. Cut the metatarsal attachments of the transversus pedis muscle, cut this muscle's nerve, and reflect the muscle over the great toe. Cut the digital arteries. Cut off the abductor pollicis muscle proximally and reflect it upon the great toe.

Cut the proximal portion of the flexor brevis pollicis and reflect the muscle distally, and cut it through between its two heads. Reflect the 5th and 6th digital arteries upon the tarsus. Observe the compound tendon of the flexor brevis pollicis, the sesamoid bones in this tendon, and the ligament between these bones and the 1st metatarsal. Cut this ligament and the muscles near the bones, and reflect the bones upon the base of the great toe. See the deep branch of the external plantar nerve.

Trace the plantar arch and return into position the 5th and 6th digital arteries.

Clean the plantar and dorsal interossei.

Cut away the sesamoid bones. Bring into place the tendons of the flexor longus pollicis and flexor longus digitorum muscles, and sew their divided ends. Extend the foot. Note the attachment of the lumbricales.

Now open the dorsal surface, dissect the tendons of the extensor longus and brevis digitorum muscles, which we left in the dissection of the dorsum of the foot.

Posterior Portion of Leg.

POSITION: Limb extended with posterior surface upwards. Blocks under pelvis, thigh, and ankle. Foot at right angles to leg.

INCISIONS: 1. Transverse above the middle of the popliteal space.

2. From the middle of 1 down the middle of posterior surface of leg to os calcis.

3. Along the inferior edge of heel.

DISSECTION: Remove the skin and note the veins and nerves.

Cut the external saphenous vein as it enters the fascia and the roots of the external saphenous nerve close to the fascia.

Remove the subcutaneous tissue and with it the veins and nerves.

Cut the fascia as we did the skin and reflect it. Note the nerves and veins. Clean the gastrocnemius muscle, retaining the nerves and the proximal portion of external saphenous vein.

Cut the small sciatic nerve and remove its distal part. Clean the boundaries of the popliteal space. Note the external popliteal nerve.

Cut the cutaneous branch of the external popliteal nerve, the external saphenous vein and nerve, the internal saphenous vein and nerve, and remove their distal portions. Partly flex the knee joint and study the popliteal space and its contents.

Draw aside the popliteal nerve and dissect out the popliteal vein.

Cut off the branches of the popliteal vien, and draw the vein aside to expose the artery.

Fig. 5.



Clean the soleus and other muscles.

Expose the tendons, vessels, and nerves of the distal portion of the leg.

Cut through transversely the gastrocnemius muscle. Reflect it upwards, and cut away the middle of the muscle. Clean the plantaris and soleus muscles. Note the nerves and vessels.

Cut the plantaris muscle below the point its vessels and nerves enter.

Cut the inner nerve to the soleus, reflect the tendon of the plantaris to the os calcis and cut it away.

Cut the tendo Achillis and reflect the soleus upwards to its origin, and cut it away.

Clean the popliteus muscle. Trace the internal popliteal nerve, the popliteal vein and the popliteal artery peripherally.

Remove the fascia from between the muscles.

Extend the foot. Clean the flexor longus pollicis and the flexor longus digitorum muscles and study them.

Draw the posterior tibial nerve inwards, remove the venæ comites of the posterior tibial and peroneal arteries. Study the tibial and peroneal arteries.

Clean the peroneus longus and brevis muscles.

Reflect the proximal portions of the flexor longus pollicis and the flexor longus digitorum muscles, and cut them off from their bony origins.

Cut the posterior tibial nerve, and also the posterior tibial artery below the origin of the peroneal artery. Trace and study the peroneal artery.

Clean and study the tibialis posticus muscle.

Gluteal and Posterior Thigh Regions.

Position : Body on face. Block under the pelvis anteriorly. Leg extended and thigh rotated inwards.

Incisions : 1. Along the crest of the ilium, and down between the buttocks.

2. From the middle of 1 down the back of the leg to above the middle of the popliteal space, where we meet the transverse incision of the previous dissection.

Dissection : Remove the skin.

Remove the subcutaneous tissue retaining the veins and nerves.

Abduct the leg. Incise the fascia lata of the gluteal region, and remove it from the gluteus maximus muscle.

Incise the fascia lata, the line corresponding to the skin incisions, reflect on each side, and retain the small sciatic nerve.

Replace the limb in its old position. Cut the small sciatic nerve and remove its peripheral portion. Study many muscles.

Cut the gluteus maximus muscle from its superior to its inferior border, through its middle, and reflect each half. See bursa, vessels and nerves. Cut away this muscle externally.

Abduct the thigh again. Expose the great sacro-sciatic ligament and various muscles, nerves, and vessels.

Remove the internal portion of the great gluteal muscle, and observe another bursa, and many important structures, ligamentous, muscular and vascular.

Cut the inferior gluteal and small sciatic nerves, and the sciatic artery, high up and remove their distal portions. Cut the origin of the articular branch and leave it. Separate from the semitendinosus muscle the long head of the biceps.

Flex the leg a little. Draw the long head of the biceps externally, and the semimembranosus and semitendinosus internally. Study the sciatic nerve.

Cut the tendons of origin of the biceps and semitendinosus muscles and reflect the muscles. Cut the semitendinosus again near its point of attachment, and the long head of the biceps again near its junction with the short head. Clean the semimembranosus and the short head of the biceps.

Cut the semimembranosus muscle and great sciatic nerve high up, and reflect their distal portions. Cut the semimembranosus near its insertion, and cut the nervous branch to the great adductor.

Cut the perforating arteries and the popliteal artery with its branches, and remove the arteries.

Clean the adductor magnus muscle. Find the femoral opening, the anastomotica magna artery and the long saphenous nerve.

Find the articular branches of the popliteal artery.

Clean the quadratus femoris muscle and note the internal circumflex artery.

Cut the gluteus medius muscle and reflect each half, and cut it away. Note the branches of the gluteal artery and the superior gluteal nerve. Study the gluteus minimus muscle.

Find between the great sacro-sciatic ligament and the inner ends of the pyriformis muscle, the internal pudic vessels and nerve.

Clean the gemellus superior and inferior and the obturator internus muscles.

Expose the trunk of the great sciatic nerve.

Section and remove the quadratus femoris muscle. Cut the nerve, clean the posterior surface of the external obturator artery. Find the internal circumflex artery.

Cut the tendons of the obturator internus and the gemelli. Reflect the muscles. Note bursa.

Cut the external obturator muscle and reflect it outwards.

Cut the gluteal artery and the superior gluteal nerve. Cut the gluteus minimus muscle and remove it. Cut the pyriformis, reflect it and remove it. Study the contents of the two sacro-sciatic foramina.

The Perineum of the Male.

Position : In the lithotomy position. Tie the knees by a rope around the body. Stuff the rectum with rags. Pass a bougie and tie the penis and scrotum to it.

Fig. 6.



Incisions : 1. From the posterior part of the scrotum to the coccyx, going around the anus.

2. A transverse incision back of the scrotum.

3. A transverse incision back of the anus.

Dissection : Remove the skin alone, to beyond the margins of the muscles of the thigh.

Clean and expose the sphincter ani muscle.

Incise the subcutaneous tissues to the perineal fascia, the lines of incision corresponding to those of the skin, and keeping clear of the margins of the sphincter.

Remove the fat external to the anterior half of the sphincter, and note the posterior edge of the perineal fascia.

Clean out the ischio-rectal fat and expose the lower edge of the gluteus maximus muscle. Note the inferior hemorrhoidal artery and nerve, the obturator fascia, and the anal fascia.

Open the obturator fascia and expose the pudic vessels and nerves.

Cut the perineal fascia by a transverse incision running out from in front of the central tendon half way to the fascia lata, and from the external portion of this incision run another forward to the posterior portion of the scrotum, a little external to the middle line on both sides. Reflect the fascia. Note the superficial perineal artery and nerves, the superficial transverse perineal muscles, the accelerator urinæ and erector penis muscles, and the central tendon of the perineum.

Cut and reflect forward the inferior pudendal nerve, and the superficial perineal artery and nerves. Cut the three abovementioned muscles, and reflect them, being careful not to remove the central tendon.

Note the crus penis and the corpus spongiosum.

Draw the bulb by a hook inwards and clean the triangular ligament.

Cut one of the crura of the penis from its origin and draw it internally, and then clean the nerves and vessels which perforate the triangular ligament.

Incise the ligament and reflect it laterally, being sure not to remove it posteriorly from the fascial junction. Note the artery of the bulb and the deep perineal artery and nerve.

Cut the deep perineal artery, reflect it internally, and clean the underlying muscles. Note the constrictor urethræ muscle and the deep transverse perineal muscle.

Restore the crus to position. Draw the bulb a little anterior.

Expose Cowper's glands by removing fibres of the deep transverse muscle.

Cut the great gluteal muscle and reflect it inferiorly.

Cut the constrictor muscles and deep transverse muscles longitudinally, reflect them out and in, and remove the outer portions.

Note the great sacro-sciatic ligament, the levator fascia, and the 4th sacral nerve, and study the membranous urethra.

Cut away the levator fascia from the levator ani et prostatæ muscle. Study this muscle and the coccygeus muscle.

Cut this muscle on each side by a semicircular sweep well external, and remove the internal portions. Note the recto-vesical fascia.

Hook the bulb (and hence the prostate and bladder) well front. Dissect the rectum from the recto-vesical fascia and base of the bladder, drawing it well out. Remove from the prostate the vesical layer of the recto-vesical fascia, and reflect it posteriorly.

Note the vasa deferentia, the venous plexus, and the seminal vesicles, the prostate, the recto-vesical fold of peritoneum, and the middle hemorrhoidal arteries. Now return the viscera, and hold them in by sewing the skin.

The Perineum of the Female.

POSITION: Same as in the male.

INCISIONS: Plug the vagina and rectum. Sew together the labia majora back of the clitoris and sew up the anus.

FIG. 7.



1. Make an incision outward, on each side, from the superior lateral aspect of the labia majora to the thigh.
2. Carry an incision around each labial base to the centre of the perineum.
3. An incision back from this point to just in front of the anus.
4. An incision back around the anus.
5. An incision from just back of the anus to the coccyx.
6. An incision is then carried from one ischial tuberosity to the other, touching the termination of No. 5.

DISSECTION: Reflect the skin out on each side.

Clean the sphincter ani muscle, expose some nerves, and the superficial artery of the perineum.

Clean the perineal fascia, remove the fat as in the male, to expose the posterior edge of this fascia.

Cut the nerves and the superficial artery well back and reflect them front, and remove the balance of the ischio-rectal fat. Note structures as in the male.

Cut and reflect the perineal fascia. Note the sphincter vaginæ, erector clitoridis, and superficial transverse muscles, also the central tendon.

Cut the superficial transverse muscle and reflect it outwards. Cut each half of the sphincter vaginæ muscle, and reflect it anteriorly. Cut the labial sutures, remove the vestibule mucous coat, remove the small labia from the large labia and clitoris, and remove the anterior one-half of the labia majora. Note the vaginal bulb.

Detach one crus of the clitoris from its origin, and draw the bulb and the cut crus over the median line. Note the triangular ligament, the artery of the bulb, and other structures.

Restore the vaginal sphincter, the crus, and the bulb into place.

Make traction upon the clitoris, incise the skin of the dorsum of this organ and the mons Veneris and reflect it. Note the superficial vessels and nerves. Remove these vessels and nerves from the clitoris. Note the anterior portion of the vaginal sphincter, cut it away, and observe the sheath and suspensory ligament of the clitoris. Cut through the sheath longitudinally and reflect it laterally on each side, and observe the deep vessels and nerves of the dorsum of the clitoris.

Cut the suspensory ligament with its vessels and nerves, remove the clitoris and vaginal bulb, and cut the clitoris transversely. Cut the triangular ligament as we did the perineal fascia. Reflect it partly backwards, partly outwards, and partly inwards. Note the artery of the vaginal bulb and the deep perineal artery and nerve. Clean the muscles of the urethrovaginal region. Note the constrictor urethræ and deep transverse muscles.

Cut away a part of this last-named muscle to expose the vulvo-vaginal gland.

Cut and reflect the great gluteal muscle as in the male. Note the 4th sacral nerve and the great sacro-sciatic ligament.

Cut the deep transverse and the urethral constrictor muscles, and reflect inwards. Preserve the central tendon, and the attached borders of the perineal fascia and of the triangular ligament. Note the levator fascia.

Remove this fascia and the fascial junction. Note the beginning of the obturator fascia. Cut the deep perineal artery and nerve and remove the tendinous centre. Note the levator ani and vaginal muscles. Incise the levator ani as we did in the male. Treat the rectal fibres as in the male, reflect the inner portion to the median line, and remove.

Note the coccygeus muscle and the recto-vesical fascia.

Upon the rectum and vagina incise this fascia and reflect it outwards.

Remove the rectal and vaginal plugs. Draw the urethra anteriorly. Separate the anterior superior and the anterior inferior vaginal walls, draw the vagina back, and find the base of the bladder. Note the ureters.

Restore the bladder, urethra, and vaginal wall to their places. Hook the posterior vaginal wall to the front and dissect the rectum from it, pulling the rectum posteriorly. Find Douglas's cul-de-sac, and note the vaginal and middle hemorrhoidal arteries. Push back the perineal contents and hold them in by sewing the skin.

Abdominal Cavity and Pelvic Cavity.

POSITION: The same as for dissecting the abdominal muscles.

DISSECTION: Remove the sheath of the rectus (if still in place), and remove the lower two-thirds of the transversalis fascia. Between this fascia and the peritoneum retain the remains of the fœtal duct and vessels.

Cut the parietal layer of the greater peritoneum a little to the left of the median line from the level of the xiphoid cartilage to a level with the umbilicus, and from the termination of this incision carry another incision outwards. Note the cavity of the greater peritoneum by the hand, feeling the stomach and other parts, finding the great omentum and the small intestine under it.

Cut the round ligament and broad ligament of the liver from the abdominal parietes. Make a transverse cut through the peritoneum on the right of the median line, and inferiorly to the left of the median line, and reflect the flaps. Make a transverse cut through the peritoneum between the stomach and the transverse colon. Push the hand into the cavity of the lesser peritoneum. Feel the foramen of Winslow, the pancreas, and other parts.

Draw the great omentum up with the transverse colon. Note the small intestine and the transverse meso-colon.

Find the ileum and jejunum, draw them forward and to the left and note the mesentery.

Cut the right side of the mesenteric attachment and reflect the mesentery to the left. Note the superior mesenteric vein. Cut off this vein, and remove it and its branches. Tie a ligature around the small intestine and another one a foot off. Inflate with a blowpipe. Note the branches of the superior mesenteric artery and the intestinal branches of the sympathetic nerve.

Incise and dissect the posterior layer of the transverse meso-colon. Note the colica media, colica dextra, and ileo-colic arteries.

Tie two strings around the ileum a few inches from the colon and cut between. Remove the ileum and jejunum up to the duodenum (cutting close to the bowel with scissors), throw two cords around the bowel and cut between. Cut the superior mesenteric artery and remove its branches.

Remove the great omentum from the colon and from the greater curvature of the stomach. Remove from the colon the anterior layer of the transverse meso-colon. Spread out the large intestine and expose the greater curvature of the stomach. Follow the inferior mesenteric artery (dissecting the peritoneum from the left side posteriorly, and reflecting the anterior layer of the sigmoid meso-colon).

From the anterior surface of the pancreas remove the posterior layer of the lesser peritoneum, removing it also from a portion of the duodenum. Draw up the left hepatic lobe. Open the gastro-hepatic omentum through its anterior layer, and remove it from the lesser curvature of the stomach, and the inferior surface of the liver posteriorly. Note the cœliac axis, the gastric, splenic, and hepatic arteries.

Remove the peritoneum from the presenting surface of the gall-bladder. Note the cystic duct and the ductus communis choledochus. Lift up the branches of the hepatic artery and expose the portal vein. Throw a string around the cardiac end of the stomach, tie it, and distend the viscus by means of a blowpipe.

Pull the right side half of the stomach up and out, and dissect the duodenum from the head of the pancreas (the ascending and part of the descending duodenum). Tie a long string around the pylorus and draw the stomach and loosened part of the duodenum up. Note the gastro-duodenal artery and the ductus communis choledochus. Pull out the lower cartilages of the ribs of the left side, clean and study the pancreas, and observe the superior mesenteric vessels, the spleen, and the splenic vessels.

Remove the anterior fold of the gastro-splenic omentum, and after turning the spleen outwards note the branches of the splenic artery. Draw the upper border of the pancreas forward to study the splenic vein.

Replace the stomach and duodenum. The sigmoid flexure must have been cut. Hook up the portion below the cut to put the rectum on the stretch. Study the pelvic peritoneum, the rectum, and bladder, and in a woman the round and broad ligaments, the uterus, the Fallopian tubes, and the ovaries in position.

Draw down the stomach, cut above the string, and draw the stomach and spleen downwards and to the right. Loosen the pancreas and duodenum posteriorly. Replace all of these parts. Trace the inferior vena cava, ligate it above the renal veins, and cut it through.

Draw the liver down and cut the ligaments near the diaphragm. Feel with the finger the vena cava entering the diaphragm, and after ligating this vessel cut it off below the string. Cut the remaining part of the posterior inferior layer of the coronary ligament. Cut the cœliac axis and the stump of the superior mesenteric artery. Take away together the pancreas, liver, stomach, and duodenum. Remove the peritoneal covering of the aorta and of the iliac arteries. Study these vessels and the sympathetic plexuses upon them.

Clean the supra-renal capsules and the vena cava. Note the renal veins and expose, by drawing these veins front, the renal arteries.

Remove the kidney fat and study the kidneys and ureters.

Note the inferior mesenteric artery, in the male the spermatic veins and arteries, and in the female the ovarian.

Cut the vena cava above the iliac veins, the aorta above the inferior mesenteric artery and the ureters on a level with the crest of the ilium, and reflect their lower portions into the cavity of the pelvis. Remove the testicles from the scrotum and place them in the pelvis, cut the sutures in the perineum, and cut the penis loose from the arch of the pubes.

In the male trace the ureters and the vasa deferentia, in the female the ureters, round ligaments and ovarian vessels. Cut these parts externally and reflect them inwards. Tear the organs loose mostly with the fingers, cutting, however, the vessels. Remove the viscera when loose through the pelvic outlet.

Clean and study the iliac arteries and veins. Remove the venæ comites of the anterior trunk branches of the internal iliac arteries. Note the obturator, superior vesical, and other arteries.

Cut the iliac, pudic, sciatic, and obturator arteries in two places and remove them. Cut and take away the iliac veins. Clean the branches of the posterior trunk of the internal iliac and the sacra media artery.

Cut the lateral sacral, gluteal, and ilio-lumbar arteries. Take away the common iliac artery above and below its bifurcation. Expose and study the lumbar ganglia of the sympathetic and portions of the lumbar arteries.

Clean and expose the psoas magnus and parvus, quadratus lumborum, posterior portion of transversalis, and iliacus internus muscles (don't cut away their nerves).

Cut the psoas magnus and parvus muscles of one side through the middle, and reflect each part. Note the lumbar plexus. Observe the ilio-hypogastric and ilio-inguinal, the external cutaneous, the anterior crural, and other nerves.

Find the 5th lumbar nerve and its branch from the 4th, the sacral nerves, and the sacral ganglia, and the ganglia impar.

Section the aorta below the cœliac axis and remove to the cut made above the inferior mesenteric artery. The ureters, the vena cava, the spermatic or ovarian vessels must be cut. The branches of the suprarenal arteries are cut from the phrenic arteries.

Find the receptaculum chyli on the 2d lumbar vertebra.

Remove the kidneys and capsules with the upper portions of these sectioned structures. Note lymphatic glands around the receptaculum, and observe the beginning of the great azygos vein.

Remove the diaphragmatic peritoneum and study the phrenic arteries and nerves. Find the openings in the diaphragm, and clean the ligamenta arcuata and the crura. Study a number of muscles and ligaments.

Abdominal Viscera.

Slit up and study the interior of the ilium and jejunum, washing it clean.

Take a small section of intestine, tack it to a board to hold it firm and stretch it and dissect the coats.

Cut the colon six or eight inches above the ilio-cæcal valve, study this region, note the valve action by pouring water in each end, open the colon externally, and study the appendix.

Slit open the colon, wash it and study.

Cut the duodenum off just below the stomach and separate the stomach and spleen from the pancreas, liver, and duodenum.

Put the liver on its anterior surface, its inferior border looking up, and turn the duodenum and pancreas on to their posterior surfaces below the superior hepatic border. Cut away the front of the 2d portion of the duodenum. Open the common duct and find its duodenal opening with a probe. Follow the splenic and superior mesenteric veins to the portal.

Cut the portal vein near its origin, and the common duct on the same line. Remove the liver, and take off of it the splenic and superior mesenteric veins. Study the surface of the liver, its vessels, ducts, etc. Free the gall bladder. Cut the cystic duct near the hepatic and the cystic artery. Open the bladder.

Section the liver.

Study the pancreas. Make a longitudinal incision in the upper half of the posterior surface of the pancreas and find the pancreatic duct.

Separate the stomach and spleen. Cut the stomach along its lesser curvature from the cardiac orifice nearly, but not quite to the pyloric orifice, invert it and study.

Remove the duodenum from the pancreas, slit the duodenum entirely open and study.

Study the spleen. Cut to find the vessels.

Study the kidneys and the suprarenal capsules. Peel off the kidney capsule to the hilus and trim it away. Section the kidney.

Pelvic Viscera of the Male.

The rectum and bladder having been removed together and stuffed (a small opening is made in the bladder anteriorly to admit of stuffing) are cleaned, the dangling nerves and arteries being retained. Note the peritoneal coverings of the bladder and rectum.

Cut the recto-vesical fold of peritoneum and remove the rectum from the bladder. Cut the vasa deferentia and ureters, and, after opening the bladder, spread it out and tack it to a board with the penis, the base of the bladder and inferior surface of the penis looking upwards. Note the seminal vesicles, vasa deferentia, ureters, the prostate and Cowper's glands, the bulb of the corpus spongiosum, and the crura of the penis.

Make a cut through the prostate in its middle line from base to apex, in order to find the ejaculatory ducts.

Take the bladder and penis from the board, turn them over and tack them down again, exposing the interior of the bladder and the dorsum of the penis. Make a longitudinal incision in the dorsum of the penis, a little out from the median line, extending from the bladder to the meatus, and opening the urethra. Study the interior of the bladder, the penis, the urethra, the prostate gland, etc.

Take the removed testicle with the portion of cord still attached to it, and pin them on a board. Reflect the sheath from the cord, and reflect the tunica vaginalis. Find the testis, the epididymis, the vas deferens, the vessels, and nerves.

Pelvic Viscera of the Female.

Remove these viscera together, stuff the rectum, bladder, and vagina.

Remove connective tissue, but retain peritoneum. Lay the organs on their lateral surfaces upon a board, maintaining their normal position of mutual relationship. Note the organs and the reflections of peritoneum.

Cut the utero-vesical fold of peritoneum, and dissect the bladder and urethra from the vagina and uterus. Cut the recto-vaginal fold and separate the rectum and vagina. Open the urethra and bladder on their anterior superior surface, and, spreading them out on a board, study their interiors.

Take the plug out of the vagina, and nail the uterus, broad ligaments, and vagina upon a board, with their anterior superior surfaces looking upwards.

Reflect the anterior layer of the right broad ligament, and note the round ligament, the Fallopian tube, the uterine and ovarian vessels, and the ovarian ligament.

Turn the organs over, thus exposing their posterior inferior surfaces. Note the uterine peritoneum, the ovary, the uterus, and vagina.

Cut the Fallopian tubes and uterine ligaments, open the uterus posteriorly from the fundus to the os uteri, find the uterine openings of the Fallopian tubes, and open them laterally, spread the uterus open and tack it. Study its interior.

Clean the rectum externally, retaining its vessels and peritoneum. Remove the plug, open the canal, and study its interior.

Anterior Portion of the Thoracic Wall and the Thoracic Cavity.

"Anterior Thoracic Region" has been dissected.

DISSECTION: Remove the great pectoral muscle from its sternal and costal cartilage origin, and, after clearing, note the sterno-clavicular articulation. Take a strong knife and cut the six upper costal cartilages on each side near the rib junctions. Cut the muscles of the six upper intercostal spaces along the lower edges of the ribs, from the cuts through the cartilages to the sternum. Raise one side of the plate included by incisions, and turn it off from the pleural surfaces, the pericardium, and the contents of the anterior mediastinum. We cut with scissors the internal mammary arteries and branches, the origins of the sterno-hyoid and thyroid muscles; the fascial attachments of the slips to the omo-hyoid muscles, and the sternal origins of the sterno-cleido-mastoid muscles. Clear the surfaces of the sternum and costal cartilages, the sterno-chondral ligaments, the perforating arteries, and the intercostal aponeurosis. Dissect the anterior intercostal aponeurosis in the 2d and 3d intercartilaginous spaces, thus exposing the anterior surfaces of some of the internal intercostal muscles. Remove the anterior sterno-chondral ligament. Turn up the sternal plate and note its interior, the origins of the sterno-hyoid and sterno-thyroid muscles, the surface of the triangularis sterni muscle, and the interior branches of the internal mammary arteries. Remove the triangularis sterni muscle and trace a part of the internal mammary arteries.

Note the parietal layers of the pleuræ and the anterior portion of the parietal layer of the pericardium. Retain the pericardial and mediastinal branches of the internal mammary arteries.

Cut away portions of the anterior pleural surfaces. Note a portion of the diaphragm. Hook the lungs outwards, and find the phrenic nerves and the comes nervi phrenici arteries. Clean the thymus gland, the innominate veins, and the superior vena cava.

Cut the superior cava a little below the right innominate. Cut the right innominate just above its termination and the left innominate just external to that portion of the thymus gland which ascends into the neck. Remove the thymus gland and the veins between their points of section. Find the pneumogastric and recurrent laryngeal nerves, and the cervical and thoracic cardiac nerves. Expose the anterior surface of the transverse aortic arch, the innominate artery and its bifurcation, and the left subclavian and left common carotid. Clear the right bronchus behind the aortic arch and the trachea above it.

Free the lungs from the thoracic wall with the hand. Separate the diaphragmatic parietal pericardium from the diaphragm. Cut the innominate artery at its bifurcation, the left common carotid and the left subclavian at the same level (and with these vessels the cervical cardiac nerves). Cut the phrenic nerves and their arteries at their superior and inferior ends, removing the sectioned portions. Cut the recurrent laryngeal nerve at its origin and where it is seen in front of the gullet, the thoracic branches of the vagus, and the trachea. Grasp the trachea just below where it was cut, and with it the arch of the aorta, draw the lungs and heart downwards and forwards, cut the trachea and bronchi from the gullet and the descending portion of the aortic arch from the chest wall, and cut the pulmonary branches of the pneumogastric nerves. Cut the great azygos vein, as it enters the posterior surface of that portion of the superior cava which is without the pericardium.

Cut the arch of the aorta where its descending portion joins the thoracic aorta. Cut the inferior vena cava at its emergence from its diaphragmatic foramen. Remove the viscera in a mass.

Study the interior of the thoracic cavity.

Clean the thoracic portion of the gullet, and note, below and upon it, the vagi and their œsophageal branches.

Expose the dorsal portion of the right sympathetic nerve, its splanchnic branches, and the branches which join the intercostals.

Clean and study the thoracic aorta, œsophageal arteries, and the intercostal vessels and nerves of the right side.

Remove from an intercostal space (near the vertebral column) the posterior aponeurosis and the posterior intercostal muscle, and note the relation between the external and internal intercostal muscles.

Cut the œsophagus on a level with the 2d interspace above, and, if the abdomen has not been dissected, cut again near the opening for the gullet in the diaphragm. Remove this section. Cut the vagi above and below and remove the sections. Open the œsophagus longitudinally and study its interior.

Cut the aorta just before it passes through the diaphragm, and its intercostal branches a little external to their origins. Remove the aorta and intercostal stumps, being careful not to injure the small azygos vein and the left intercostal veins. Study the dorsal portion of the left sympathetic nerve and trace the course of the intercostal nerves and vessels of the left side. Find the upper thoracic portion of the thoracic duct on the 3d dorsal vertebra, and trace it up to the neck.

Remove the pleura from the superior surface of the diaphragm. Study the diaphragmatic openings and their contents and trace the thoracic duct from its passage through the diaphragm to the 3d dorsal vertebra.

Thoracic Viscera.

Lay the lungs and heart upon their posterior surfaces. Hook the lungs aside and note the pericardium. Open the parietal layer of the pericardium from the anterior portion of the beginning of the aorta to the cardiac apex. Make an incision outwards in both directions from the upper part of this incision. Reflect flaps.

Study the anterior surface of the heart and the great vessels at its base.

Turn the heart and lungs so as their posterior surfaces are looking upwards. Cut the aorta between the left carotid and left subclavian. Remove the parietal layer of the right and left pleuræ posteriorly. Draw the lungs aside. Note the bronchi, the trachea, the heart and vessels, and the pericardium.

Clean the trachea and bronchi posteriorly. Retain the nerves and veins, and the branches of the bronchial arteries.

Draw out superiorly and towards the right the superior cava. See the stump of the great azygos vein.

Below the bifurcation of the trachea find the right pulmonary artery. Find the left pulmonary artery on a plane anterior to the left bronchus. Between the arch of the aorta and the left pulmonary artery observe the ductus arteriosus. Study the roots of the lungs.

Cut the pulmonary arteries and veins, remove from the heart the trachea and bronchi, and cut again the pulmonary arteries, veins, and nerves, close to the lungs. Study the lungs.

Lay the lungs on their posterior surfaces and study their anterior surfaces.

Turn the lungs over and study their posterior surfaces.

Study the orifices of the pulmonary vessels. Slit open the bronchi and vessels and trace them into the lungs.

We must now investigate the heart.

Place the heart with its right side looking up. Remove the pericardium and adipose tissue from the right surface. Note the coronary artery and its branches. Expose the anterior surface of the beginning of the aorta and pulmonary artery, and the ending of the two vena cavæ. Study the outer surface of the right side of the heart.

Turn the organ so that its left side is up. Remove the pericardium and adipose tissue. Note the coronary vein and artery. Expose the aorta and pulmonary artery posteriorly. Study the outer surface of the left heart.

Open the right heart by making an incision from the anterior surface of the termination of the superior cava to near the base of the right auricle, sweeping it along the base and terminating it at the ending of the inferior cava. Raise the flap and study the interior of the auricle.

Open the right ventricle by slitting up the pulmonary artery on its anterior surface, continuing the incision to near the furrow between the ventricles towards the apex, and ending it by a semicircular sweep upwards. Raise the flap and study the interior of the ventricle.

Open the left auricle by an incision running from the termination of the inferior vena cava posteriorly into the appendix, and another incision extending up from the middle of this and terminating between the upper pulmonary veins.

Replace the left auricular flaps.

Open the left ventricle by an incision splitting the aorta, running beneath the appendix of the left auricle, sweeping along the left posterior surface to near the apex and then recurring by an upward sweep to the middle of the posterior lateral surface. Hold open the flaps and study the interior.

Dissection of the Back.

POSITION: The body lies prone, with a block under the chest and another under the pelvis.

FIG. 8.



INCISIONS: 1. From the external occipital protuberance to the sacrum.

2. From the termination of No. 1 outwards on each side along the iliac crests.

3. From the external occipital protuberance outwards to the middle of the ear posteriorly.

4. From the acromion around the posterior surface of the arm;

5. Along the spine of the scapula to the acromion;—if the arm has not been dissected posteriorly.

DISSECTION: Reflect the skin from the middle line. Note the vessels and nerves of the subcutaneous tissues. Remove the subcutaneous tissues, leaving the nerves and arteries.

Note the occipital artery, subcutaneous arteries, a number of important nerves, and the superficial fascia.

Remove the superficial fascia. Note the scapular fascia, the trapezius and latissimus dorsi muscles.

Cut the origin of the trapezius from the middle line and superior curved line of the occipital bone, and turn it out laterally.

Dissect it loose (to do this) from its perforating nerves and arteries.

Cut the latissimus dorsi muscle from the lumbar fascia and from its spinal origin, treat its nerves and vessels as we did those of the trapezius, and turn it outwards. Clean and study the two rhomboid and the levator anguli scapulæ muscles, and the arteries between and upon them.

Clear the serratus posticus inferior, and the internal posterior portions of the external and internal oblique muscles.

Cut the two rhomboid and the levator anguli scapulæ from their vertebral origins, and turn the muscles externally. Clean and study the serratus posticus superior muscle.

Remove this last-mentioned muscle, and clean and study the splenius capitis and colli muscles.

Remove from their attachments the splenius capitis and colli muscles and the serratus posticus inferior. Clean the fascia upon the erector spinæ mass from the cranium to the sacrum.

Remove this fascia and note the erector spinæ muscle. Trace the sacro-lumbalis and the longissimus dorsi divisions superiorly.

Turn the superior portion of the sacro-lumbalis and observe the musculus accessorius, reflect this externally and note the cervicalis ascendens.

Remove the two last-named muscles and find the transversalis colli, trachelo-mastoid, complexus and spinalis dorsi muscles.

Remove from their attachments the complexus trachelo-mastoid and transversalis. Cut from the ribs and from the dorsal vertebræ the longissimus dorsi, and from the ribs the sacro-lumbalis. Remove from the ilium and lumbar vertebræ the erector spinæ mass. Remove the spinalis dorsi muscle, and clean the semi-spinalis dorsi and the semi-spinalis colli muscles.

Remove the two last-named structures. Clean the levatores costarum. the multifidus spinæ, the interspinales and intertransversales muscles and the aponeurosis of the transversalis abdominis. Note the posterior divisions of the dorsal and lumbar nerves and their companion arteries.

Clean the rectus capitis, posticus minor and major, the inferior and superior oblique muscles. Trace the deep cervical and princeps-cervical and posterior branches of the vertebral arteries, and the posterior branches of the cervical spinal nerves.

Remove the recti muscles, the two oblique and the multifidus spinæ muscles. Clean the vertebræ, the sacrum, the coccyx, and the posterior intervertebral and costo-transverse ligaments.

Saw the posterior portion of the occipital bone on each side, downwards and inwards into the middle of the edges of the upper quadrants of the foramen magnum.

Saw the cervical vertebræ on each side within their articular processes.

Saw the dorsal and lumbar vertebræ on each side at the roots of their transverse processes, and the sacrum internally to its posterior foramina. Remove the pieces. Note the dura mater of the cord, the cauda equina, and the roots of the spinal nerves.

Open the dura from the foramen magnum to its lower end.

Note the subdural space and the cord covered by the arachnoid and the pia mater.

Remove laterally upon one side the cervical dura mater and note the anterior and posterior spinal nerve roots and trace them. Find the spinal accessory nerve, and observe the ganglia upon the posterior roots of the spinal nerves.

Anterior Scapular Muscles (Dissect either with arm or back).

Position : Body as above.

Dissection : Expose the serratus magnus and subscapularis muscles.

To remove the upper extremity from the trunk cut the great serratus muscle (if the anterior of the thorax and neck, and the region back have been dissected).

Scalp, Superio-lateral Portion of Brain Membranes. Interior of Base of Cranium.

Position : As for dissection of the thorax. Head twisted upon one side. Shave the scalp.

Incisions : 1. From between the two superciliary ridges longitudinally back to the external occipital protuberance.

2. Around the head, joining the beginning and termination of No. 1 (running above the eyebrows).

DISSECTION : Reflect the skin. Find in the subcutaneous tissue the frontal and the anterior branches of the superficial temporal arteries, and the supratrochlear nerve. Expose the anterior portion of the occipito-frontalis muscle.

FIG. 9.



Cut out from the above-named muscle the supra-orbital artery and nerve, and trace them.

Find laterally in the subcutaneous tissue the superficial temporal artery and vein.

Cut from beneath the superficial temporal fascia the branches of the facial nerve. Find and trace the auriculo-temporal nerve. Expose the occipital and posterior auricular arteries, and the small occipital nerve, and a branch from the 3d cervical nerve.

Remove the occipito-frontalis muscle (cutting nerves and vessels).

Reflect the superficial temporal fascia with the ear muscles downwards, and cut it off. Note the deep temporal fascia, and incising it just below the temporal ridge, cut it away. Observe the temporal muscle. Cut away the lower one-half of the temporal muscle and observe the deep temporal vessels and nerves.

Saw the bones, the lines corresponding to the skin incisions.

Finish with a chisel. Drag off the skull cap.

Note the dura mater, the meningeal arteries and the Pacchionian bodies.

Incise the dura mater longitudinally in the median line and expose the interior of the superior longitudinal sinus.

Cut away the dura from one hemisphere (its exposed part). Note the subdural space and arachnoid membrane. From the posterior one-half of this hemisphere remove the arachnoid. Note the subarachnoid space and pia mater. Remove a portion of the pia posteriorly, and observe the cortex.

Cut the dura around on a level with the bony section. Lift the anterior cerebral lobes and cut the falx from the ethmoid. Lift up the posterior lobes and cut the tentorium at its circumference, not opening the lateral sinuses. Lift up the anterior lobes again, loosen one olfactory nerve and leave it on the brain, cut the other and leave it upon the ethmoid bone. Cut the optic nerves behind the commissure and the internal carotid arteries. Open the coronary sinus and turn out the pituitary body. Continuing to raise the brain cut the 3d, 4th, 5th, 6th, 7th, and 8th nerves, the auditory and the vertebral arteries, the pneumogastric, glossopharyngeal, and hypoglossal nerves, the spinal accessory nerve, and the spinal cord. Pass the fingers of one hand into the posterior fossa, and drawing the organ forward lift out the brain.

Trace the sinuses and arteries on the surface, especially at the base.

Cut the arteries, remove the pia, and note the fissures, the gyri and the sulci. Study the parts at the base.

Cut off the vertex of one hemisphere and the anterior border of the opposite temporo-sphenoidal lobe, to study the gyri and sulci.

Section the uncut vertex on a level with the cut one. This plane is the centrum ovale of Vieussens. Observe the corpus callosum.

Make an antero-posterior section of the corpus callosum on each side of the raphé, and turn off the roof of the lateral ventricles.

Cut away the roof of the ventricles, and cut off an additional piece of the occipital lobe of the right hemisphere.

Study the lateral ventricles and their cornua, the foramen of Monro, the septum lucidum, the corpora striata, and the tænia semicircularis.

Cut the corpus callosum. Reflect its anterior portion and expose the 5th ventricle, and lift its posterior portion from the fornix. Note the choroid plexuses and the optic thalami.

Section the fornix anteriorly and reflect each half. Observe the velum interpositum, the anterior pillars of the fornix, and the veins of the interior of the ventricles.

Cut the veins which pass into the venæ Galeni and remove the intraventricular veins. Turn off the velum interpositum and the choroid plexuses posteriorly. Note the venæ Galeni, the choroid plexus of the 3d ventricle, the 3d ventricle, the pineal gland, and optic thalami.

Return into place the velum interpositum and fornix. Section the fornix and choroid plexuses transversely and remove their anterior portions, cut transversely through the outer wall of the left lateral ventricle to the floor of the middle cornu. Continue a curved incision anteriorly on a level with the floor. Remove by a transverse cut a piece of cerebrum.

Make an anterior transverse section of the basal ganglia of the right hemisphere.

Make a posterior transverse section of the basal ganglia of the right hemisphere.

Cut the posterior pillar of the fornix and the splenium of the corpus callosum of the left hemisphere, and carry the cut into the posterior part of the longitudinal fissure. Remove by this cut the left temporo-sphenoidal and occipital lobes.

Remove the same lobes of the right hemisphere. Remove the left frontal lobe. Expose the 4th ventricle.

Cut the peduncles of the cerebellum. Cut away the right amygdala. Study the surface of the cerebellum.

Section the cerebellum by a transverse antero-posterior cut of the middle of the vertical diameter of the right hemisphere, and a transverse vertical cut at the middle of the antero-posterior diameter of the left hemisphere.

Cut away the cerebrum anterior to the pons. Note the optic tract and geniculate bodies, the corpora quadrigemina, the processi e cerebello ad testes, the valve of Vieussens, the 4th pair of nerves, the cerebral peduncles, the pons, the medulla, and many nerve roots.

Open the 4th ventricle by two longitudinal cuts along the lines of junction of the valve of Vieussens and the processi e cerebello ad testes. Reflect the valve forward.

Superficial Region of the Face.

Cut off the hair, stuff the nostrils, push oakum under the lids and sew the lids together. Push oakum between the teeth and the cheeks and lips and sew the lips together.

POSITION : Body same as for thorax. Place head laterally on a block.

INCISIONS : 1. From between the eyebrows down the centre of the nose to the mucous membrane of the lip, around the mouth to the middle of the lower lip, down to the middle of the inferior border of the lower jaw.

2. Outwards along the lower edge of the body and upwards along the ramus of the lower jaw, carrying the ascending line to meet the circular line around the scalp.

DISSECTION : Remove the skin.

Remove the subcutaneous tissue carefully with scissors and note the risorius and platysma myoides muscles.

Cut the last-named muscle along the border of the lower jaw and reflect its superior portion upwards and inwards (risorius with it). This exposes fascia. Clear this fascia from one-half the orbicularis oris from the depressor of the angle of the mouth, the depressor of the lower lip, the zygomatici, the levator labii superioris alæque nasi, and orbicularis palpebrarum muscles.

Note the submental artery and vein, the nasal and frontal arteries and the supra-trochlear nerve. Cut out from the fascia over the masseter muscle the branches of the facial nerve. Dissect and trace the facial artery and vein.

Clean the surface of part of the pyramidalis nasi compressor naris, lifter of the upper lip and the angle of the mouth, buccinator and masseter muscles.

Trace the superficial temporal artery, the auriculo-temporal nerve, branches of the facial nerve, and the transverse facial artery.

Remove the fascia from the external surface of the parotid gland and note Steno's duct.

Cut Steno's duct just before it perforates the buccinator muscle. Reflect the duct back to the gland, reflect the gland externally and section it so as to leave in place its posterior portion. Reflect this portion outwards. Trace the facial nerve and branches, the transverse facial and superficial temporal arteries and the bifurcation of the external carotid artery.

Clean the masseter muscle and the raiser of the upper lip.

Note the facial vein and artery (facial portions) and the branches of the artery.

Expose these vessels throughout their course.

Clean the compressor naris and pyramidalis nasi muscles.

Remove branches of the facial nerve. Cut the masseter muscle through transversely and remove it (noting its nerve and artery).

Observe the lateral ligament of the temporo-maxillary articulation.

Remove the remaining portions of the zygomatici, the levator labii superioris alæque nasi, and the levator labia superioris muscles.

Cut the facial vein and artery on the body of the lower jaw, and remove their superior portions. Note the infraorbital artery and nerve, and the buccal branches of the inferior maxillary.

Clean the levator anguli oris and buccinator muscles.

Cut the depressor anguli oris and depressor labii inferioris muscles transversely across their centres, and remove their superior portions.

Note the mental nerve, the mental artery and the levator menti muscle.

Remove the orbicularis palpebrarum muscle and note the corrugator supercilii muscle, tarsal ligaments, etc. Cut the lid sutures and remove the oakum. Study the conjunctival sacs and the puncta lachrymalia, and slit up the latter to find the lachrymal canals and the lachrymal sac.

Open the palpebral ligament and note the supraorbital and frontal arteries, the supraorbital and supratrochlear nerves, and portions of two of the orbital muscles and the lachrymal gland.

Remove the inferior portion of the palpebral ligament and note the inferior oblique muscle, the nasal artery, and the infratrochlear nerve.

The Cavity of the Orbit.

EXPOSURE: Detach the orbital periosteum at the superior orbital border and push it off from the bone. Saw through the supraorbital portion of the frontal bone into the orbit, the lines being external to the internal angular process and internal to the external angular process. Chisel out a V-shaped piece from the orbital roof (the brain has been removed). Remove this piece and with it the supraorbital portion of the frontal bone. Incise the orbital periosteum and clean out fat and find the frontal nerve and supraorbital artery, the lachrymal artery, nerve, and gland, and three orbital muscles.

Remove the inner portion of the lesser wing of the sphenoid bone. Expose the 1st, 3d, and 4th nerves, the ophthalmic division of the 5th nerve, and the ophthalmic artery.

Cut the frontal nerve and supraorbital artery and remove their anterior portions. Also in the same way the lachrymal nerve and artery, carrying the gland with their anterior portions. Cut the levator labii superioris, remove its anterior portion and with it the upper tarsal cartilage. Find the infraorbital fibrous ring. Clean the rectus superior muscle. Find the ophthalmic vein and artery and nasal nerve.

Cut the rectus superior near its origin, and the superior oblique near its origin, and again just before it reaches its pulley. Cut the ophthalmic branch of the 5th pair and the nasal branch of the ophthalmic nerve. Remove the nerves and muscles. Cut the fibrous ring, reflect the internal portion of the ring with the origins of the superior oblique and levator labii superioris internally. Dissect away the external portion. Carry externally the superior attachment of the external rectus. Remove the ophthalmic vein. Trace the 6th nerve. Cut away the branch from the ophthalmic artery to the external rectus muscle and so find the ophthalmic ganglion. Trace the 3d nerve, the nasal nerve and its branches, the ophthalmic artery and its branches, and the optic nerve. Clean part of the superior surfaces of the external and internal rectus muscles.

Cut the ophthalmic artery, the motor root of the ciliary ganglion, the nasal nerve, the ciliary nerves and arteries. Remove the ganglion, the ophthalmic artery, and the nasal nerve. Note the central artery of the retina and cut it. Cut the optic nerve between its entrance into the orbit and the arteria centralis retinæ, and remove this portion of the nerve.

Cut the external rectus muscle and remove its anterior part.

Lift the eyeball posteriorly, note the inferior branch of the 3d nerve, clean the internal rectus and the two inferior eyeball muscles.

Middle Fossa of the Cranium.

Peel off the dura mater on one side. Notice the stumps of the facial nerve-roots, the ganglion of Gasser with its divisions, the middle meningeal artery, and the large and the external superficial petrosal nerves.

Cut the superior and inferior maxillary divisions of the sensory root of the 5th pair, remove the ganglion and a part of the motor root. See the internal carotid artery and the large superficial petrosal nerve.

Deep Region of the Face.

Incisions: Saw through the temporal zygoma and through the malar bone.

DISSECTION: Remove the sectioned bone. Clean externally the inferior portion of the temporal muscle.

Cut the temporal muscle from the coronoid process (retaining artery and nerve stumps). Saw through the inferior maxillary bone at the neck of the condyle and in the middle of the ramus below Steno's duct. Remove the sectioned portion. Remove the remaining portion of the parotid gland. Remove the skin posterior to the ramus. Find the posterior belly of the digastric muscle and the stylo-hyoid muscle and a part of the external carotid artery. Trace the internal maxillary artery and its branches, note the temporal, buccal, and masseteric nerves, the temporo-maxillary articulation, the external pterygoid muscle, part of the internal pterygoid muscle, the gustatory, and the inferior dental and mylo-hyoid nerves.

Cut the internal maxillary artery and the alveolar branch, and again just after the origin of the middle meningeal. Remove the included portion. This exposes the external pterygoid muscle.

Remove the condyle of the lower jaw. Note the joint and its ligaments. Remove the fibro-cartilage of the joint.

Remove the external pterygoid muscle from its anterior attachments. Find its nerve and leave it. Remove the fat from the upper outer surface of the internal pterygoid. Trace the deep temporal, gustatory, and other nerves. Find and trace the chorda tympani nerve. Expose the middle and small meningeal arteries.

Draw the inferior maxillary division of the 5th nerve internally. Note the Otic ganglion.

Cut the external carotid artery above the origin of the posterior auricular. Cut the middle meningeal and other branches of the internal maxillary and remove. Cut the auriculo-temporal and chorda tympani nerves. Remove the Otic ganglion with its roots.

Remove the internal lateral ligament of the lower jaw. Expose the tensor palati, levator palati, and inferior constrictor muscles (in part), and the internal carotid artery, the stylo-glossus and stylo-pharyngeus muscles, and the glosso-pharyngeal and spinal accessory nerves.

The Neck, Anteriorly and Laterally.

Position : Same as for dissection of the arm. Put the lateral surface of the head on a block.

Incisions : 1. From the symphysis of the jaw down the middle line of the neck to the top of the sternum.

2. Along the body and ramus of the jaw (if the face is not dissected).

3. Outward along the clavicle (if the anterior thoracic region is not dissected). Turn the head and incise the opposite side.

Dissection : Reflect the skin and the subcutaneous tissue from the platysma myoides muscle. Study this muscle.

Cut the right platysma myoides along the body of the lower jaw and reflect it downwards, respecting the external jugular vein.

Remove the fascial sheath covering the outer surface of the sterno-cleido-mastoid muscle, leaving in place the external jugular vein, and the nerves of this region.

Trace the external jugular vein and branches of the cervical plexus. Remove the lower and outer portion of the spinal accessory nerve.

Lift up the anterior border of the trapezius muscle. Note the great occipital nerve, the occipital artery, and a branch from the 3d cervical nerve.

Clear the space between the anterior border of the trapezius and the posterior edge of the sterno-cleido-mastoid, leaving in place the nerves and vessels (removing the fascia).

Note a number of muscles, many nerves, the subclavian artery and vein, the superficial cervical, supra-scapular, and other arteries.

Remove the fascia in front of the sterno-cleido-mastoid and study this space. Find the submaxillary gland, the internal jugular vein, the carotid arteries and their branches, the hypoglossal, descendens noni, and superior laryngeal nerves, and many muscles.

Clean the mylo-hyoid, anterior belly of the digastric, and adjacent structures.

Turn the head and dissect the other side of the neck to the same stage. Draw the head back in the middle line with the block removed.

Study the hyoid bone, the thyro-hyoid (in part), the omo-hyoid, the digastric and sterno-hyoid muscles, the facial artery and vein, the submental artery, and other structures, vascular and nervous.

Again place the head on its left side with a block. Cut the facial and external jugular veins, the superficial cervical and the great and small occipital nerves, and the nerve to the trapezius. Remove the nerves and reflect the veins. Cut both heads of the sterno-cleido-mastoid muscle near their points of origin. Reflect the muscle upwards. Remove the inner layer of its sheath. Note the middle of the omo-hyoid muscle with its slip of cervical fascia, the sterno-hyoid muscle, the right lymphatic duct, the phrenic and other nerves, the common carotid and other arteries, and the internal jugular vein. Clean the muscles on the floor of the posterior triangle.

Hook up the submaxillary gland and the parotid (if remaining). Note the mylo-hyoid and digastric and portions of the stylo-hyoid and stylo-glossus muscles. Find the laryngeal arteries.

Turn the head and dissect the opposite side of the neck to the same stage.

Cut both sterno-hyoid muscles on a level with the hyoid bone and reflect downwards. Put the head in the middle line and take away the block. Clean the sterno-thyroid muscles, note the fascia of the trachea, the larynx, the cricoid artery, the hyoid bone and thyro-hyoid membrane, and part of the thyroid gland.

Remove the fascial slip to the left omo-hyoid and expose the left lymphatic duct and the 1st part of the subclavian artery.

Cut the anterior belly of the left digastric near its insertion and reflect it, cut the fascial slip of this muscle. Remove the whole of the digastric, the slip, and the stylo-hyoid muscle.

Cut the left mylo-hyoid longitudinally, near the raphé, from the lower jaw to the hyoid bone. Cut it outwards from the hyoid bone, reflect it up, and cut it from the jaw. Cut away a part of the left submaxillary gland and trace its duct. Find the stylo-glossus muscle, gustatory nerve, and hypoglossal nerve. Clean the genio-hyoid and hyoglossus muscles.

Cut the thyro- and omo-hyoid muscles of the right side, and put the head again on its left side. Cut the communicans and descendens noni nerves and the external laryngeal artery, and reflect with them the omo-hyoid and thyro-hyoid muscles. Cut the internal jugular vein just below the posterior belly of the right digastric and reflect it downwards. Expose the pneumogastric and its branches.

Clean the innominate artery in front and the trachea. Retain the inferior thyroid vein and artery, and the recurrent laryngeal nerve.

Study the origin and bifurcation of the right carotid. Study the first portion of the right subclavian artery with its branches. Cut the anterior belly of the right digastric muscle above the fascial loop and reflect it forwards. Cut the posterior belly and the stylo-hyoid back of the fascial loop and reflect them backwards.

Clean the hypoglossal nerve at its upper part, the lower part of the internal carotid artery, and the lower part and branches of the external carotid. Note the constrictors of the pharynx and the stylo-glossus muscle, and the glosso-pharyngeal nerve. Dissect the left side of the neck to this stage.

Put the head in the middle line without the block.

Cut the right common carotid artery near its origin and the pneumogastric nerve at the same level, and reflect them upwards. Remove the posterior layer of the deep fascia. Note the right sympathetic nerve.

Clean the 1st and 3d portions of the subclavian artery, the subclavian vein, and the brachial plexus.

On the left side remove the common carotid and the lower portion of the external carotid with its branches, and the lower portion of the internal carotid. Remove the left pneumogastric nerve. Find the left sympathetic nerve and remove it. Note the thyro-hyoid muscle, the larynx and hyoid bone, the thyro-hyoid membrane, thyroid gland, and inferior thyroid veins.

Expose the left recurrent laryngeal nerve, the thoracic duct, the 1st portion of the left subclavian artery and its branches, parts of the œsophagus and brachial plexus, and the longus colli muscle.

Reflect downwards the remains of the left submaxillary gland. Note the submaxillary ganglion with its branches.

Cut at the root of the neck the inferior thyroid veins and arteries, the recurrent laryngeal nerves and the œsophagus. Raise the trachea and gullet, dissecting them from the vertebræ and muscles. Continue dissecting up to the base of the cranium, removing the pharynx from the vertebræ and muscles. Reflect to the base of the skull the internal jugular veins and the internal carotid arteries, the 9th, 10th, 11th, 12th, and sympathetic nerves. Saw the cranial walls on each side back of the external auditory meatus. Drag the reflected parts anteriorly and saw the base of the skull posterior to the styloid processes of the temporal bones. Remove the front of the head with the reflected parts. Study the rectus capitis anticus major muscles. Remove the lower portion of these muscles. Clean the right longus colli, the lower portion of the left rectus capitis anticus minor, and part of the left rectus lateralis. Remove the lower portion of the right small anterior rectus, and the rectus lateralis. Remove the left longus colli muscle. Note the superior cervical nerve (anterior branches) and the left vertebral artery. Note the scaleni muscles.

The remaining undissected portion comprises the pharynx and its nerves and arteries, the soft palate, the hard palate, the tongue, the larynx and adjacent parts.

POSITION: The anterior sawn portion of the head with its attached structures contains the above-named parts.

Saw the right and left ramus of the lower jaw bone just above the angle. Stuff the pharynx with oakum. Hang up the anterior portion of the head to the top bar of a square frame (like a large slate frame), fasten the lower portion to the lower bar and make tense.

DISSECTION: Cut away the left internal carotid artery, the upper part of the left external carotid, the left hypoglossal, pneumogastric, spinal accessory, and glosso-pharyngeal nerves and their branches. Remove the left lobe of the thyroid gland, the inferior laryngeal artery and recurrent laryngeal nerve of the same side. Remove the upper end of the right external carotid artery and the right facial artery (leaving the ascending palatine branch).

Study the 9th, 10th, 11th, and 12th nerves and their branches, the pharyngeal plexus, the ascending pharyngeal artery, and the internal carotid artery.

Expose the constrictors of the pharynx, the stylo-pharyngeus muscle and stylo-maxillary ligament.

Study the palate and tongue.

Cut away the lateral pharyngeal walls and the upper part of the œsophagus. Look into the larynx from above.

Dissect the larynx, study its arteries, nerves, muscles, cartilages, etc. Saw near the median line through the base of the cranium and study the nares.

THE END.

ESSENTIALS OF HUMAN ANATOMY.

Give the derivation and meaning of the term anatomy.

From two Greek words, literally meaning dissection, but it is used to indicate the study of the physical structure of organized bodies.

How is human anatomy divided?

Into two great divisions, viz.: 1. General or descriptive anatomy, which deals with the separate parts of the human body; and 2. Surgical or regional anatomy, which describes the relations individual parts, such as muscles, nerves, arteries, etc., bear to each other, in order to enable the surgeon to find or avoid important structures when operating.

What is osteology?

A subdivision of general anatomy describing the number, form, structure, and uses of the bones.

What is the chemical composition of osseous tissue (bone)?

About one-third (33.30) is organic or animal matter, resolvable into gelatine after prolonged boiling, with traces of chrondrigen (the proximate principle of cartilage), and two-thirds inorganic, or mineral, consisting of calcium phosphate (tribasic) 51.04, calcium carbonate 11.30, calcium fluoride 2, magnesium phosphate 1.16, and sodium chloride with traces of sodium dioxide 1.20. Either the organic or inorganic matter may be removed without affecting the form of the bone; the former by exposure to heat with free access of air, after which slight force will reduce the bone to powder; the latter by steeping in dilute hydrochloric acid, which will render a long bone as pliable as a strip of rubber.

Does an increase of the mineral constituents take place in old age; rendering the bones more brittle than in youth?

No; although this is the common statement in text-books, for while equal *bulks* of young and old bones do show marked differences in the proportion of earthy and animal constituents, equal *weights* do not, so that the elasticity in youth and the brittleness in age depend upon the greater sponginess of texture in young bones.

Is bone a homogeneous substance?

No; for while the exterior is composed of a *compact* (hard, ivory) layer, the inner portions are formed of *spongy* or *cancellous* tissue; while the interior of long bones is hollow, forming the medullary (marrow) canal. The compact tissue is used upon the exterior or in the shaft of long bones where "cross-strain" is greatest, whereas cancellous tissue enables the articular end of the bones to be large, for security, yet light, this tissue being capable of bearing enormous pressure without yielding, but incapable of bearing much "cross-strain."

Describe the microscopic structure of bone.

In transverse section, with a low power, a number of holes will be observed, averaging $\frac{1}{500}$th inch in diameter, surrounded with a series of tolerably concentric circles, consisting of an interrupted series of dark spots. With high powers, the holes, called *Haversian canals* for the passage of vessels, are seen to be surrounded by a series of concentric lines, termed lamellæ, while the dark spots reveal themselves as cavities in the bone, called *lacunæ*, intercommunicating with each other and the central Haversian canal, by means of delicate radiating hollow tubes, called *canaliculi:* this aggregation of structures is called an *Haversian system.* The lacunæ contain bone corpuscles, processes of which extend into the *canaliculi.* As each Haversian system communicates by its canaliculi with those of its neighbor, the exterior of the bone, all portions of its most compact tissue and the medullary canal freely intercommunicate and receive nutrient material. As these systems are circular they would leave interspaces where not in contact;

Pattie

but these gaps are filled up by layers of bone with *lamellæ* and *canaliculi*, these portions being called *interstitial lamellæ*. In addition to the *concentric lamellæ* around the Haversian canals, an examination of a cross-section of a bone long exposed to atmospheric influences will show that it is composed of a series of circumferential *fundamental lamellæ* running around the whole circumference of the bone.

What is the periosteum?

It is a fibrous membrane serving as a scaffolding to enable the bloodvessels to reach all portions of the exterior of the bone except its articular ends and the points of attachment of strong tendons, the layer in contact with the bone containing many *osteoblasts* or bone-forming cells upon which the growth in thickness of the bone depends.

What is the endosteum?

This term is applied to the delicate connective tissue lining of the medullary and cancellous tissues which contains numerous bone-forming cells (osteoblasts).

Describe the medulla or marrow.

There are two varieties. In adult bones—except the cancellous tissue of their articular extremities, that of the vertebral bodies, the sternum, ribs, and cranial bones—the marrow is *yellow*, consisting almost solely of fat (adipose) tissue (ninety-six per cent.). In the other situations just mentioned the marrow is red (fœtal), contains not more than one per cent. of fat, and numerous oval, many-nucleated cells (①myeloplaxes). This kind of marrow is that found in the fœtus and infants.

h. ent. [illegible]

Do bones receive blood only from vessels in the periosteum?

No, for the medullary tissue of all long bones receives a good-sized artery (the nutrient artery), which obliquely penetrates the compact tissue, after which it divides into two main branches, one ascending the other descending in the medullary canal; the veins chiefly emerge through numerous openings near the articular ends of the bones.

① Marrow - plaque

Describe the process of ossification.

There are two methods, viz., the *membranous* and the *cartilaginous*. In the former, at the centre of ossification of a parietal bone for instance, the bone-forming cells (*osteoblasts*) arrange themselves—about the second month of intrauterine life—along the thick bundles of fibrous tissue which radiate from the centre of the future bone; by the deposition of lime-salts in these osteoblasts, the deposit of bone shoots out in needle-like rays toward the circumference. Ossification by cartilage is essentially similar; for *protection of vital centres*, as at the base of the skull, or to maintain *rigidity of parts and proper tension of muscles*, as in the extremities, certain bones are laid down in cartilage. Just before ossification commences the cartilage cells multiply, arrange themselves in columns—especially at the ① epiphyseal ends—and the intercellular material becomes infiltrated with lime-salts. Now ossification proper commences by the ingrowth from the periosteum of buds of young connective tissue covered with bone-forming cells, which, after causing absorption of the cartilage, become converted into bone. The first bones to ossify are 1, the clavicle; 2, the inferior maxilla (fifth to seventh fœtal week).

Of how many bones is the adult human skeleton composed?

Two hundred, including the os hyoides, but excluding the teeth, Wormian bones, all ② sesamoid bones, except the patellæ, and the ossicles of the middle ear.

Into what classes are bones divided, and give a few instances of each?

Long bones, as femur, tibia; *flat*, as those of the vault of the skull, scapula; *irregular*, as the vertebræ; and *short*, such as the carpal or tarsal bones.

What are Wormian bones?

Irregular fragments, developed from supplementary centres, situated at the junction of two or more cranial sutures, where, during infancy, a membranous interval existed, viz., a *fontanelle*. From their triangular form they are often called *ossa triquetra*.

1.) [illegible] - to grow (2 bones + Cart, later all bone)
2.) Sesame seed - like

What are sesamoid bones ?

Those developed in the substance of tendons, whereby the muscles obtain additional *leverage—i. e., power;* the patellæ are, by some, classed with these bones.

Of what parts do long bones consist, and what are their uses ?

Of a shaft (body,① diaphysis), two articular extremities, and various processes; they are developed in cartilage, from one principal, and one or more additional (epiphyseal) centres of ossification; they serve as supports and levers for power and progression.

Where are short bones employed, and why ?

In the carpus and tarsus, where strength with limited motion is required. They ossify in cartilage.

Describe the structure and uses of flat bones.

They consist of two layers of compact tissue with interposed cancellous tissue, called ② *diploë;* they serve to protect important viscera, as brain, etc., and afford extended surfaces for origin of muscles; for the most part, they ossify in membrane.

How are bony prominences named ?

Processes, and are described as *articular* and *non-articular.*

Give the names and characteristics of the chief articular processes.

Head, a convex smooth projection, with a constriction or neck beneath; found in freely moving joints.

③ *Condyles,* double projecting processes, may have a constriction or neck—*i. e.,* neck of condyle of jaw.

④ *Trochanters,* short projecting levers near articulations to facilitate *rotation* of the bone on its long axis.

⑤ *Tuberosities,* roughened, broad prominences.

Tubercles, similar to the above, but small with reference to the size of the bone.

Spines and spinous processes, more or less pointed projections.

Apophysis, strictly speaking, any bony process which develops from the primary centre of ossification, commonly used, however,

① between — to grow
② double = a fold.
③ knuckle
④ wheel, pulley.
⑤ bump, swelling bone

for any process, even if an epiphysis, *after* it has coössified with the mass of the bone.

What is a diaphysis ?

The main portion of a bone (shaft in a long bone, body in an irregular one) between the *epiphyses*.

What is an epiphysis ?

A supplementary centre, usually to provide for growth in *length*, developed in cartilage, which remains separated by a layer of *epiphyseal cartilage* until the growth of the bone is completed, when it coössifies with the *diaphysis*, and all further growth ceases. Epiphyseal centres appear after birth: they coössify in the inverse order of their appearance, except that of the lower end of the fibula. This process of coössification commences about puberty, and the last to unite are those of the upper end of the tibia and the vertebral bodies—as late as twenty-five years.

Mention some of the non-articular processes, with their meaning.

Azygos, without a fellow; *coronoid*, or *coracoid*, like a crow's beak; *mastoid*, like a nipple; *rostrum*, a beak; *styloid*, pen-like; *squamous*, scaly; *vaginal*, ensheathing.

Name some of the articular cavities of bone.

Cotyloid, when they resemble a deep cup; ② *glenoid*, when they have a shallow-cup form; *trochlear*, pulley-like ③ *facet*, when smooth, like one of the surfaces of a cut gem; *sigmoid*, when curved in two opposite directions.[1] like letter S

What are the principal non-articular cavities called ?

Fossæ, shallow depressions ④ *sinuses*, deep cavities, communicating with the exterior by small openings; *grooves*, long narrow depressions; *fissures*, cracks; *notches*, deficiencies of edges of bones; *foramina*, holes through bones for transmission of nerves, etc.

① [illegible]

② glen a Cavity –

③ Facette = little face

④ Gulf, hollow

BONES OF THE HEAD.

Name the bones composing the head.

They are twenty-two in number: eight of which (*cranial*) compose the brain-case, viz., one frontal, two parietal, two temporal, one sphenoid, one occipital, and one ethmoid; fourteen are *facial:* two superior maxillary, two malar, two nasal, two lachrymal, two palate, two inferior turbinated, one vomer, and one inferior maxillary; the *ethmoid* also enters largely into the formation of the nasal cavities.

The Frontal Bone.

Of what parts does this bone consist?

Of a vertical and horizontal portion.

FIG. 1.



Describe the points on the outer surface of the vertical portion.

On each side of the median line are two *frontal eminences*[1], between which are the remains of, or the completely obliterated, *inter-frontal suture*, leaving a slight linear *depression*, which, above the root of the nose, terminates in a rounded, projecting *nasal eminence*^{6}. Extending outward from this on each side are two curved, rounded *superciliary ridges*^{7}, situated beneath which, between the two tables of the skull, lie the *frontal sinuses*^{14} (Fig. 2). The junction of the vertical and horizontal portions forms on each side a curved margin, the *supra orbital arch*$^{3-4}$, notched or perforated toward its inner part by the *supra-orbital notch*^{5}, or *foramen*, transmitting the artery, vein, and nerve of the same name. Each orbital margin terminates by two stout processes, called *internal angular*^{3} and *external angular*^{4} processes. Between the inner projects the *nasal spine*^{9}, in the rough, uneven space, called the *nasal notch*; the margin of the external angular process extends upward as a *temporal ridge*^{8} (Fig. 1).

FIG. 2.



Describe the points on the inner surface.

In the median line a *vertical groove*^{10} (Fig. 2) (for the longitudinal sinus) exists, whose edges coalesce below to form the *frontal crest*^{11}, which terminates as a notch, or perhaps complete *foramen cæcum*^{13}, which when pervious transmits a small vein.

Describe the horizontal portion.

This consists of two *orbital plates*^{12} of a triangular outline, separated by a quadrilateral *ethmoidal notch* (for articulation with that bone). A shallow *lachrymal*[13] depression (for gland) exists at the outer part of the orbit, also a smaller one or sometimes a small *tubercle*[15] at the anterior inner part for the pulley of the superior oblique muscle. Several half cells are seen along the margins of the ethmoidal notch, which complete the ethmoidal cells when the ethmoid is in position, as well as two *grooves*, which are likewise converted into the *~~anterior and~~ posterior ethmoidal ~~canals~~*, the former for the nasal nerve and anterior ethmoidal vessels, the latter for the posterior ethmoidal vessels. On each side of the nasal spine open the *frontal sinuses*[14] (absent in children), a part of the nasal cavities, into which they open by the *infundibulum*. The inner surfaces of both vertical and horizontal portions present numerous depressions for the convolutions of the brain, and branching grooves for the anterior meningeal artery.

What centres of ossification has this bone ?

Two, near the orbital arches, in ~~membrane~~, seventh or eighth fœtal week.

With what bones does it articulate ?

With two parietal, the ~~sphenoid~~, the ~~ethmoid~~, two nasal, two superior maxillary, two lachrymal, and two malar—~~twelve~~ in all.

What muscles arise from it ?

The corrugator supercilii, orbicularis palpebrarum, and temporal, on each side.

Parietal Bones.

Describe a parietal bone.

It is of a quadrilateral form, convex externally, the most prominent point called the *parietal eminence*. Its upper serrated border forms with its fellow the *sagittal suture*[1]; the anterior serrated margin articulates with the frontal, forming part of the *coronal suture*[3]; its anterior inferior angle[7], more prolonged than the others, articulates inferiorly with the sphenoid (*spheno-parietal suture*); the remainder of the inferior border[2], ~~bevel edged~~, is ~~overlapped by the squamous plate of the temp~~oral (*squamo-parietal suture*); the ~~posterior serrated~~ border[4] articulates with the occipital (*occipito-parietal* part of *lambdoid suture*); while the posterior inferior angle[8] joins the mastoid process of the temporal (*masto-parietal suture*). Crossing the middle of the bone, from before backward, is a curved line, the *temporal ridge*[5], and near the posterior superior angle is often a *parietal foramen*[6] transmitting a small vein to the superior longitudinal sinus (sometimes a small branch of the occipital artery).

FIG. 3.



Describe the internal surface.

This is concave, marked with numerous smooth *digital depressions* for the brain convolutions, with a shallow *half groove*[5] (Fig. 4) along the superior border for the ~~superior longitudinal~~ sinus, and

FIG. 4.



several *Pacchionian depressions*. At the anterior inferior angle a deep *groove*, branching in various directions, sometimes converted into a foramen, is found^{7}, for the middle meningeal artery, and at the posterior inferior angle a broad well-marked *groove*^{8} for part of the ~~lateral si~~nus.

How is it developed?

In membrane, from one centre at the eminence appearing at the fifth to sixth fœtal week.

Give muscular attachments, and articulations.

Temporal muscle: articulates with five bones, parietal, occipital, temporal, frontal, and sphenoid.

Occipital Bone.

Of what parts does this bone consist?

Of a curved plate of trapezoidal form, whose anterior blunted angle forms the *basilar process*^{1} (Fig. 5), and two lateral *jugular processes*^{3}, arising opposite the two *articular processes* called *condyles*^{12}, which lie on either side of a large opening, the *foramen magnum*^{11}; the long axes of the *condyles*^{12} converge toward each other from behind forward and inward.

Fig. 5.



What points should be noted on the external convex surface?

Midway between the summit and margin of the foramen magnum the prominent *external occipital protuberance*^{7} for attachment of the ligamentum nuchæ; a median *crest*^{8} extending from this to the margin of the foramen magnum; a pair of *superior curved lines*^{6} extending from the protuberance toward the lateral angles of the bone; a second pair of parallel *inferior curved lines*^{9}, starting about midway along the crest.

Describe the condyles.

They are two ~~kidney-shaped~~ convex articular surfaces on each side of the foramen magnum, having on their inner sides a rough *tubercle* for the check ligaments.

What foramina are situated near them?

The *anterior condyloid*^{2} (constant) for the hypoglossal ~~nerves~~, and *posterior condyloid foramina*^{10} (inconstant) for a small vein.

Fig. 6.



Describe the basilar process.

A strong quadrilateral plate in front of the foramen magnum with a median tubercular ridge-like *pharyngeal spine.* Its cerebral surface is smoothly *grooved* longitudinally for the ~~medulla oblon~~gata. At its junction with the jugular processes are the deep jugular *notches* forming part of each *jugular foramen* (foramen lacerum posterius).

Describe the cerebral surface of the bone.

It is deeply concave, divided into *four fossæ*[16-17] by a *crucial grooved ridge*[13-14-15], crossing at a prominent *internal occipital protuberance*[13]; the two lateral and superior grooves lodge, respectively, between the layers of the tentorium cerebelli and the falx cerebri, the lateral, and part of the superior longitudinal sinuses; the inferior groove, or *internal occipital crest*[14], has attached the *falx cerebelli*, and lodges in two grooves, the *occipital sinuses.* The upper surfaces of the jugular processes are deeply *grooved*[x] for the lateral sinuses.

What do the upper two fossæ lodge?

The posterior lobes of the cerebrum.

What do the lower fossæ accommodate?

The hemispheres of the cerebellum.

With what bones does it articulate?

Two parietal, two temporal, sphenoid, and atlas; ~~six~~ in all.

What centres of ossification has it?

One in membrane, at the occipital protuberance (eighth fœtal week), one for the basilar, and one for each condyloid portion in cartilage (seventh or eighth fœtal week): in four pieces at birth; in sixth year forms one bone.

What muscles are attached to the superior curved line?

Occipito-frontal, trapezius, and sterno-cleido-mastoid.

What muscles are attached between the curved lines?

Complexus and splenius capitis.

What muscles are attached below the inferior curved line?

Rectus capitis posticus major and minor, obliquus superior.

What muscles are attached to the basilar process?

The superior constrictor of the pharynx, the rectus capitis anticus major and minor.

What muscle is attached to the jugular process?

The rectus capitis lateralis.

The Sphenoid Bone.

What are its divisions and where is it situated?

A *body*^{2}, two *lesser wings*^{7}, two *greater wings*, two *pterygoid plates*, two *spinous processes*^{10}-^{11}, six *clinoid processes*, and three *minor prominences*; it is situated at the base of the cranium articulating with all the cranial bones, and many of the face. Named from a Greek word meaning *wedge*, in the process of development this is its function, serving to enlarge the cranial, nasal, orbital, and oral cavities at puberty.

Describe the body of the sphenoid.

It is large, cuboidal, hollowed out (in adult) into the *sphenoidal sinuses*^{20} (Fig. 8). Superiorly, in front, is a sharp projecting *ethmoidal spine*^{6} for the ethmoid; behind this, a smooth surface with a longitudinal eminence producing two shallow *olfactory grooves* on either side, bounding which behind is a transverse *optic groove*^{8} (Fig. 7), lodging the commissure of the optic nerves, terminating in two *optic foramina*^{12} (Fig. 7) for the optic nerves and ophthalmic arteries. Behind the groove is the projecting *olivary process*^{3} (Fig. 7) forming the anterior boundary of a marked depression, the *sella turcica*^{8} (Fig. 7), lodging the pituitary gland and circular sinus. On each side, in front, are two small projecting *middle clinoid processes*, while behind is a square-shaped projecting plate terminating at either angle by the tubercular *posterior clinoid processes*, and to them are attached portions of the tentorium cerebelli. The sides of this bony plate are *notched* for the sixth pair of nerves, and the superior surface, or *dorsum*^{2} (Fig. 7), with a broad shallow depression, slopes obliquely downward and backward, becoming continuous with the basilar portion of the occipital bone; it supports the pons Varolii. A broad *cavernous groove*^{4} (Fig. 7) for the carotid artery and cavernous sinus exists on either side of the body. The *posterior*[1] (Fig. 7) rough quadrilateral surface articulates with the basilar portion of the occipital bone, coössifying from the eighteenth to twenty-fifth year.

FIG. 7.



A vertical plate, the *rostrum*[21-22] (Fig. 8), projects from the *inferior surface* of the body forming part of the nasal septum, having the irregular openings of the *sphenoidal sinuses*[20] on either side (absent in child), which open either into the posterior ethmoidal cells or directly into the nasal cavities. Two thin-curved *sphenoidal turbinated*[20] bones narrow these orifices to a round opening at their upper part; these plates articulate with the palate and ethmoid bones.

The *inferior surface* presents a continuation of the *rostrum* for the vomer, having on either side a thin plate, the *vaginal process*[22], under which slips the alæ of the vomer, and the *pterygo-palatine grooves*, which by articulation with the sphenoidal processes of the palate bones are converted into the *pterygo-palatine canals* for the arteries and nerves of the same name.

Describe the greater wings.

Two strong processes arising from the sides of the body, curving upward, outward, and backward. The concave *cerebral surface*[14] of each wing forms part of the middle fossæ of the skull, presenting a circular opening at its antero-internal part, the *foramen rotundum*[14] for the second branch (sup. maxillary) of the fifth pair of nerves; a second larger ovoidal foramen, the *foramen ovale*[15] (Figs. 7 and 8), transmitting the third (inf. maxillary) division of the same nerve (sometimes the small petrosal nerve and small meningeal artery), while piercing the sharp, pointed, posterior extremity of the wing, or *spinous process*, is seen a short canal (sometimes double), the *foramen spinosum*[16] (Fig. 7), for the passage of the middle (great) meningeal artery. Occasionally, the *foramen Vesalii*, a small aperture to the inner side of the oval foramen, opposite to the root of the pterygoid process, transmitting a small vein, is detectable.

The *external surface* presents the *pterygoid ridge* dividing the temporal from the zygomatic fossa, and at its posterior part the *spinous process*[9] (Fig. 8), giving origin to the laxator and tensor tympani muscle, and the internal lateral ligament of the lower jaw.

FIG. 8.



The *anterior* or *orbital surface*, smooth, quadrilateral, forming part of the outer orbital wall, bounded above by a serrated edge for articulation with the frontal, below by a rounded border which enters into the formation of the *spheno-maxillary fissure*, internally aiding in the formation of the *sphenoidal fissure*[13], presents a serrated articular margin for the malar bone, and has a *notch* at the upper part of the inner border for a branch of the ophthalmic artery, a small *spine* at its lower part, for part of the origin of the external rectus muscle, and (at times) one or more *external orbital foramina*. The *circumference* has been described above.

Describe the lesser wings.

These[5] (processes of Ingrassias) two thin, triangular processes each arise from the upper and lateral parts of the body, project transversely outward, to terminate internally in the *anterior clinoid process*[x] (Fig. 7), externally in a sharp point. By their anterior borders they articulate with the orbital plates of the frontal bone, while by their posterior free margin they divide the anterior from the middle cerebral fossæ. Between their roots on each side is the *optic foramen*[12] (Fig. 7) for the optic nerve and ophthalmic artery. Their inferior surface forms on either side the back part of the orbital roof, and the upper boundary of the ~~*sphenoidal*~~ *fissure*[13] (Fig. 7), or foramen lacerum anterius, which is bounded internally by the sphenoid body, and inferiorly by the orbital margin of the greater wing, transmitting the third, fourth, ophthalmic branch of the fifth and sixth nerves, with the ophthalmic vein, filaments from the cavernous sympathetic plexus, also branches of the lachrymal and middle meningeal arteries, with a process of the dura mater.

Describe the pterygoid processes.

These, consisting of an *internal*[11] and *external*[10] plate separated behind by a notch—the *pterygoid fossa*—descend on each side perpendicularly from the junction of the greater wing with the body. At the base of the internal plate is the ~~*scaphoid fossa*~~, giving origin to the ~~tensor palati~~ muscle, above which is the orifice of the *Vidian canal*[17]; this plate terminates by a curved *hamular process*[11], around which plays the tendon of the tensor palati muscle. The external pterygoid plate, forming part of the inner wall of the zygomatic fossa, has arising from its inner surface the internal pterygoid, and from its outer the external pterygoid muscle. The *triangular notch* below has fitting into it the tuberosity of the palate bone.

Name the bones with which the sphenoid articulates.

The other seven cranial bones and five facial, viz., the two palate, two malar, and vomer.

By how many centres of ossification is this bone developed?

By ~~fourteen~~ centres: two for the internal pterygoid plates in *membrane*, two for the greater wings and external pterygoid plates, two for the lesser wings, two for the anterior part of the body, two for the sphenoidal turbinated bones, two for the posterior part of the body, two for the *lingula*, *i. e.*, posterior part of the outer margin of the cavernous groove, appearing from the second fœtal month to the third year in *cartilage*. From the tenth to the twelfth year the parts are all ossified, except a portion of the sphenoidal turbinates, which completely unite by the twentieth year.

What muscles have their origin from the sphenoid bone?

The temporal, external and internal pterygoids, superior pharyngeal constrictor, tensor palati, laxator tympani, levator palpebræ, obliquus superior, superior, internal, external, and inferior recti.

The Temporal Bones.

Where are they situated?

At the sides and base of the skull.

What parts does each bone present for examination?

A *squamous* (scale-like), a *mastoid* (nipple-like), and a *petrous* (stony hard) portion.

Describe the squamous portion.

Fig. 9.



The main portion is an irregularly semi-circular, smooth plate,[1] grooved posteriorly for the deep temporal arteries, convex externally, concave internally, and here grooved for the middle meningeal artery, with *digital depressions* for the cerebral convolutions. Posteriorly, a curved ridge (part of temporal) is seen. Springing from its lower part is the long, twisted, arched *zygomatic process*[2], arising by three roots, viz., a posterior, forming part of the temporal ridge, one running directly inward in front of the glenoid fossa, convex, covered with cartilage, forming the *eminentia articularis*[5], and a middle, the outer margin of the *glenoid fossa*[4], the obliquely transverse, ovoidal socket for the condyle of the lower jaw, divided into two parts by the *Glaserian fissure*[6] (Fig. 11), which transmits the laxator tympani muscle, the tympanic artery, and lodges the processus gracilis of the malleus: part of the parotid gland occupies that portion posterior to the fissure. Parallel to this fissure, in the retiring angle between the squamous and petrous portions, is the opening of the *canal of Huguier* for the chorda tympani nerve. At the junction of the articular eminence with the zygomatic process is the *zygomatic tubercle* for the external lateral ligament of the lower jaw.

Describe the mastoid portion.

This projects[3], rough, and perforated by numerous foramina (one large—the *mastoid foramen*[9]—transmits a vein to the lateral sinus, and a small artery), from the postero-inferior portion of the bone. The interior portion of the bone contains the *mastoid cells*, lined with mucous membrane continuous with that of the tympanum. The conical tip is termed the *mastoid process*[3x], upon whose inner side a deep grooved *digastric fossa*[26] (Fig. 11) exists for the digastric muscle, while parallel, but more internal, lies the *occipital groove*[27] (Fig. 11), lodging the artery of the same name. A deep, curved *groove*[13] (Fig. 10) exists on the inner surface of the mastoid portion for part of the lateral sinus, in which can be seen the inner orifice of the mastoid foramen.

Fig. 10.



Describe the petrous portion.

This is a dense, pyramidal mass, wedged in between the sphenoid and occipital bones, containing the organs of hearing. Its long axis is directed from without inward, forward, and a little downward. Its apex, three surfaces, and three borders must be studied. The *apex*[14] (Fig. 10) lies in the angle formed by the posterior border of the greater sphenoidal wing and the basilar process of the occipital, forms the posterior and external boundary of the foramen lacerum medium, and presents the internal orifice of the carotid canal.

The *base*, its upper half covered by the squamous and mastoid portions, presents the oval, expanded orifice of the *meatus auditorius externus*[8], its upper margin smooth, but surrounded for the greater part of its circumference by the rough-edged (for the cartilage of the external ear), curved plate of bone called the *auditory process*[8].

The *anterior surface* forming the posterior part of the middle cerebral fossa, presents an *eminence*[19] (Fig. 10) near the centre indicating the situation of the vertical semi-circular canal of the ear; external to this a *depression* over the tympanum; a shallow groove (sometimes double) leading backward and outward to an oblique opening, the *hiatus Fallopii*[20], for the petrosal branch of the Vidian nerve; a smaller opening (occasionally seen external to the latter) for the smaller petrosal nerve; the termination of the *carotid canal*[14] near the apex; above this canal a shallow *depression* for the Gasserian ganglion of the fifth nerve.

The *posterior surface* forms the anterior part of the posterior cerebral fossa, presents about its centre the large orifice of the short (about four lines) *meatus auditorius internus*[15] directed outward (closed at its bottom by a vertical perforated plate), which transmits the facial and auditory nerves (seventh and eighth pairs cranial), the auditory artery, and is lined by the dura mater; behind the meatus is a small slit leading to the *aquæductus vestibuli*[16], transmitting a small artery and vein, and lodging a process of the dura mater.

The *inferior* (basilar) *surface*, from apex to base, presents a rough quadrilateral surface for the origin of the tensor tympani and levator palati muscles; the circular orifice of the *carotid canal*[21] (Fig. 11) ascending vertically, then abruptly passing horizontally forward and inward for the passage of the internal carotid artery with its sympathetic nerve plexus; to the inner side of this —a small, triangular opening, the *aquæductus cochleæ*[17], transmitting a vein from the cochlea to the internal jugular; behind these openings a deep depression, the *jugular fossa*[24] (Fig. 11), which lodges the lateral sinus, and, with the jugular notch of the occipital bone, forms the *foramen lacerum posterius*, transmitting the jugular vein and ~~eighth~~ 9, 10, 11 pair of cranial nerves; a small *foramen for Jacobson's nerve*[23] in front of the bony ridge separating the carotid canal from the jugular fossa; on the outer wall of the jugular fossa a small *foramen for Arnold's nerve*[25]; posterior to the jugular fossa, a square-shaped, smooth facet (covered in the fresh state with cartilage), for articulation with the jugular process of the occipital; a broad, sheath-like, bony plate extending from the carotid canal to the mastoid process, the *vaginal process*[7], which splitting into two laminæ, encloses the base of a long, sharp spine, the *styloid process*[10] (Figs. 10 and 11), from which arise the stylo-pharyngeus, stylo-glossus, and stylo-hyoideus muscles, and the stylo-hyoid and stylo-maxillary ligaments; between this process and the mastoid, a large orifice, the *stylo-mastoid*[26] foramen, for the exit of the facial nerve, and the entrance of the stylo-mastoid artery; and, finally, between the vaginal and mastoid processes, the *auricular fissure* for the exit of Arnold's nerve.

FIG. 11.



At the angle formed by the junction of the petrous and squamous portions, two *canals*[28] exist, separated by a thin, bony septum, the *processus cochleariformis*, the upper lodging the tensor tympani muscle, the lower being the bony *Eustachian tube*[28].

What additional points of interest do the borders present?

The *superior* is *grooved* for the superior petrosal sinus, and has attached to it the tentorum cerebelli; the *posterior* has a *half-groove* which, with another on the occipital, accommodates the inferior petrosal sinus.

From how many centres is this bone developed?

From ~~ten:~~ one for the squamous and zygoma (in membrane), four for the petrous, two for the mastoid, two for the styloid, and one for the auditory process (all latter in cartilage), from eighth fœtal week to sixth fœtal month—bone coössifies during first year, except the styloid process.

With what bones does each temporal articulate?

The occipital, one parietal, sphenoid, one malar, and the i~~nferior~~ ~~max~~illary.

What muscular attachments has it?

Squamous, the temporal; *zygoma*, the masseter; the *mastoid portion*, the occipito-frontal, sterno-mastoid, splenius capitis, trachelomastoid, digastricus, and retrahens aurem. *Styloid*, the stylopharyngeus, stylo-hyoideus, stylo-glossus. *Petrous*, the levator palati, tensor tympani, tensor palati, and stapedius.

Ethmoid Bone.

Of what parts does this bone consist?

Of a *horizontal cribriform plate*[3] (sieve-like, whence name), of two *lateral masses*[10] of cells depending from this on either side, between which lies the *perpendicular plate*[2] forming part of the septum of the nose; the whole bone is of a cuboidal form.

Describe the cribriform plate.

Springing from the upper surface is the *crusta galli*[1] (cock's-comb), a vertical plate running from before backward, whose anterior border is notched to complete the foramen cæcum of the frontal bone. Two deep, longitudinal *olfactory grooves*, for the olfactory bulbs, lie on either side, their bottoms perforated by three rows of small foramina for filaments of the olfactory nerves, and on each side in front is a small slit for the nasal branch of the o~~phthalmic~~ nerve; the ~~crest~~ gives attachment to the ~~falx cerebr~~i.

Describe the perpendicular plate.

It is thin, usually deflected to one or other side, forms part of the nasal septum, and has upon each side numerous grooves and canals to accommodate the olfactory nerve filaments.

Of what are the lateral masses formed?

Of thin-walled cavities, the *ethmoidal cells*, whose outer limits are the vertical *orbital*[4] plates, while their inner walls form part of the nasal fossæ; above they are *grooved* to form with grooves on the frontal the two ethmoidal foramina. The upper outer margins of each mass present a number of half-cells completely closed in when articulated with the ethmoid notch of the frontal bone; posteriorly are also half-cells completed by the sphenoidal turbinated and palate bones; again, in front and below, the cells are completed by the lachrymal and superior maxillary bones; inferiorly an irregular hook-like plate projects, the *unciform process*[5], which closes in the upper part of the orifice of the antrum. A narr~~ow oblique~~ fissure, the *~~superior~~ ~~nasal~~ meatus*, subdivides the inner surface, bounded above by the thin-curved *superior turbinated bone*[6], below by the convex scroll-like *middle turbinated bone*[7]; the *~~posterior~~ cells* open into the upper part of this fissure, the *~~anterior~~* into the fron~~tal~~ sinus above, and below by the flexuous *infundibulum* they communicate with the *middle* nasal *meatus*, which is bounded above by the middle turbinated bone.

FIG 12.



How many centres of ossification has this bone?

Three; one for each lateral mass (fourth to fifth fœtal month), one for the perpendicular and cribriform plates in *cartilage* during the first year; the ethmoidal cells, completing the bone, form about the fourth or fifth year.

With how many bones does it articulate?

~~Fifteen~~: the sphenoid, two sphenoidal turbinated, frontal, two nasal, two superior maxillary, two lachrymal, two palate, two inferior turbinated, and the vomer.

Nasal Bones.

Describe them.

They are of oblong form, their *outer* surfaces concave from above downward, convex from side to side, marked by many small arterial furrows and usually presenting at their centres a single or double venous foramen. Their *inner* surfaces are convex from above downward and concave from side to side, traversed longitudinally by a *groove*, sometimes a *canal*, for a branch of the nasal nerve which escapes by a notch about the middle of the lower thin margin.

With what bones does each nasal articulate ?

With the frontal, ethmoid, its fellow nasal, and the superior maxillary. It has no muscular attachments.

How is each developed ?

From one centre in membrane (eighth fœtal week).

The Superior Maxillary Bones.

Describe them.

They each consist of a hollow body and four processes; together they form the whole upper jaw.

Describe the body.

It is cuboidal in form and hollowed out into a pyramidal cavity, *the antrum of Highmore*[11] (Fig. 14). Of its four *surfaces*, the *external* (facial), directed forward and outward, presents, just above the incisor teeth, the *incisive* (myrtiform) *fossa*[7] (Fig. 13), more externally the larger *canine fossa*[8] (Fig. 13), separated from the former by the *canine eminence* formed by the socket of the canine tooth; above this the *infra-orbital foramen*[9] (Fig. 13) giving egress to the infra-orbital artery and nerve, and, above all, the *margin of the orbit.*

The posterior surface is convex, forms part of the zygomatic fossa, has about its centre several apertures of the *posterior dental canals* for the dental vessels and nerves; at its lower part (especially after the wisdom tooth is formed) is a rounded eminence, *the maxillary* *tuberosity*[2] (Fig. 13), which articulates by its rough inner surface with the tuberosity of the palate bone; and just above the rough surface is an oblique *groove* running down on the inner side, which by the apposition of the palate bone forms the *posterior palatine canal*[15] (Fig. 14).

The *superior* (orbital plate) *surface*[5], thin, smooth, and triangular, forms part of the floor of the orbit, articulates internally by an irregular margin (from behind forward) with the palate bone, os planum of the ethmoid and lachrymal; externally, by its smooth, rounded edge, it helps to form the spheno-maxillary fissure (sometimes articulating in front with the orbital plate of the sphenoid), and in front, part of the orbital margin. Commencing at the middle of the outer border of this surface is the deep ~~*infra-orbital groove*~~[6] (Fig. 13), which in front becomes a *canal* of the same name, opening by the infra-orbital foramen; a small canal branching from this passes into the substance of the anterior wall of the antrum—the ~~*anterior dental canal*~~; at the inner forepart of this orbital surface is a minute *depression* for the origin of the inferior oblique muscle of the eye. ?

The *internal surface*, unequally divided horizontally into two parts by the *palate process*[12] (Fig. 14), forms above this, part of the outer wall of the nasal fossæ; below, a portion of the cavity of the mouth. The nasal surface presents a large irregular opening into the antrum, above which are irregular cellular cavities closed in, when articulated, by the ethmoid and lachrymal bones. Below the opening is the smooth surface forming part of the inferior meatus

FIG. 13.



FIG. 14.



of the nose; behind a *rough surface*[15] (Fig. 14) **for the palate bone with a groove passing from above downward and forward, the** *posterior palatine*[15] (Fig. 14), **which, closed in by the palate bone, forms the posterior palatine canal; in front, a deeper vertical** *groove*[10], **which by the apposition of the lachrymal and inferior turbinated bones forms the** *lachrymo-nasal duct*; **and still more anteriorly the well-marked horizontal, rough-ridged** *inferior turbinated crest*, **for the inferior turbinated bone.**

Describe the antrum of Highmore (maxillary sinus).

It is triangular, with apex outward formed by the malar process, its base the outer nasal wall, and opens, in the fresh state, by a small orifice of the size of a probe point into the middle nasal meatus. In the non articulated bone there is a large, irregular opening, which is narrowed when articulated by portions of the palate, ethmoid, and inferior turbinated bones. On its posterior walls are the *posterior dental canals*, while its floor presents several conical *prominences*, produced by the fangs of the first and second molar teeth; sometimes the floor is actually perforated. The walls are very thin.

Describe the malar process.

It is a rough, triangular eminence[4], concave on its facial and zygomatic surfaces, and roughly serrated above for the malar bone.

Describe the nasal process.

This is thick, triangular[3], rising upward, inward, and backward, with an external concave smooth surface, an internal surface articulating with the frontal above, and by a *rough surface* with the ethmoid, below this is the transverse ridge-like *superior turbinated crest* for the middle turbinated bone. Next comes a smooth, concave surface, part of the middle nasal meatus; then the inferior turbinated crest (see *ante*), and, most inferior of all, the concavity forming part of the inferior nasal meatus. The *anterior border* is thin, serrated for articulation with the nasal bone, and the *posterior border* thick, and grooved for the lachrymal duct, which is directed *backward and a little outward*; where the outer margin of the groove joins the orbital surface, is the *lachrymal tubercle*.

Describe the alveolar process.

This is thick, spongy, with eight cavities of varying depth, that for the canine tooth being deepest, those for the molars widest and subdivided.

Describe the palatine ~~process~~.

It is a horizontal plate[13] (Fig. 14), thickest in front, forms most of the roof of the mouth and floor of the nostril; has an upper and lower concave surface. In front, upon the upper surface, is the upper orifice of the *anterior palatine canal*, which conducts to a fossa formed between the two bones, transmitting the anterior palatine vessels through the *foramina of Stenson*, the ~~naso-palatine~~ nerves passing through the intermaxillary suture by the *foramina of Scarpa*; upon the under surface is a longitudinal *groove* (sometimes a canal), leading from the posterior palatine canal, for the posterior palatine vessels and nerve, while in front is seen the lower orifice of the *anterior palatine fossa* (foramen) presenting four openings; the inner border is thickened into a ridge on the upper surface forming the *nasal crest*, between which and its fellow fits the vomer, and which in front forms the *anterior nasal spine.*

How is this bone developed?

Probably by four centres; one for nasal and facial, one for orbital and malar, one for palatal in *membrane*, and one for incisive portion in *cartilage* at sixth to seventh fœtal week. Antrum commences at the fourth fœtal month.

With what bones does it articulate?

Frontal, ethmoid, nasal, malar, lachrymal, inferior turbinated, palate, vomer, and the other superior maxillary.

Give the muscular attachments.

Orbicularis palpebrarum, obliquus inferior of eye, levator labii superioris alæque nasi, levator labii superioris, levator anguli oris, compressor nasi, depressor alæ nasi, dilator naris posterior, masseter, buccinator, internal pterygoid, and orbicularis oris.

Lachrymal Bones.

What is their shape and where are they situated?

They are two small bones situated at the front part of the inner wall of the orbit, resembling in form, thinness, and size a ~~finger-n~~ail, hence called ~~*os unguis*~~. The external *orbital*¹ surface presents a vertical *ridge* giving origin to the tensor tarsi muscle, in front of which is a concave longitudinal *groove*² forming part of the *lachrymal duct* below, lodging the lachrymal sac above. Below is a hook-like process⁷ articulating with the inferior turbinated bone, assisting to form the lachrymal canal. A depressed furrow is seen on the nasal surface corresponding to the ridge externally; the surface in front forming part of the middle nasal meatus, the posterior closing in the anterior ethmoidal cells.

FIG. 15.



With what bones does it articulate?

With the frontal⁴, ethmoid⁵, superior maxillary⁶, and inferior turbinated.

How is it developed?

From one centre in membrane, at the eighth fœtal week.

What muscles arise from it?

The ~~tens~~or tarsi of Horner.

Malar Bones.

Describe them.

Two small q~~uadran~~gular bones placed at the upper outer part of face, forming the prominence of the cheek, parts of the orbit, temporal, and zygomatic fossæ. The *external surface*¹ is convex, smooth, perforated by one or more *malar foramina*², for the passage of nerves and vessels; it gives origin to the two zygomatic muscles. The *internal*⁶ concave surface has a rough triangular surface for the superior maxillary bone. It has superiorly a thick serrated-edged *frontal process*³ articulating with the external angular process of the frontal bone; a thick, strong plate-like *orbital* *process*, projecting backward, smooth and concave above, forming part of the floor and outer wall of the orbit, convex below, forming part of the temporal fossa, and bounding the spheno-maxillary fissure in front by its free margin, having upon its upper surface the orifices of one or more *temporo-malar canals* for filaments of the orbital branch of the superior maxillary nerve. Externally, projecting backward is the *zygomatic process*[7] articulating with the zygomatic process of the temporal bone by a serrated margin.

The *upper border*[4-5], smooth and arched, forms the outer and inferior margin of the orbit; the *lower border*[9], thick and rough, gives origin to the masseter muscle; the *anterior border*[6], roughly bevelled, articulates with the superior maxillary; while the *posterior border*[8], like an italic *f*, is continuous above with the temporal ridge, below with the upper border of the zygoma.

Fig. 16.



How is it developed?

From one centre (some say two), at the eighth fœtal week *in membrane.*

With what bones does it articulate?

With the frontal, sphenoid, temporal, and superior maxillary.

What muscles arise from it?

The levator labii superioris, zygomaticus major and minor, masseter, and temporal.

The Palate Bones.

Describe them.

These two bones, situated at the back part of the nasal fossæ, assist in the formation of the floor and ~~outer~~ wall of ~~the nose~~, the roof of the mouth, the floor of the orbit, the inner wall of the

antrum, and aid in forming the spheno-maxillary and pterygoid fossæ.

Of what parts does each bone consist?

Of a *vertical plate*[2], a *horizontal plate*[1], and three processes, the *pterygoid*[3], *sphenoidal*[9], and *orbital*[8].

Describe the points on the vertical plate.

This thin oblong plate, directed upward and a little inward, presents internally, at its upper part, the *superior turbinated crest*[2], lower a second ridge, the *inferior turbinated crest*[6] for the middle and inferior turbinated bones, between which lies the *groove* for the middle nasal meatus, and below the inferior crest another *groove* for the inferior meatus. The ~~*posterior border* articulates with the pterygoid process of the sphenoid;~~ on its external surface is a deep *groove* forming the posterior palatine canal by articulation with the superior maxilla, transmitting the vessels and nerve of the same name, while the upper and back *smooth surface* helps to form the inner wall of the spheno-maxillary fossa, the *anterior smooth* small lamina springing from the anterior border of the bone opposite the inferior turbinated crest—the *maxillary process*—narrows the orifice of the antrum, between which is the rough *maxillary surface.* The posterior surface at the lower portion is pyramidal, forming the *pterygoid process*[3], is deeply *grooved*[10], forming part of the pterygoid fossa, with a V-shaped, rough margin for articulation with the pterygoid plates.

FIG. 17.



The *superior border*, deeply notched by the *spheno-palatine foramen*[7] (or notch), forms the triangular hollow *orbital process*[8], in front, articulating *anteriorly* with the maxilla, *posteriorly* with the sphenoid (its cells here usually opening into the sphenoidal sinus), *internally*, with the ethmoid, and has a free *orbital* and *zygomatic* surface, the latter opening into the zygomatic fossa, with a *rounded border* forming part of the spheno-maxillary fissure; behind, projecting upward and inward is the *sphenoidal process*[9], *grooved* on its upper surface by a *pterygo-palatine groove* to help to form the same-named canal; articulating here with the sphenoid bone, externally and posteriorly with the pterygoid process, and having an internal concave surface forming part of the outer wall of the nasal fossa.

Describe the horizontal plate.

This, completing the nasal floor and hard palate, presents on its inferior surface a transverse *ridge* for the ~~tensor palati~~ aponeurosis; a deep *groove* assisting in the formation of the posterior palatine canal[11]; near this the orifices of several *accessory posterior palatine canals*; an *anterior border*, serrated and bevelled for the palate process of the superior maxillary; a smooth concave *posterior border*, for the attachment of the soft palate, terminating in the median line by a projection, which, with its fellow on the other bone, forms the *posterior nasal spine*[5], for the ~~azygos uvulæ mu~~scle; and an internal thickest border, whose upper edge is raised to form, with its fellow, a *crest*[4] articulating with the vomer.

How is this bone developed?

By one centre *in membrane* at the angle of junction of the two plates (seventh to eighth fœtal week).

With what bones does it articulate?

The sphenoid, ethmoid, superior maxillary, inferior turbinated, vomer, and opposite palate.

Give the muscular attachments.

Tensor palati, azygos uvulæ, internal and external pterygoids, and superior constrictor of the pharynx.

The Inferior Turbinated Bones.

Describe them.

They are two thin, scroll-shaped bones, extended horizontally along the outer walls of the nasal fossæ, just below the orifice of the antrum, and are attached to the inferior turbinated crests of the superior maxillary[1] and palate[2] bones, and present three processes for study, a *lachrymal*[5], in front of the upper border, forming part of the nasal duct by its junction with the lachrymal and superior maxillary bones; further back, the *~~ethmoidal process~~*[6], joining the ~~unciform process~~ of the ethmoid; and from the lower border of this process the *maxillary process*[3] projects, curving downward, and hooking over and narrowing the orifice of the antrum. The outer surface (from the median plane of the head) is concave; its ~~inner~~ convex, roughened, and grooved for arteries and veins.

FIG 18.



Name the bones with which it articulates, and its method of development.

It *articulates* with the ethmoid, lachrymal, palate, and superior maxillary bones; it is developed by one centre *in cartilage* (fifth fœtal month).

Vomer.

Describe it.

It is single, plowshare-shaped, vertically placed posteriorly in the nasal fossæ (usually deviated to one side), and forms part of

FIG. 19.



the nasal septum. Its *superior border*[1] is deeply *grooved* for the rostrum of the sphenoid, with lateral *wings* (alæ), which slip under the vaginal processes of the same bone. The *anterior border*[4] is grooved for the vertical plate of the ethmoid and the nasal septal cartilage. The *inferior border*[2] is received in the groove formed by the nasal crest of the superior maxillary and palate bones. The *posterior border*[3] is free, concave, thick above and thin below, and separates the nasal fossæ behind. On each lateral surface is the *~~naso-palatine~~ groove*[6] (sometimes a canal) for the nerve of the same name, and also small *furrows* for the lodgement of bloodvessels.

Name the bones with which it articulates, and its centres of ossification.

The sphenoid, ethmoid, both superior maxillary, and both palate bones. It has but one centre (eighth fœtal week *in membrane*), which includes between its two layers the septal cartilage; ossification is not completed until after puberty.

The Inferior Maxillary Bone.

Enumerate its general characteristics.

It consists of a *body*[1], with two *rami*[2], which are surmounted on either side by a curved beak-like *coronoid process*[8] in front, and an articular *condyloid process*[7] posteriorly, the constricted portion beneath which is termed the *neck* of the condyle; the depression separating these two processes is called the *~~sigmoid~~ notch*[9], crossed by the masseteric vessels and ~~nerve~~.

Describe the body of the bone.

It is horseshoe-shaped, and presents the following points for examination: the prominent triangular eminence in front, the *mental process*[3], extending upward from which is a median vertical ridge the *symphysis*[3], marking the junction of the two halves of the bone; on either side, just below the roots of the incisor teeth, is the *incisive fossæ*, giving origin to the levator menti; and more externally, below the root of the second bicuspid tooth, the *mental foramen*[6], for the exit of the mental artery and nerve. Extending outward from the base of the mental process is a distinct *ridge*[1], the *external oblique line*, continuous posteriorly with the anterior border of the ramus; to it are attached anteriorly the depressor labii inferioris and depressor ~~anguli~~ oris. The lower border[4], just in front

LANE LIB

of the attachment of the masseter, is *grooved* for the facial artery. The internal surface presents an indistinct *linear depression*, indicating the point of junction of the two halves of the bone, on either side of the centre of which are four prominent *genial tubercles*, disposed in pairs, sometimes blended into two, or even one irregular mass, the upper giving origin to the genio-hyo-glossus, the lower pair to the genio-hyoid muscles. Upon each side of these tubercles is the oval *sublingual fossa* for the glands similarly named, and beneath each a rough *depression* for the origin of the digastric muscle. Extending obliquely upward and backward from the sublingual fossa is the *internal oblique line* (mylo-hyoid ridge) for the same named muscle; beneath its edge is the oblong *submaxillary fossa*, for the submaxillary gland; above it the alveolar portion pitted with alveoli (sockets) for sixteen teeth in adult, ten in child.

FIG. 20.



Describe the rami.

These ascending, flattened, quadrilateral plates present in front the curved *coronoid process*[8], giving attachment to the temporal muscle; the deep *sigmoid notch*[9], forming whose posterior boundary is the *condyloid process*[7] surmounting its narrow *neck*, in front of which is a depression for the tendon of the external pterygoid muscle, and externally a small *tubercle* for the external lateral ligament. The articular surface is oblong, with its long axis transversely oblique from behind forward and outward, and is convex from behind forward, and from side to side. Upon the inner surface of the coronoid process commences a longitudinal *ridge*, with a *groove*, for the temporal above, the buccinator muscle below. The outer surface of the ramus has various *ridges* for the attachment of the masseter muscles; its inner surface presents a central oblique aperture, that of the *inferior dental canal* (communicating with each alveolus and terminating at the mental foramen) for the vessels and nerve; in front of this opening a prominent *ridge*, terminating behind in a sharp *spine* for the long internal lateral ligament; below the spine the *myo-hyloid groove* for the same named vessels and nerves, and behind the groove a *rough surface*, for the internal pterygoid muscle.

At the junction of the posterior and inferior margins of the ramus is the *angle*[5] marked with rough oblique ridges externally for the masseter, internally for the internal pterygoid muscle.

With what bones does the inferior maxillary articulate?

With the two temporals.

What ossific centres has it?

It is the second bone of the skeleton to ossify (clavicle first), and is probably developed by only two centres, chiefly in *membrane*, partly in *cartilage*, one for each lateral half, which coössify about the first year.

Compare the forms of the old and adult jaw.

In old age the alveolar portion being absorbed, the angle formed by the ramus with the body is very obtuse, and the dental canal is near the upper surface (important surgically in operating for neuralgia); in adult age the ramus is almost vertical, and the dental canal lies about the middle of the bone.

Give the muscular attachments.

Fifteen pairs; to the *ramus* the masseters, temporals, internal and external pterygoids; to the *inner surface* of the *body*, the genio-hyoglossus, genio-hyoid, mylo-hyoid, digastric, superior pharyngeal constrictor; to the *outer surface* of the body the depressor labii inferioris, depressor anguli oris, levator menti, orbicularis oris, platysma myoides, and buccinator.

The Sutures and Fontanelles.

What is a suture?

Rows of interlocking, tooth-like processes, projecting from the ~~external table~~ of either bone (the inner tables are merely apposed)

How many sutures are there?

~~Eighteen.~~

Name those at the vertex.

The interparietal (sagittal), the fronto-parietal (coronal), and the occipito-parietal (lambdoid).

Name those at the sides.

Spheno-parietal, squamo-parietal, and masto-parietal.

Name those at the base.

One baso-sphenoidal, two petro-occipital, two masto-occipital, two petro-sphenoidal, and two squamo-sphenoidal. The points of junction of the sphenoid with the frontal and ethmoid, and the frontal with the ethmoid, are not usually described as sutures.

Which only of the facial sutures has received a name?

The *transverse,* that extending from one external angular process of the frontal to the other, joining that bone with the malar, sphenoid, ethmoid, lachrymal, superior maxillary, and nasal bones.

Why does the skull consist of so many pieces?

To ~~admit of continuous growth~~ of the bones at their edges, pari passu with the growth of the brain, hence premature coössification of all the sutures results in *idiocy.* Obliteration of the sutures occurs at variable periods after maturity.

What are the fontanelles?

They are membranous intervals in the infant's skull, corresponding to the junction of the four angles of the parietal with the contiguous bones. They number six, viz., the *anterior,* of lozenge-shape, at the junction of the sagittal and coronal sutures, usually closed not later than the second year; the *posterior,* triangular, at the junction of the sagittal and lambdoid sutures, closed a few months after birth; and the *lateral,* one at the antero-inferior angle, another at the postero-inferior angle of each parietal bone, which are closed soon after birth.

How are these spaces closed?

By a gradual extension of the ossifying process, or by the development of ad~~ditional cen~~tres, forming so-called ~~*Wormian bones*~~.

The Cerebral Fossæ.

Describe the general characteristics of the anterior fossæ with their foramina [a].

Formed by the orbital plates of the frontal[1], the cribriform plate of the ethmoid with its spine[9], and the lesser wings[2] of the sphenoid, they are convex on either side and concave in the median line. The foramina are one median, the *foramen cæcum.* which, if pervious, transmits a vein from the nose to the superior longitudinal sinus; two *slits*[9] on each side of the crista galli for the nasal nerves; three rows of olfactory foramina on each side for the olfactory filaments; on the outer side of each olfactory groove, the *anterior and posterior ethmoidal foramina*, the former transmitting the anterior ethmoidal artery and the nasal nerve, the latter, the posterior ethmoidal artery and vein.

Give the boundaries and the foramina of the middle fossæ [b].

In front, they are bounded by the posterior margin of the lesser wings of the sphenoid[2], the anterior clinoid processes[11], and anterior margin of the optic groove; behind, by the upper borders of the petrous portion[5] of the temporal bones and basilar suture; externally by the squamous plates of the temporal[4] and anterior inferior angles of the parietals[7]. The *foramina* are on each side, the *optic*[14] for the optic nerve and ophthalmic artery. The *sphenoidal fissure* (foramen lacerum anterius), transmitting the third, fourth, ophthalmic branch of the fifth and the sixth cranial nerves, and filaments of the sympathetic, the ophthalmic vein, branches of the lachrymal and middle meningeal arteries with a process of the dura mater; immediately behind this, the *foramen rotundum*[15], for the superior maxillary division of the fifth cranial nerve; more posteriorly, the *foramen ovale*[16], for the inferior maxillary division of the fifth nerve, the

FIG. 21.



small ~~meningeal~~ artery and the ~~small petrosal~~ nerve; between the two, internally, the *foramen vesalii* (often absent), for a small vein; piercing the posterior inferior angle of the greater sphenoidal wing, the *foramen spinosum*[17], for the middle meningeal artery, meningeal veins, and sympathetic nerve branches from the cavernous plexus; on the inner side of the oval foramen the *foramen lacerum medium*[18] (filled in below by a plate of cartilage in the fresh state), for the internal carotid ~~artery,~~ carotid sympathetic ~~plexus~~, large petrosal nerve (Vidian), and a ~~small meningeal~~ branch from the ascending pharyngeal artery; on the anterior surface of the petrous portion of the temporal, the *hiatus Fallopii*[23], and beneath it, a small *foramen* for the small petrosal nerve.

What are the posterior fossæ [c]?

They are deeply concave, and each is formed by the occipital[8], petrous[5] and mastoid[6] portions of the temporal, and the posterior inferior angle of the parietal[7].

What foramina and other openings does each present?

The *meatus auditorius internus*[19] for the facial and auditory ~~nerves~~, and the ~~auditory artery~~; the *aquæductus vestibuli* for a small artery and vein; the *foramen lacerum posterius*[20] transmitting the glossopharyngeal, pneumogastric, and spinal accessory nerves, the internal jugular vein, and meningeal branches of the ascending pharyngeal and occipital arteries; the ~~*mastoid*~~ *foramen* (often absent) for a ~~vein~~; the *posterior condyloid foramen* (often absent) for a ~~vein~~ of the same name; the *anterior condyloid foramen*[21] for the ~~hypoglossal nerve~~, and a ~~meningeal~~ branch from the ascending pharyngeal artery: and separating the two fossæ in the median line, the *foramen magnum*[13], accommodating the medulla oblongata and membranes, the spinal accessory nerves and ~~vertebral arteries~~.

What additional foramina or openings are found at either side of the base?

The orifice of the Eustachian tube[33] admitting air to the middle ear.

The canal for the tensor tympani muscle[33] above the former.

The posterior orifice of the Vidian canal for Vidian nerve and vessels.

Glaserian fissure[15] for processus gracilis of the malleus, the laxator tympani muscle, the tympanic artery; and close by

The orifice of the canal of Huguier, transmitting the chorda tympani nerve.

The canal for Jacobson's nerve, the tympanic branch of the glosso-pharyngeal.

The aquæductus cochleæ, for a small artery and vein running to the cochlea.

The canal for Arnold's nerve, the auricular branch of the pneumogastric.

The auricular fissure, for the exit of Arnold's nerve.

The stylo-mastoid foramen[36], for the exit of the facial nerve, and the entrance of the stylo-mastoid ~~arte~~ry.

The *anterior*[26] and *posterior*[27] foramina of the palatal region have already been sufficiently described, and this region does not properly belong to the *base of the skull*, while those opening externally at the *base* and upon the *face* have been described sufficiently under the temporal bone, and the superior and inferior maxillary bones.

FIG. 22.



The Orbital Cavities.

Describe them.

They are two quadrilateral ~~pyramidal~~ cavities, with bases outward, their long axes directed from in front inward and backward, in such directions that if prolonged they would meet about the sella turcica of the sphenoid. ~~Seven~~ bones contribute to the formation of each, viz., the frontal[12], ethmoid[27], sphenoid[23], lachrymal[24], superior maxillary[6], palate[25], and malar[22]; they communicate with the *cranial cavity*, behind, by the optic foramina[18] and sphenoidal fissures[23], with the *nasal fossæ* through the lachrymo-nasal duct, and below, externally, by the spheno-maxillary fissure^{26} with the *temporal*, *zygomatic*, and *spheno-maxillary fossæ.*

FIG. 23.



What bones compose the roof, and what is its form?

The orbital plate of the frontal, anteriorly, the ~~lesser~~ sphenoidal wing behind; it is concave, and directed downward and forward, presenting internally a *depression*^{28} for the fibro-cartilaginous ~~pulley~~ of the superior oblique muscle of the eye, externally the ~~*lachrymal*~~ *fossa*^{12} for the gland.

Describe the floor, and of what bones formed.

Nearly flat, formed chiefly by the orbital plate of the superior maxillary^{6}, and, to a less extent, by the orbital processes of the malar^{19} and palate bones; it presents, just external to the lachrymal canal, a ~~*depression*~~ for the inferior ocular oblique muscle; externally, the *malo-maxillary suture;* near the middle the *infra-orbital groove;* and posteriorly, the *palato-maxillary suture.*

Give the bones forming, and the points of interest upon the inner wall.

It is flattened, formed by the nasal process of the superior maxillary, the lachrymal [24], os planum of the ethmoid [27] and sphenoidal body. It presents in front the *lachrymal groove* [20], bounded behind by the *lachrymal crest*; further back, respectively, the *lachrymo-ethmoidal* and *ethmo-sphenoidal sutures.*

What forms the outer wall?

The orbital plate of the malar and the greater wing of the sphenoid, and on it are seen the openings of one or two *malar canals* [19] and the *spheno-malar suture.*

Describe the chief points connected with the superior external angle of the orbit.

Posteriorly, the *sphenoidal fissure* (foramen lacerum anterius) [28] for the *entrance* of the third, fourth, ophthalmic branch of the fifth and sixth nerves, and branches of the lachrymal and middle meningeal arteries, a process of the dura mater, sympathetic nerve filaments, and the *exit* of the ophthalmic vein; also the *fronto-malar* and *fronto-sphenoidal sutures.*

What points does the superior internal angle present?

The suture between the lachrymal, ethmoid, and frontal bones; between the junction of the two latter bones, the *anterior ethmoidal foramen* [2] for the anterior ethmoidal artery and nasal nerve, and the *posterior ethmoidal foramen* [17] for the posterior ethmoidal artery and vein.

What points does the inferior external angle present?

The *spheno-maxillary fissure* [26] transmitting the infra-orbital vessels and superior maxillary nerve, the ascending branches from the spheno-palatine ganglion, and the orbital branch of the superior maxillary nerve.

How is the inferior internal angle formed?

By a *suture*, the union of the lachrymal and os planum of the ethmoid with the superior maxillary and palate bones.

What foramen does the orbital margin present?

The *supra-orbital* [8] at the junction of the inner and middle thirds, transmitting the supra-orbital artery, veins, and nerve.

What foramen opens at the apex of the orbit ?

The *optic*[18], ~~between the two roots of the lesser wing~~ of the sphenoid, transmitting the optic nerve and the ophthalmic artery.

Recapitulate the openings communicating with the orbit.

The optic[18], sphenoidal fissure[23] (foramen lacerum anterius), spheno-maxillary fissure[26], infra-orbital canal[4], anterior[2] and posterior[17] ethmoidal foramina, malar foramina[19], supraorbital foramen, and lachrymal canal[20] (occasionally, in addition, one or more external orbital foramina).

The Nasal Fossæ.

Describe them.

They are two irregular cavities, extending from the base of the cranium above to the roof of the mouth below[17], separated in the median line by a thin osteo-cartilaginous septum, opening upon their facial aspect by two large apertures, the *anterior nares*[4, 10], and into the pharynx by the *posterior nares*[15]. Each fossa communicates with four sinuses and four cavities.

Of what parts is the nasal septum composed ?

In front, the crest of the nasal bones, and the frontal nasal spine; its middle portion, the vertical plate of the ethmoid; behind, the rostrum of the sphenoid and the vomer; below, the crests of the superior maxillary and palate bones.

What points does the roof of each fossa present ?

In front, the *slit* for the nasal nerve; numerous *foraminæ* for the olfactory filaments; most posteriorly the opening of the sphenoidal sinus[24].

The floor ?

In front, the *anterior nasal spine*[10]; behind this the upper opening of the *anterior palatine canal*[14]; internally the *nasal crest*[12-17] of the superior maxillary and palate bones.

Describe the chief points of interest in the outer wall of each fossa.

This presents three irregular longitudinal passages, formed by three projecting bony plates, called the *superior* [19], *middle* [20], and *inferior meatuses* [21].

Describe each meatus.

The *superior* [19] occupies the posterior third of the wall, lies between the superior[6] and middle[7] turbinated bones (processes of the ethmoid), and has opening into it two *foramina*, the *spheno-palatine* [25] posteriorly, the *posterior ethmoidal* cells in front part of the upper wall.

FIG. 24.



The *middle* [20] occupies the posterior two-thirds of this wall, lies between the middle and inferior turbinated bones, and has opening into it, in front, the *infundibulum*; in its centre, the *antrum* [26].

The *inferior* [21] lies between the inferior turbinated bone and the nasal floor, extends along the whole length of the outer wall, and has opening into it, in front, the lower orifice of the lachrymo-nasal duct (and the anterior palatine canal in the macerated bone, *not* in the natural state).

Describe the position and boundaries of the temporal fossa.

Situated at the lateral region of the skull, each fossa is marked out upon the skull-cap by the *temporal ridge,* which extends in a curved line first upward and backward from the external angular process of the frontal bone, then downward behind to form the posterior root of the zygomatic process. Its anterior boundaries are the frontal, malar, and greater wing of the sphenoid, above and behind the temporal ridge, below the pterygoid ridge on the greater wing of the sphenoid, externally the zygomatic arch; it opens below into the zygomatic fossa, and is filled by the temporal muscle, and is traversed by grooves, for branches of the deep temporal artery.

Describe the zygomatic fossa, its boundaries, and communicating fissures.

It is bounded, *anteriorly*, by the tuberosity of the superior maxilla, and the ridge descending from its malar process; *superiorly*, by the pterygoid ridge of the greater sphenoidal wing and squamous plate of the temporal; *behind*, by the posterior border of the pterygoid process; *below*, by the alveolar border of the superior maxilla; *internally*, by the external pterygoid plate; and *externally*, by the ramus of the lower jaw and the zygoma. The *spheno-maxillary* and *pterygo-maxillary fissures* open into its inner upper part.

What does it lodge?

The internal and external pterygoid and part of the temporal muscle, the internal maxillary artery and inferior maxillary nerve and their branches.

Describe the spheno-maxillary fissure.

It runs horizontally, opens into the outer back part of the orbit, lying between the lower orbital border of the greater wing of the sphenoid and the outer border of the orbital plate of the superior maxillary and a small part of the palate bone; externally is a small part of the malar, while, *internally*, it joins the pterygo-maxillary fissure at a right angle; it transmits the infra-orbital artery, the superior maxillary nerve, with its orbital branch, ascending branches from Meckel's ganglion, and serves to connect the orbit with the spheno-maxillary, temporal, and zygomatic fossæ.

What are the boundaries of the pterygo-maxillary fissure, and what passes through it?

It is bounded, in front, by the maxillary tuberosity; behind, by the pterygoid plate of the sphenoid; descends at right angles from the inner extremity of the spheno-maxillary fissure, transmits branches of the internal maxillary artery, and connects the zygomatic and spheno-maxillary fossæ.

Describe the spheno-maxillary fossa.

Situated at the junction of the spheno-maxillary and pterygo-maxillary fissures, it is bounded, *above*, by the under surface of the body of the sphenoid, and by the orbital process of the palate; in *front*, by the superior maxillary bone; *behind*, by the pterygoid process of the sphenoid; *internally*, by the vertical plate of the palate. In this fossa terminate the sphenoidal, spheno-maxillary, and pterygo-maxillary fissures. The *orbital*, *nasal*, and *zygomatic fossæ* communicate with it, and also the *cranial* cavity.

How many foramina open into it?

Five (sometimes seven or eight); three in the back wall, viz., above, the *foramen rotundum*; more internal and inferior, the *Vidian*; and lowest and most internal, the *pterygo-palatine*; on the inner wall is seen the *spheno-palatine foramen*; below, the upper orifice of the *posterior palatine canal*, and sometimes two or three *accessory posterior palatine canals*.

BONES OF THE TRUNK.

The Vertebral Column.

How are the vertebræ divided?

By regions, viz., into cervical (7); dorsal (12); lumbar (5); sacral (5); and coccygeal (4); 33 in all.

What are the sacral and coccygeal vertebræ called to distinguish them from the remaining bones?

False, the others being called true (false described with pelvic bones).

What parts are common to all vertebræ?

A *body*[1]; posteriorly on each side a *pedicle*[2], supporting two *laminæ*[3], which, joining behind enclose the *spinal foramen*[4]; from the junction projects a *spinous process*^{5}: from other parts six additional *processes* arise, viz., two *transverse*^{6}, and four *articular*$^{7\,7'}$.

Describe a vertebral body.

Composed of cancellous bone, with a thin, compact layer externally, its sides are concave from above downward, while its *upper surface* is, in the *cervical* region, concave laterally, forming a lip at either side, while the *lower surface* is convex from side to side, and concave from before backward, forming an anterior lip; in the *dorsal* region it is flat above and below; and in the *lumbar region*, flattened, or slightly concave above and below. Articular facets and demi-facets for the heads of the ribs, mark the *dorsal* bodies, neither the lumbar nor the cervical bodies having these.

Describe the pedicles.

They project backward in all but the cervical vertebræ, where they are directed obliquely outward. They present *intervertebral notches* above and below (deepest *above* in the cervical vertebræ; *below* in the dorsal and lumbar spine), forming, when articulated, *intervertebral foraminæ* for the exit of the spinal nerves, and the entrance of vessels.

Fig. 25.



What are the laminæ ?

Two broad plates closing in the spinal foramen, roughened at their upper margins and their inferior internal surfaces for the ligamenta subflava.

Describe the transverse processes.

They are bifid in the cervical and perforated by the vertebral foramen for the vertebral vessels; thick, strong, and long, with anterior articular facets, in the dorsal; long and slender in the lumbar: they spring from the junction of the pedicle and lamina, and also from the side of the body in the *cervical* region.

Describe the articular processes.

Two are superior and two inferior, projecting from the junction of the laminæ and pedicles. The upper pair look upward and backward in the cervical region; backward and outward in the dorsal, inward and slightly backward in the lumbar; the lower pair are exactly the reverse of the upper in each region.

Describe the spinous processes.

Bifid, short, and horizontal in the cervical; long, triangular, directed obliquely downward in the dorsal; and thick, broad, quadrilateral in the lumbar.

Describe the spinal foramen.

It is largest and triangular in the cervical, smallest and round in the dorsal, medium and triangular in the lumbar.

Name the peculiar vertebræ of each region, and describe them.

The first cervical, or atlas; the second cervical, or axis; the seventh cervical, or vertebra prominens; the first, ninth, tenth, eleventh, and twelfth dorsal, and the fifth lumbar.

The atlas supports the head, and is formed of two *lateral masses* joined by an *anterior* and *posterior arch*; the former presents a *facet* posteriorly for the odontoid process. Upon the upper surface of each lateral mass is an *articular facet*, looking upward, inward, and backward for the occipital condyles, while the *inferior facet* looks downward and directly inward; a small tubercle represents the spinous process. *Development* is by one centre for anterior arch (first year); one for each lateral mass (sixth fœtal week).

The *axis* has surmounting the body the *odontoid process*, with a *facet* in front for articulation with atlas, another behind for the transverse ligament, with a *roughened apex*, to which are attached the check ligaments; on each side of the odontoid, facing upward and outward, are the superior articulating processes. *Development* is like other vertebræ, except three additional centres for odontoid, two appearing at the sixth fœtal month.

The vertebra prominens is so called because of its long, spinous process, to which is attached the ligamentum nuchæ. *Development*, like other vertebræ, except one additional centre for anterior part of transverse process.

The dorsal vertebræ.

The first has an *entire facet* and a *demi-facet* on body; the ninth has a *demi-facet* on body above, and a *facet* on the transverse process; the tenth has *one facet* on the body and *one* on the transverse process; the eleventh and twelfth, *one facet* on the body, none on the transverse processes, the latter also closely resembling those of a lumbar vertebra.

The fifth lumbar has a markedly wedge-shaped body, with the base forward.

How are the ordinary vertebræ developed?

By one centre for the body, one for each lamina (sixth to eighth fœtal week); at sixteen years, one centre for the tip of each transverse, and two for the spinous process; and at twenty-one years, a plate upon the upper and lower surface of the body; the lumbar vertebræ have two additional centres tipping the superior articular processes; coössification at thirty years.

The Thorax.

Give its structure, form, and boundaries.

It is conical in form, and its osseo-cartilaginous framework is formed by the dorsal vertebral bodies behind, the ribs laterally, and the costal cartilages and sternum in front; its base is formed by the diaphragm. Through its apex, the great cervical vessels, the pneumogastric, phrenic, and sympathetic nerves, the trachea, œsophagus, and thoracic duct pass (it is also said, the apices of the lungs during inspiration).

What are the most important structures this cavity contains?

The trachea, primitive bronchi and lungs, the heart, aorta and its primary branches, the internal mammary arteries, the venæ cavæ, bronchial and azygos veins, the pneumogastric, phrenic, and splanchnic nerves, the œsophagus, thoracic duct, lymphatic vessels and glands.

Hyoid Bone.

Describe this bone and its development.

It consists of a body and two *greater* and two *lesser cornua*; the *greater cornua* project backward from the lateral surfaces of the body, and have attached to their tubercular ends the thyro-hyoid ligaments; the *lesser cornua*, attached to the junction between the body and greater cornua, give attachment, by their apices, to the stylo-hyoid ligaments; it is *developed* by five centres: one for body, and one for each greater horn toward the end of fœtal life; one for each lesser horn some months after birth.

Give the muscular attachments.

The sterno-hyoid, thyro-hyoid, omo-hyoid, digastric, stylo-hyoid, mylo-hyoid, genio-hyoid, genio-hyo glossus, hyo-glossus, middle pharyngeal constrictor, and sometimes the lingualis.

The Sternum.

Describe it.

The breast-bone consists of three segments, viz., the manubrium (handle), the gladiolus (sword), and the ensiform (xiphoid) cartilage. Upon the upper border of the *manubrium*[8] is the *inter-clavicular notch*[11]; upon either side of which is the *facet*[12] for the clavicle; lower, another for the first rib; below, a *demi-facet*[2] for the second rib.

The gladiolus[9] has a *demi-facet*[7] above for the second rib, another below for the seventh[7], and, between, *facets* [3], [4], [5], [6], for the third fourth, fifth, and sixth ribs.

The ensiform cartilage[10] has a *demi-facet*[7] above for the cartilage of the seventh rib.

How is this bone developed?

By six centres; one for manubrium, four for gladiolus, one for ensiform cartilage (fifth fœtal month to eighteen years); the three pieces rarely coössify.

FIG. 26.



What muscles are attached to this bone?

Above, the sterno-cleido-mastoid, the sterno-hyoid, and sterno-thyroid; below, the rectus abdominis, external and internal oblique, transversalis, and diaphragm; in front, the pectoralis major; behind, the triangularis sterni.

Ribs.

What is their number, and how are they divided?

Twelve on each side, seven of which are *true*, or those articulating with the sternum by a separate cartilage; five *false*, three of which indirectly articulate through the medium of the seventh cartilage; while the two lowest, having their anterior extremities free, are called *floating ribs.*

Of what parts does a rib consist ?

A head, neck, shaft, angle, and tuberosity, except the eleventh and twelfth ribs.

Describe these parts.

The *head*[1] (except first, tenth, eleventh, and twelfth ribs having only one facet) is divided into two facets by a ridge for the inter-articular ligament. The *neck*[2] is flattened, *roughened* upon its upper border for the anterior, and upon its posterior surface for middle costo-transverse ligaments; in front it is smooth. The *tuberosity*[3] (absent in eleventh and twelfth), placed at the base of the neck, presents a facet for the transverse process of the next lower vertebra, and a *rough surface* for the posterior costo-transverse ligament. The *shaft*[5] is twisted on its long axis (except first and second), externally convex, its upper border rounded, its lower *grooved*[4] for the intercostal vessels and nerve, and presents at its anterior extremity an *oval depression*[6] for the costal cartilage. The *angle*[4], at a variable distance in front of the tuberosity, is indicated by a *rough line*.

FIG. 27.



How are the ribs developed ?

By three centres (except eleventh and twelfth, two centres), one each for head, shaft, and tuberosity (for shaft sixth fœtal week); for head and tuberosity (sixteen to twenty years); coössifies at twenty-fifth year.

Describe the peculiar ribs.

They are the first, second, tenth, eleventh, and twelfth ribs.

First rib [c], broad, short, flat, one facet on head, angle absent; on upper surface two parallel grooves, the anterior for the subclavian vein, the posterior for the artery, and between them a *tubercle*[7] for the anterior scalene muscle a *sure guide to the subclavian artery.*

Second rib, also flattened, the tuberosity and angle nearly coincide, and presents near its middle a *rough eminence,* for the attachment of part of the first, and all of the second digitation of the serratus magnus.

Tenth rib [d], one facet on head.

Eleventh rib, one facet on head, no tuberosity, no neck, slight angle.

Twelfth rib, one facet on head, neither neck, angle, nor tuberosity.

The Pelvic Bones.

What are they?

The ossa innominata, the sacrum, and the coccyx.

Describe the innominate bones.

They are formed by the union, about puberty, of three bones, the ilium[1], ischium[2], and pubes.[3] At their point of junction is the *acetabulum*[19] (cotyloid cavity) for the head of the femur—the pubes forming one-fifth, the ilium and ischium each about two-fifths. A rough central depression at the bottom of the acetabulum lodges a vascular mass of fat covered with synovial membrane, while the anterior lower margin is interrupted by the *cotyloid notch,* bridged across by the transverse ligament, beneath which passes the nutrient vessels and ligamentum teres, which latter arises from either side of the notch externally; the whole margin has attached to it the cotyloid ligament. Between the pubes and ischium, on the anterior surface, is the *obturator foramen*[20] (thyroid), closed by the membrane of the same name, except above externally where the obturator vessels and nerve escape; the foramen is a large ovoidal opening in the male, a smaller and triangular one in the female.

Describe the ilium.

This is the upper expanded portion of the bone, presenting along its upper border the *crest*[4] with its outer and inner lips, terminating in front by the *anterior superior spinous process*[8], giving origin to Poupart's ligament, the sartorius and tensor vaginæ femoris muscles, and behind by the *posterior superior spinous process*[10], to which are attached part of the erector spinæ muscle and the oblique band of the sacro-iliac ligament. Below both the anterior[8] and posterior spines[10] is a process called, respectively, the *anterior*[9] (for straight tendon of the rectus femoris muscle) and *posterior inferior spinous*[11] (for great sacro-sciatic ligament); by these projections a *notch* is formed in front and behind, the former giving partial origin to the sartorius muscle and transmitting the external cutaneous nerve. Between the posterior inferior spine and the spinous process of the ischium is situated the *greater sacro-sciatic notch*[13], giving egress to the pyriformis muscle, the greater and lesser sciatic, superior gluteal and pudic nerves, also one to the obturator externus muscle, and the sciatic, gluteal and pudic vessels. About two inches from the posterior superior

FIG. 28.



FIG. 29.



spine, passing downward and outward from the crest, is the *superior curved line*[7], from the surface back of which arise the gluteus maximus muscles and a few fibres of the pyriformis; about an inch behind the anterior superior spine, passing from the crest downward and backward to the upper part of the great sacro-sciatic notch, is the *middle curved line*[5], the gluteus medius arising from the space between these two lines; passing downward and backward, from the upper part of the anterior inferior spine to the front of the sacro-sciatic notch, is the *inferior curved line*[6], between which and the middle arises the gluteus minimus. Above the acetabulum is a *groove* for the reflected tendon of the rectus femoris muscle. The inner surface presents the large, smooth concave *Venter* (internal iliac fossa)[1] (Fig. 29), limited below by the prominent *ilio-pectineal line*, and behind the iliac fossa is the rough *auricular surface*[9], the lower part for articulation with the sacrum, the upper for the posterior sacro-iliac ligaments.

Describe the ischium.

It consists of a body, tuberosity, and ramus, and forms the lowest part of the innominate bone. The external surface of the *body*[2] forms two-fifths of the acetabulum; below this is a *groove* for the obturator externus tendon; its inner surface forms the lateral boundary of the true pelvis; from the posterior border, below the centre, projects the *spine of the ischium*[19] (Fig. 29), above and below which are the *greater*[22] and *lesser sacro-sciatic notches*[23], the latter giving egress to the obturator externus muscle and its nerve, and ingress to the pudic vessels and nerve. The lowest portion presents a *tuberosity*[18], with an outer and inner lip—to the latter being attached the greater sacro-sciatic ligament; above is a *groove* for lodgement of the internal pudic vessels and nerve. Passing upward and inward from the tuberosity to join the ramus of the pubes, and bounding the obturator foramen in front, is the thin, flattened *ascending ramus*[17].

Describe the pubes.

This bone (also called pectineal) consists of a *body*[2] or *horizontal*[12] and a *descending* (perpendicular) *ramus*[13]. The outer end forms one-fifth of the acetabulum; above, a rough *ilio-pectineal* *eminence*[20] indicates the point of junction with the ilium; the inner end is the oval *symphysis*[11] with eight or nine *ridges* for attachment of the fibro-cartilage; the upper triangular surface presents posteriorly the pectineal portion of the *ilio-pectineal line*[16]; the anterior surface presents the *crest*[14], ending externally in the *pubic spine*[15] giving attachment to Poupart's ligament, internally in the *angle*; below is a *groove* for the obturator vessels and nerve; the *descending ramus*[13], thin and flat, joins that of the ischium, completing the anterior boundary of the obturator foramen.

How are these bones developed?

By three primary centres, one for each bone (from the eighth fœtal week to fifth fœtal month); and one for the crest, one for the tuberosity, one for the anterior inferior spine, one for symphysis, and one Y-shaped, joining the three pieces forming the acetabulum, appearing about puberty; bone coössifies completely about twenty-fifth year. (For order of junction, see Gray.)

What muscular attachments has the innominate bone?

Those of the abdomen, some of the thigh, those of the perineum, pelvic floor and rotators of the thigh.

Describe the sacrum.

It is composed of five consolidated vertebræ, is of triangular form with broad base and blunted apex and lateral expanded masses or alæ; its anterior surface is concave, its posterior convex, and with the coccyx it forms the posterior wall of the true pelvis. Its anterior surface is marked by four transverse *ridges*[1], indicating the lines of junction of the segments; eight *anterior sacral foramina*[2] with broad shallow *grooves* for the anterior sacral nerves open on this surface; the point of junction with the last lumbar vertebra forms the *promontory* (sacro-vertebral angle); and upon each side are the *alæ*[7]—the expanded portions of the bone. Posteriorly are three or four median tubercles (rudimentary spinous processes); externally are the laminæ, those of the fifth and sometimes the fourth being deficient; outside these is a row of rudimentary articular processes. On each side of the spine is a broad *sacral groove,* lodging the origin of the erector spinæ muscle; externally are the four *posterior sacral foramina* on each side; at the posterior inferior portion of the bone are the two *cornua*—articulating surfaces for the coccyx; each lateral surface has on its anterior upper part an *auricular surface*[4] for articulation with the ilium; on each side of the apex below is a deep *notch,*[10] [11], for the fifth sacral nerve; the *base*[8] resembles the upper surface of a lumbar vertebra, with the last of which it articulates; the *apex*[9], directed downward and forward, has an oval concave articular surface for the coccyx; the *sacral canal* runs the whole length of the bone, triangular above, small and flattened below and deficient in its posterior wall at the lower part; it lodges the sacral nerves, and into it open the anterior and posterior sacral foramina.

FIG. 30.



FIG. 31.



What centres of ossification has the sacrum?

Thirty-five, appearing from the eighth fœtal week to the twentieth year, coössification being complete from the twenty-fifth to thirtieth year. The bodies have each three centres; each lamina one centre; the lateral masses three centres each; the lateral surfaces two each.

Give the muscular attachments.

The pyriformis, coccygeus and iliacus in front, the gluteus maximus, latissimus dorsi, multifidus spinæ, erector spinæ, and sometimes the extensor coccygis behind.

Describe the coccyx.

Usually composed of four rudimentary vertebræ, more or less coössified, it forms a triangular bone whose *base*[1] articulates with the sacral apex. The first piece presents two *cornua*[2], projecting upward from either side of the base for articulation with the sacral cornua, their junction completing the fifth sacral foramina for the posterior branches of the fifth nerves. The *apex*[6] is rounded and occasionally bifid or turned to one side; two rudimentary *transverse processes*[3] are seen on the first piece.

Describe its development.

From four centres: one for each segment, the first piece commencing at birth; second, five to ten years; third, ten to fifteen years; fourth, fifteen to twenty years; coössification varies as to time and manner.

Give the muscular attachments.

Laterally the coccygei; behind the gluteus maximus and extensor coccygeus (when present); apex, sphincter ani; in front levator ani.

The Pelvis.

Describe the pelvis.

Formed by the two innominate bones, the sacrum and the coccyx, all above the ilio-pectineal line is called the *false pelvis*, consisting of the two *iliac fossæ*; all below, the *true pelvis*.

Describe the true pelvis.

Its *brim*, or inlet, somewhat heart-shaped, is formed by the linea ilio-pectinea at the sides, completed in front by the spine and

FIG. 32.



crest of the pubes, behind by the anterior margin of the base of the sacrum and promontory of the sacrum. Its average diameters in the female are, antero-posterior^{d}, four and one-half inches; transverse^{tr}, five and a quarter inches; the oblique^{ob}, five inches; its long axis, if extended, would pass from the middle of the coccyx to the umbilicus; in the male these measurements are diminished by at least one-half inch.

Describe the cavity.

This is bounded in front by the symphysis pubis, behind by the concavity of the sacrum and coccyx, on either side by the broad, smooth, quadrangular inner surface of the body of the ischium, forming a curved canal wider in the middle than at its outlets, measuring in depth at the symphysis one and one-half inches, three and one-half inches in the middle axial line, and four and one-half inches posteriorly, perhaps as much as five and one-half inches in males.

Describe the lower circumference of the pelvis.

This is called the *outlet*, is bounded on each side by the tuberosities of the ischium, the pubic arch in front and tip of the coccyx behind. Its diameters are four and one-quarter to four and three-quarters inches transverse, antero-posterior and oblique four and one-half increased to five by pressure on the coccyx; in the male, the diameters average three and one-half inches.

What are the chief differences between the male and female pelves?

The strength of the bones, distinctness of the muscular impressions, the depth and narrowness of the cavity, and large obturator foramina mark the *male pelvis*: the lighter bones, broader iliac fossæ, the less-curved sacrum, the wider pubic arch, and the universally greater diameters, distinguish the *female pelvis*.

Bones of the Upper Extremity.

Name the bones.

The clavicle, scapula, humerus, radius, ulna, scaphoid, semilunar, cuneiform, pisiform, trapezium, trapezoid, os magnum, unciform, five metacarpal, and fourteen phalangeal bones.

Describe the clavicle.

It is a long bone, curved like the italic letter *f*, its outer third flattened from above downward, and concave anteriorly; the inner two-thirds are cylindrical and convex anteriorly, and it extends almost horizontally between the sternum and scapula, the two extremities being respectively termed the *sternal*[1] and *acromial*[2].

FIG 33.



Describe the chief points presented by the clavicle, commencing at the outer extremity.

The upper surface has *impressions*, that in front for the deltoid,[7] that behind for the trapezius; at the outer end is a *facet*[2] articulating with the acromium process of the scapula; at the posterior border, beneath, is the *conoid tubercle*,[4] just above the coracoid process of scapula, for the conoid ligament; extending from the tubercle, forward and outward, nearly to the outer end of the anterior border is the *oblique line* for the trapezoid ligament; occasionally at the centre of the anterior border of the outer third is the *deltoid tubercle*. The under surface of the middle portion presents the *subclavian groove*[3] for the same named muscle—here appears the *nutrient foramen* directed outward; the inner third of the superior border bears an *impression* for the sterno-mastoid muscle[5] (A, Fig. 33), while the inner half of the anterior margin has another *impression* for the pectoralis major muscle. Beneath the posterior border of the sternal end is the *rhomboid impression*,[5] for the rhomboid or costo-clavicular ligament. The *sternal end*[1] is triangular in form, its inner surface for articulation with the sternum, this surface being continuous with a facet beneath, for articulation with the first costal cartilage.

Give its development.

By two centres; one for the shaft earliest in the body (thirtieth day); one for sternal end (eighteenth to twentieth year); uniting about the twenty-fifth year.

Give the muscular attachments.

Sterno-cleido-mastoid, trapezius, pectoralis major, deltoid, subclavius, sterno-hyoid, and platysma.

Describe the scapula.

The shoulder-blade, a large, flat, triangular bone, extends from the first to the eighth ribs on the back and side of the thorax, its posterior margin lying nearly parallel to, and one inch from the spinous processes of the vertebræ.

The *venter* (anterior surface) or *subscapular fossa*^{1} is concave, presents some transversely oblique ridges $^{1}, ^{1}, ^{1}$, for the tendinous intersections of the subscapular muscle, and a *marginal surface*^{15} along the inner border—triangular above and below, linear between —for the serratus magnus muscle. The *angle* is the transverse depression at its upper part where the fossa is deepest.

FIG. 34.



FIG. 35.



The *dorsum*^{2} (posterior surface) (Fig. 35) is divided by a prominent bony ridge, the *spine*^{10}, affording attachment to the deltoid and trapezius muscles and ending in the *acromion process*[12], into the *supra-*[1] and *infra-spinatus*[2] *fossa*, for the origin of similarly named muscles; in the latter is the *nutrient foramen*[13]. The external border presents a *marginal surface*[5], divided by the *oblique line* at the junction of the lower and middle thirds, into two surfaces, the lower for the teres major, the upper for the teres minor muscle; about the junction of the middle and upper thirds of this border is a *groove* for the dorsalis scapulæ vessels. The spine posteriorly presents a *triangular surface*[11], over which plays the trapezius muscle.

The *acromion process*[12] (summit of the shoulder) is a triangular flattened process, curving outward, forward, and upward, to overhang the glenoid fossa; giving attachment along its outer margin to the deltoid; its inner margin, the trapezius; by its *apex* to the coraco-acromial ligament; and having on its inner margin, behind the apex, an articular *facet*[9] (Fig. 34) for the clavicle.

Describe the coracoid process of the scapula.

The *coracoid process*[6] (Fig. 34)[14]; (Fig. 35) (like a crow's beak) rises by a broad base from the upper part of the neck of the scapula, curving over the inner upper part of the glenoid cavity. To the anterior margin, near the tip, is attached the pectoralis minor muscle, and from its apex arise the short head of the biceps and the coraco-brachial muscle. To the inner side of the root is a rough *impression* for the conoid ligament, whence runs obliquely forward and outward on its upper surface a *ridge* for the trapezoid ligament.

Describe the scapular borders.

The superior border[3]—the shortest—presents, at the base of the coracoid, the *suprascapular notch*[4] (Fig. 35), becoming a foramen when the transverse ligament is in situ, *through* which passes the suprascapular nerve, *above* which passes the suprascapular artery; from the border, just internal to the notch, arises the omo-hyoid muscle. The axillary border[5]—the thickest—presents just below the glenoid fossa a *rough surface*[8] for the long head of the triceps muscle, succeeding which is a longitudinal *groove* for part of the subscapular muscle. The vertebral border[9]—the longest—presents an *anterior lip* for the serratus magnus muscle, a *posterior lip* for the supra- and infra-spinatus muscles, and an *intermediate space* for the levator anguli scapulæ above, for the rhomboideus minor from the edge of the triangular base of the spine, and for the fibrous arch of the rhomboideus major muscles below.

Give the points of interest connected with the angles.

The *superior angle*[4] (Fig. 34) affords partial attachment to the serratus magnus, to the levator anguli scapulæ, and supra-spinatus muscles. The *inferior angle*[7] (Fig. 35) affords attachment to the teres major muscle, part of the serratus magnus, and (at times) a few fibres of the latissimus dorsi. The external angle or head[6] presents a shallow pyriform glenoid *fossa*[10] (Fig. 34) for the head of the humerus, deepened, in the fresh state, by the glenoid ligament attached around its circumference, from the upper part of which the long head of the biceps flexor cubiti arises. The *neck*[8] (Fig. 35) is the slightly depressed surface surrounding the head; the *surgical* neck is well posterior to the head, passing through the suprascapular notch[4-8] (Fig. 35).

How is this bone developed?

(7° Rib)

By seven centres; one for body (second fœtal month); two for coracoid (one at first year, one at fifteenth to seventeenth year); two for acromion; one for vertebral border; one for inferior angle (all these last, fifteen to seventeen years); coössification (twenty-two to twenty-five years).

Give the muscular attachments.

Subscapular, supra- and infra-spinatus, deltoid, trapezius, omohyoid, serratus magnus, levator anguli scapulæ, rhomboideus minor and major, triceps, teres minor and major, biceps, coraco-brachial, pectoralis minor, platysma, occasionally latissimus dorsi.

The Arm.

Describe the chief processes and the general characteristics of the humerus.

This, the only *arm* bone, articulates above with the scapula,

Fig. 36.



below with the ulna and radius. The upper extremity, the *head*[2], nearly hemispherical, facing upward, inward, and slightly backward, forms, with the glenoid fossa of the scapula, the shoulder or scapulo-humeral joint. Best marked superiorly is the constriction called the *anatomical neck*[3], indicating the capsular attachment *above*; external to the head is the *greater tuberosity*[4] with three facets for the supra- and infra-spinatus and teres minor muscles; in front, directed inward and forward, is the *lesser tuberosity*[5] for the subscapular muscle; commencing between these two projections, and extending for the upper third of the bone, is the *bicipital groove*, passing obliquely downward and inward, lodging the long head of the biceps muscle; into the *anterior lip*[7] of this groove is inserted the pectoralis major muscle, into the *posterior lip*[8] the latissimus dorsi and teres major; while about the centre of the inner border of the bone is a *rough impression* for the coraco-brachial muscle, just below which is the *nutrient canal*[10], directed downward. The constriction beneath the tuberosities is called the *surgical neck*. The *shaft*, cylindrical above, prismoid and flattened below, presents upon the middle of its outer surface a *roughness*[9] for the deltoid muscle, and below it, winding from behind forward and downward, on the back of the bone, is the *musculo-spiral groove*, for the musculo-spiral nerve and superior profunda artery, internal and external to which arise the inner and outer heads of the triceps muscle. The lower extremity presents an *inner*[14] (very prominent) and an *outer condyle*[13], extending from each of which, upward on the shaft, are the *internal*[16] and *external condyloid*[15] *ridges*[15]; from the external ridge arise the external lateral ligament and *extensor* and *supinator* muscles; from the inner condyle and ridge arise the internal lateral ligament and the *flexor* and *round pronator* muscles. Projecting from the lower front portion of the outer condyle is the *radial head*[11] (capitellum), for the head of the radius; while internal to this, extending from the anterior to the posterior surface of the bone, is the pulley-like *trochlear surface*[12] for the greater sigmoid cavity of the ulna. The *coronoid fossa*[17] in front of the trochlea receives the coronoid process of the ulna during flexion of the forearm, while the *olecranon fossa*, similarly placed behind, accommodates the tip of the olecranon during extension.

How is the humerus developed?

By seven centres: one for shaft (fifth fœtal week); one for head (first to second year); one (sometimes two) for tuberosities (second to third year; by fifth year the centres for head and tuberosities have coössified); one for radial head (second year); one for internal condyle (fifth year); one for trochlea (twelfth year); one for external condyle (thirteenth to fourteenth year). The lower centres are all coössified with the shaft by the sixteenth or seventeenth year except that for the internal condyle, which unites at eighteen years, while the head unites at the twentieth year.

Give the muscular attachments.

The supra- and infra-spinatus, teres minor, subscapular, pectoralis major, latissimus dorsi, teres major, deltoid, coraco-brachial, brachialis anticus, triceps, subanconeus, pronator radii teres, flexor carpi ulnaris, palmaris longus, flexor digitorum sublimis, flexor carpi radialis, supinator longus, extensor carpi radialis longior and brevior, extensor communis digitorum, extensor minimi digiti, extensor carpi ulnaris, anconeus, and supinator brevis.

The Forearm.

Describe the ulna.

This, the inner forearm bone[1], longer than the radius, forms the greater part of the articulation with the humerus, while it is excluded from the wrist-joint by the interarticular fibro-cartilage. Its upper extremity forms the point of the elbow. The *olecranon process*[4] presents upon its anterior part a saddle-shaped articular surface which, with its continuation on the coronoid process, is the *greater sigmoid cavity*[2] for the trochlea of the humerus; continuous with this upon the outer side is the *lesser sigmoid cavity*[3] for the head of the radius; to the posterior surface is attached the triceps tendon, and the *apex* is accommodated by the olecranon fossa of the humerus during extension. Projecting forward below the olecranon is the *coronoid process*^{5}, fitting into the same named fossa of the humerus during flexion, its upper surface forming part of the greater sigmoid cavity; it presents, where it joins the shaft, a tubercle for the oblique ligament, and a *rough impression* for the brachialis anticus muscle above and internally. At the front is a small *eminence* for the flexor sublimis digitorum, whence descends a *ridge* for the pronator radii teres. The *shaft*^{1}, large and prismatic above, smaller and rounded below, has on its anterior surface the *nutrient foramen*, directed upward, and by its *external sharp border*^{7} affords attachment to the interosseus ligament. The lower carpal extremity, or *head*^{8}, articulating, by its outer surface, with the sigmoid cavity of the radius, and, by its lower, with the triangular fibro-cartilage, has projecting internally and behind, the *styloid process*^{9}, to whose apex is attached the internal lateral ligament, to a depression at its base the fibro-cartilage; upon the posterior surface is a *groove* for the tendon of the ulno-carpal extensor.

Fig. 37



How is this bone developed?

By three centres; one for shaft (eighth fœtal week); one for head (fourth year); one for olecranon (tenth year, joining shaft at sixteenth year); head coössified with shaft by twentieth year.

Give the muscular attachments.

Triceps, anconeus, flexor carpi ulnaris, brachialis anticus, pronator radii teres, flexor sublimis and profundus digitorum, flexor longus pollicis (occasionally), pronator quadratus, flexor and extensor carpi ulnaris, anconeus, supinator brevis, extensor ossis metacarpi and secundi internodii pollicis, and extensor indicis.

Describe the radius.

Shorter than the ulna, situated upon the outer side of the forearm, with a small upper extremity, forming only a small part of the elbow-joint, its lower end is large, forming the chief part of the wrist-joint. It is slightly curved, and of a prismatic form. The *head*[11] (upper extremity) is cylindrical, with slightly cupped upper surface, articulating with the radial head of the humerus, by its sides, with the lesser sigmoid cavity of the ulna and the orbicular ligament by which it is embraced; the constriction beneath the head is the *neck*[12]; below, and to the inner side, is the *tuberosity*[13], rough toward its posterior margin for the biceps tendon, smooth in front for a bursa; the *shaft*[10] is prismoid, curved outward, smaller above than below, having a sharp internal *interosseous border*[7] for the interosseous ligament, with the *nutrient foramen* directed upward at the junction of the middle and upper thirds of the anterior surface.

The lower *carpal extremity*[15] has on its lower face an *articular surface* divided by a slight ridge into two facets for the semilunar and scaphoid bones; upon its inner side the shallow *sigmoid cavity* for the ulnar head; externally the *styloid process*[16] giving attachment by its apex to the external lateral ligament, by its base to the supinator longus muscle; and on its posterior and external convex surfaces are five *grooves* for the extensor tendons.

How is this bone developed?

By three centres; one for shaft (eighth fœtal week); one for carpal extremity (second year); and one for head (fifth year, joins shaft at puberty); bone coössified by about twentieth year.

Give the muscular attachments.

The biceps, the supinator brevis, flexor digitorum sublimus and longus pollicis, pronator quadratus, the extensor ossis metacarpi and primi internodii pollicis, pronator radii teres, and supinator longus.

The Hand.

Into what segments is the hand divided?

Into the carpus (eight bones); metacarpus (five bones); and phalanges (fourteen bones); total, twenty-seven bones.

- Silly sailors can't pin their ties; man urethra.

Describe the arrangement of the carpal bones.

They are placed in two rows of four each; thus, enumerating from the radial to the ulnar side, with palm upward; First, or *proximal row*: scaphoid^{s}, semilunar^{l}, cuneiform^{c}, pisiform^{p}.

Second, or *distal row*: trapezium^{t}, trapezoid^{1},^{t}, os magnum^{7}, unciform^{u}.

FIG. 38.



With how many bones does each articulate?

Scaphoid, five; semilunar, five; cuneiform, three; pisiform, one; trapezium, four; trapezoid, four; os magnum, seven; unciform, five; total, thirty-four.

Describe the chief peculiarities of each bone.

The *scaphoid*^{s} (boat-shaped) has on the thumb side a tuberosity. a transverse groove on the dorsum parallel to the convex articular surface for radius; facets for os magnum and semilunar bones on its inner lower face; on its lower for the trapezium and trapezoid.

The *semilunar*^{l} (half-moon) presents a convex facet above for radius; on its outer face a semilunar facet for scaphoid; on its lower, a concave one for os magnum; a quadrilateral one on the inner face for cuneiform.

The *cuneiform*^{c} (wedge-shaped) has an oval facet in front for pisiform; one external for semilunar; an inferior concave facet for unciform; a superior convex facet for interarticular fibro-cartilage.

The *pisiform*[p] is rounded, with one ovoidal facet for cuneiform.

The *trapezium*[t] is obliquely grooved on its palmar surface for the tendon of the flexor carpi radialis; superiorly is a concave facet for scaphoid; below, a saddle shaped one for thumb-metacarpal; internally, one large concave facet for trapezoid, and a smaller for second metacarpal.

The *trapezoid*[1t] is wedge-shaped, apex palmar, has four articular surfaces touching, separated by sharp edges, the external for trapezium; inferior (like a ridge-roof), for second metacarpal; internal, for os magnum; superior, for scaphoid.

The *os magnum*[7] has superiorly a convex head for scaphoid and semilunar, a neck and body; below, facets for three metacarpal bones; externally, one for trapezoid; internally, one for unciform.

The *unciform*[u] is triangular, has a sigmoid internal articular facet for cuneiform; two facets below for fourth and fifth metacarpals; one, external, for os magnum; from the palmar surface projects the curved unciform process.

When do these bones ossify?

Os magnum and unciform, during first year; cuneiform, third year; trapezium and semilunar, fifth year; scaphoid, sixth year; trapezoid, during eighth year; pisiform, about twelfth year.

(The muscular attachments to the carpal and tarsal bones will be given under the Muscular System.)

Describe the metacarpal bones.

They are five long bones, with shaft, head, and base. The *first metacarpal* (sometimes considered a phalanx) is shorter, has only one facet on base, and has one ossific centre for shaft (eighth fœtal week), and one for *base* (third year). The *second metacarpal* has four facets on base for trapezium, trapezoid, os magnum, and third metacarpal; it, in common with the rest, has one ossific centre for shaft (eighth fœtal week), and one for *head* (third year; they coössify by twentieth year). The *third metacarpal* articulates only with the os magnum, on the ulnar side has two small facets for fourth metacarpal, a single facet on the radial side for second metacarpal, and the outer angle of its base is much prolonged. The *fourth metacarpal* has two facets for unciform and os magnum, a single facet on ulnar side for fifth metacarpal, and two small ones on radial side for third metacarpal. The *fifth metacarpal* articulates with unciform by a concavo-convex facet, has only a lateral facet on the radial side for fourth metacarpal, and on the ulnar side a prominent tubercle for the extensor carpi ulnaris tendon.

Describe the phalanges.

Fourteen in number—three for each finger, two for thumb—they are long bones, having a shaft, base, and condyles, except those of the distal row. The bases of first row, cup-shaped, articulate with heads of the metacarpals; those of second and third have a double concavity, separated by a median ridge, and articulate with the condyles of the row above; the distal extremities of the ungual phalanges have rough, horse-shoe shaped tubercles on their palmar surfaces for attachment of the pulp of the finger; they ossify by one centre for the shaft (eighth fœtal week), and one for base (third to fifth year); uniting, from eighteen to twenty years.

LOWER EXTREMITY.

The Thigh.

Describe the femur.

It is the longest bone of the skeleton, and inclines toward its fellow to bring the knee-joint near the centre of gravity during walking, this obliquity being greater in the female, from the breadth of the pelvis, and in a short than a tall person of either sex.

The *head*[2], forming rather more than a hemisphere, directed upward, inward, and forward, has behind and below its centre an ovoid *depression* for the ligamentum teres. The *neck*[3], joining the head with the shaft, is flattened and pyramidal; in the adult male it forms an obtuse angle with the shaft; in the female it approaches a right angle; in very old and debilitated subjects its direction becomes horizontal. The *great trochanter*[4] is a large, irregular, quadrilateral eminence, directed upward, outward, and backward, marked on its external surface by a *diagonal line* for the gluteus medius; below and behind this is a smooth surface for a bursa beneath the gluteus maximus; in front, from above downward, are inserted the internal obturator and gemelli muscles, the pyriformis, and the gluteus minimus; upon its inner surface is the *digital fossa* for the tendon of the obturator internus muscle.

FIG. 39.



The *lesser trochanter*[6], small and conical, projects from the lower back part of the base of the neck, receiving the insertion of the psoas muscle above, and the iliacus below.

The *anterior*[5] *and posterior intertrochanteric lines* connect these processes, the latter being the more prominent, while to the former is attached the anterior portion of the capsular ligament. Extending from the middle of the posterior intertrochanteric line, for about two inches down the shaft, is the *linea quadrati* for the quadratus femoris muscle.

The *shaft*[1], broad and cylindroid at either extremity, narrow and triangular in the centre, slightly curved forward, has its *nutrient foramen* at the junction of the middle and lower thirds of its posterior surface, directed upward; and from its anterior surface arise the crureus and subcrureus muscles.

The *linea aspera*, a prominent longitudinal ridge occupying the middle third of the posterior surface, has an external and an internal lip, and an intermediate space; above, it divides into three lines, one directed upward to base of greater, one to base of lesser trochanter, a third, the most internal continuous with the anterior *intertrochanteric line*, forming, with it, the *spiral line*[5], while below the linea aspera bifurcates to inclose the smooth *popliteal space*, the inner division *grooved* for the femoral vessels. In general terms, this line and its subdivisions have attached the following muscles: the vastus internus and externus, the pectineus, the three adductors, the biceps and gluteus maximus.

Of the two *condyles*, the *internal*[8] is the longer by about half an inch, to bring both condyles on the same horizontal plane in the normally oblique position of the femur; above each condyle, behind, is a *depression* for the gastrocnemius (above the *external* the plantar muscle also originates); separating them is the *intercondyloid notch*, to whose sides are attached the crucial ligaments; in front, the condyles form a continuous cartilaginous-covered articular surface; the outer, as well as the inner condyle, presents a *tuberosity*[9]-[11] upon its free surface for the lateral ligaments, beneath which, on the outer condyle, is a *groove*[10] for the tendon of origin of the popliteus muscle; above the *internal tuberosity*[11] is a small *tubercle* for the tendon of the adductor magnus.

Give the development of the femur.

By five centres; one for shaft (fifth fœtal week); one for condyles (ninth fœtal month); one for head (end of first year); one for greater trochanter (fourth year); one for lesser trochanter (thirteenth to fourteenth year); all coössified by twentieth year in the *reverse order of their appearance.*

Give the muscular attachments.

The three glutei, pyriformis, two obturators, gemelli, quadratus, psoas, iliacus, two vasti, biceps, pectineus, three adductors, crureus, subcrureus, gastrocnemius, plantaris, and popliteus.

The Patella.

Describe the patella.

It is flat, triangular, placed at front of the knee-joint, and, being developed in the quadriceps tendon, is considered a sesamoid bone by some; the convex *anterior surface* is roughened; the *posterior surface* is divided by a vertical *ridge* into two smooth *facets* for either femoral condyle, the outer being the broader and deeper; the apex gives attachment to the ligamentum patella; and the superior and lateral borders to the rectus femoris, crureus, and vasti muscles. It is developed by one centre (about three years); the muscular attachments have been already given.

The Leg.

Describe the tibia.

FIG. 40.



The shin-bone, situated at the inner front part of the leg, is only second in length to the femur; the *head*[2,3], or upper extremity, is large and expanded on each side into two lateral *tuberosities*[2,3], bearing upon their upper surfaces smooth concave *ovoidal facets* for the femoral condyles, between which is the vertical, bifid *spine*[4] for the extremities of the semilunar fibro-cartilages, the depressions in front and behind its base giving origin to the crucial ligaments. Below, medianally, in front of the head, is the *tubercle*[5] for the ligamentum patellæ; separating the tuberosities behind is the *popliteal notch*, giving attachment to the posterior crucial ligament; on the posterior surface of the inner tuberosity is a transverse *groove* for the insertion of the semimembranosus tendon; upon the back of the outer tuberosity, facing downward, is a *facet* for the head of the fibula; running obliquely from this facet, downward and inward, on the posterior surface, is the *oblique line* for the popliteal fascia, parts of the soleus, flexor longus digitorum, and tibialis posticus muscles; just below the line, directed downward, is the *nutrient* canal, the largest in the skeleton.

The prismoid shaft[1] has three borders, of which the anterior, called *crest* or *shin*[6], and the external or *interosseous* ridge, for the interosseous membrane, only are of importance. The *lower extremity*[7], smaller than the upper, has an inferior concave surface for the astragalus, an external rough triangular surface for articulation with the fibula; it is grooved posteriorly for the flexor longus pollicis tendon; has projecting downward, internally, the *internal malleolus*[8], which articulates by its outer surface with the side of the astragalus, is grooved behind for the tibialis posticus and flexor longus digitorum tendons, and has attached to its tip the internal lateral ligament.

How is this bone developed?

By three centres: One for shaft (seventh fœtal week); one for head (at birth); one for lower end (second year); bone coössified by twentieth year (some authors say twenty-fifth year).

Give the muscular attachments.

Semimembranosus, tibialis anticus and posticus, biceps, extensor and flexor longus digitorum, sartorius, gracilis, semitendinosus, popliteus, soleus, and quadriceps femoris.

Describe the fibula (peroneal bone).

It is long, slender, and is placed externally nearly parallel to the tibia.

The head[10] or upper end articulates by a flattened facet with the external tibial tuberosity[3], and is prolonged upward behind into the *styloid process* for the biceps tendon and the external lateral ligament; below and behind is attached the long external lateral ligament.

The shaft[9], triangular and twisted, has three ridges, the internal or *interosseous ridge* being for same-named ligament; about the middle of the anterior internal surface is the *nutrient foramen* directed downward. The *external malleolus*[11] forming the lower extremity, longer than the internal, articulates by its inner surface with the outer side of the astragalus; is grooved behind for the peroneus longus and brevis tendons; to its summit is attached the middle, and to rough *depressions* in front and behind the anterior and posterior fasciculi of the external lateral ankle-ligament.

Describe its development.

By three centres: One for shaft (eighth fœtal week); one for malleolus (second year); one for head (fourth year); bone coössified by twenty-fifth year, but, *contrary to rule*, the lower epiphysis unites *first*.

Give the muscular attachments.

Biceps, soleus, three peroneals, the extensor and flexor of the great toe, and extensor longus digitorum, and tibialis posticus.

The Foot.

Into what segments are the bones of the foot divided?

Into the *tarsus* (7); *metatarsus* (5); and *phalanges* (14); total, 26 bones.

Name the tarsal bones.

Calcaneum[3] (os calcis), astragalus[1], cuboid[8], scaphoid[4], internal[5], middle[6], and external[7] cuneiform.

Give the chief peculiarities of each tarsal bone.

The *astragalus*[1] has a large rounded *head*[2], a *neck*, a *body*[1], superiorly a *trochlear surface* broader in front than behind, for the tibia, and continuous with it on either side *facets* for the internal and external malleolus; the under surface presents two articular *facets* —that behind for the calcis, that in front partly for calcis but chiefly for the calcaneo-scaphoid ligament—separated by a groove for the calcaneo-astragaloid ligament, running obliquely forward and outward; it articulates with the tibia, fibula, os calcis, and scaphoid, and is *developed* by one centre (seventh fœtal month).

The *os calcis*[3], the largest tarsal bone, forms by its *tuberosity*[3] behind the heel, has a *groove* on its upper surface to correspond to that of the astragalus; behind and in front of which are two articular *facets* for the same bone: on the inner side projects the *sustentaculum tali* supporting the internal articulating surface; beneath this process the inner surface of the bone is deeply concave for the flexor tendons, plantar vessels, and nerves; its anterior concavo-convex surface articulates with the cuboid; on the under surface are an *inner* and *outer tuberosity*; it articulates with the astragalus and cuboid, and is *developed* from two centres, one for main mass (sixth fœtal month); one for tuberosity (tenth year); union after puberty.

The *cuboid* has one articular surface each for os calcis, external cuneiform, the fourth and fifth metatarsals, and sometimes the scaphoid; upon the under surface is a deep *groove* for the peroneus longus tendon, and behind this a *ridge* terminating externally in a *tuberosity*; it is *developed* from one centre (ninth fœtal month).

The scaphoid, situated internally, is concave behind for the head of the astragalus; is convex in front with three *facets* for the three cuneiform bones; externally is a *facet* for the cuboid; and internally, below, is the *tuberosity* for part of the posterior tibial tendon; it is *developed* from one centre (fourth year).

FIG. 41.



The internal cuneiform, the largest, is placed at the inner side of the foot, has its base downward, upon which is the *tuberosity* for part of the tendon of the posterior tibial muscle; in front is a kidney-shaped *facet* for first metatarsal; externally are two *facets* for the second metatarsal in front, the middle cuneiform behind; posteriorly a *facet* for the scaphoid; it is *developed* by one centre (third year).

The middle cuneiform, the smallest, has its base upward, a triangular *facet* in front for second metatarsal, another behind for the scaphoid; along the posterior and superior borders of the inner face a *facet* for the internal cuneiform; and externally a smooth facet for the external cuneiform; it is *developed* by one centre (fourth year).

The external cuneiform is intermediate in size with base upward, has an anterior triangular *facet* for third metatarsal; another posterior for the scaphoid; two upon internal surface for second metatarsal and middle cuneiform; and two upon outer surface for fourth metatarsal and for cuboid: it is *developed* by one centre (first year).

Describe the metatarsal bones.

These five long bones have prismoid shafts[9], anteriorly a *head* for articulation with the phalanges, posteriorly a *base* articulating with the tarsus and one another.

The first metatarsal is shorter, much stouter, and articulates only with the internal cuneiform: *developed* by one centre for *shaft* (seventh fœtal week); one for *base* (fifth year), united by the twentieth year.

The second metatarsal, the longest, articulates posteriorly with the middle cuneiform, and laterally with the other cuneiforms, presenting, therefore, three *facets* on base; *developed* by one centre (seventh fœtal week) for *shaft*, one for *head* (third year), united at twenty years.

The third metatarsal has a *facet* on base for external cuneiform, two on its inner side and one on its outer for the contiguous metatarsal; *developed* like second.

The fourth metatarsal articulates behind with cuboid, has a *facet* on inner side divided into anterior portion for third metatarsal, a posterior for external cuneiform, and externally one *facet* for fifth metatarsal: *developed* like second.

The fifth metatarsal has a triangular oblique surface for the cuboid, continuous internally with one for fourth metatarsal; externally a tubercular *eminence*; *developed* like second.

Describe the phalanges.

They resemble closely those of the hand, except that they are strongly compressed from side to side, instead of from before backward: ossification also, similar but later. Thus, the shaft centres appear from two to four months, except distal at seventh fœtal week.

THE ARTICULATIONS.

How are the articulations classed ?

In three divisions; 1. *Synarthroses*, immovable, as most of cranial articulations; 2. *Amphiarthroses*, including *synchondroses*, or *symphyses*, yielding (limited motion), as those between the vertebral bodies, the pubic and sacro-iliac symphyses; (4) 3. *Diarthroses*, freely movable.

Into what classes are the synarthroses divided ?

1. *Sutura vera*, consisting of ~~interlocking~~ serrations, including three sub-classes of *S. dentata*, when the bony projections are tooth-like, as the inter-parietal suture; *S. serrata*, when like a fine saw, as the inter-frontal; *S. limbosa*, when the edges are bevelled in addition to dentation, as the fronto-parietal. The *Sutura notha* (false

(1) together – joint
(2) around (both) –
3.) – Cartilage.
thro' –

sutures) include *S. squamosa*, formed by two overlapping bevelled edges, as the squamo-parietal suture; and the *S. harmonia*, mere apposition of roughened surfaces, as the two superior maxillary bones.

1. 2. *Schindelysis*, where a thin edge is received into a cleft or fissure, as the vomer between the superior maxillary bones.

2. 3. *Gomphosis*, the insertion of a conical process into a socket, as ~~teeth~~ in their alveoli (not really a bony articulation, as teeth are *not* bones).

How are diarthroses classed?

As ~~*arthrodia, gliding joi*~~nts—the articular processes of the ~~verte~~bræ; *enarthroses*, ball-and-socket joint—hip- or shoulder-joints: *ginglymus*, hinge-joint—elbow-joint; *diarthrosis rotatoria*, or *lateral ginglymus*, a pivot turning within a ring, or a ring on a pivot, as the superior radio-ulnar and atlo-odontoid. (Trochoides (like pulley, [illegible] Saddle (Reciprocal recep. - thumb) Condyloid ([illegible]

What varieties of motion do joints enjoy?

Flexion, extension, adduction, abduction, circumduction (a combination *in succession* of the four preceding), rotation, and gliding.

What structures are essential to the formation of each of the three classes of articulations?

(1) For synarthroses, two or more bones, an interposed layer of fibrous tissue (sutural ligament) or, perhaps, cartilage (base of the skull); *symphyses*, an interposed bond of fibro-cartilage, with strong bands of white fibrous tissue, *i. e.*, *ligaments* (ligaments are sometimes composed of yellow elastic tissue, as the ligamenta subflava or ligamentum nuchæ); (2) *diarthroses*, two or more cartil~~aginous-coated~~ surfaces (reducing friction), sometimes inter-articular fibro-cartilages to deepen joint surfaces, as those of the knee- and temporo-maxillary joints, a ~~complete fibrous capsule~~, and often additional ligamentous bands, some inter-articular, *i. e.*, within the joint cavity, and a s~~ynovial (serous)~~ membrane lining the interior of the ~~capsu~~le, but not extending upon the cartilages; a similar membrane also forms sacs (~~bursæ) o~~utside the joints, with which they often communicate, serving to reduce friction of the tendons, ligaments, etc.; the layer of ~~bone~~ beneath the articular

Synde[illegible] F Ch.

[illegible]

cartilage ~~is denser,~~ contains neither Haversian canals nor canaliculi, and has larger lacunæ.

What is the rule as regards the nerve supply of joints?

The ~~interior~~ of the joint, the ~~muscles~~ moving it, and the ~~skin~~ over their insertions, are supplied by the ~~same trunk~~ or trunks of nerves (this explains the ~~reflex contractions of diseased~~ joints).

Describe the vertebral articulations.

Formed by the contiguous surfaces of the vertebral bodies and articular processes, their ligaments are as follows:

1. An *intervertebral connecting fibro cartilage*, between the bodies of all true vertebræ, except the atlas and axis.

1. An *anterior common ligament*[3] passing medianally over the fronts of the vertebral bodies, most firmly attached to their margins.

1. A *posterior common ligament*, similarly disposed behind.

Short intervertebral ligaments, fibres running at most over three vertebræ, firmly uniting the bodies where the anterior and posterior common ligaments are deficient.

Ligamenta subflava, of yellow elastic tissue, connecting the laminæ.

Supra- and inter-spinous, the former connecting the tips, the latter the remainder of the spinous processes.

2. *Capsular*[6], enclosing the articular processes, and lined with synovial membrane.

2. *Inter-transverse*, connecting transverse processes; *nerves*, spinal in each region; *arteries*, vertebral and ascending cervical arteries in neck, intercostals in dorsal region, lumbars in loin.

Describe the occipito-atloid articulation.

It is a ~~ginglymo-arthrodial~~ joint formed by the condyles of the occipital bone and the superior articulating processes of the atlas. Its ligaments are,

1. *An anterior occipito atlantal*[2] (Fig. 42), extending from the anterior margin of the foramen magnum to the anterior arch of the atlas, about one inch broad, blending on either side with the capsular ligaments.

1. *A posterior occipito-atlantal*, much broader, from the posterior margin of the foramen magnum between the condyles, to the postero-superior border of the posterior arch of the atlas, and is incomplete on each side for the ingress of the vertebral artery, and egress of the suboccipital nerve.

3- Two *capsular ligaments*[7], lined with synovial membrane, surrounding the articular surfaces.

2. *Two lateral* (or *anterior oblique*), passing upward and inward from the ~~transverse~~ process beyond the vertebral foramen to the inner edge of the jugular foramen; *nerve*, suboccipital; *arteries*, from vertebral.

FIG. 42.



Describe the atlo-axoidean points.

The lateral joints are ~~arthrodia~~, that between the atlas and odontoid process a double diarthrosis rotatoria. The ligaments of the lateral joints and arches are,

1. The *anterior atlanto-axoidean*[4], membranous, passing between lower front border of atlas to front of the axis.

1. The *posterior atlanto-axoidean*, stretching between the postero-inferior edge of the ring of the atlas to the superior edge of the arch of the axis behind; it is pierced on each side by the second spinal nerve.

The *anterior*[1-3] and *posterior common* ligaments are continued over the median portions of the above to the occiput.

3. Two *capsulars*[56], synovial-lined, surrounding the articulating processes.

The ligaments of the central atlanto-axoidean joints are:

5. The *transverse*[2] (Fig. 43), extending between the tubercles on the inner surface of each lateral mass of the atlas; it ~~holds the odon-toid~~ process in place, and between the two is a synovial membrane surrounded by a *capsule*, the *odonto-transverse joint;* between the odontoid process and the posterior surface of the anterior arch of the atlas, is the atlo-odontoid articulation, surrounded by a *capsule* lined with synovial membrane; passing upward and downward, are two strong vertical bands, attached above to the cranial surface

of the basilar margin of the foramen magnum, below, to the upper half of the body of the axis, forming the *occipito-axiodean ligament*[3-4], or vertical portion of the *crucial ligament; nerves*, all these joints are supplied by the second cervical or loop between it and the suboccipital; *arteries*, are branches of vertebral.

What other ligaments connect the axis and occiput?

The *posterior common ligament*[1] extends upward, to be attached far up the ~~basilar process~~; beneath this is,

The *occipito-cervical* or *cervico-basilar ligament*, attached above to the basilar groove of the occiput, below, to the third cervical body and that of the axis; (next comes the vertical part of the crucial already described); deepest of all are the,

FIG. 43.



FIG. 44.



Occipito-odontoid, or *check*[5], extending from the sides of the apex of [illegible] the odontoid to inner edge of the ~~anter~~ior portion of the occipital condyles, while stretching between the odontoid tip and the under surface of the basilar process close to the foramen magnum, is the

Central occipito-odontoid, or *ligamentum suspensorium*.

Describe the temporo-maxillary articulation.

The upper compartment is ~~arthro~~dial, formed by the glenoid fossa and eminentia articularis of the temporal bone and the upper surface of the fibro-cartilage; the ~~lower is~~ ginglymoid, between the under surface of the fibro-cartilage and the condyle of the lower jaw. The ligaments are *capsular*, passing between maxillary and temporal bones near their margins, consisting of ligamentous fibres, which are thicker at certain parts, and are described as

2. The *external lateral ligament*[5], attached above to nearly the whole length of the lower edge of the zygoma and its tubercle, below to the outer side of the neck of the condyle of the jaw.

1. The *short internal lateral ligament*, extending between the ~~spine~~ of the sphenoid and inner edge of the glenoid fossa, ~~and a~~ ridge on the inner side of the neck of the condyle.

2. The *long internal lateral ligament*, stretching between the spine of the ~~sphenoid~~ and the forepart of the tip of the inferior dental foramen.

The *inter-articular fibro-cartilage*, concavo-convex on the upper surface, concave obliquely transversely below, dividing the joint into two separate synovial cavities.

The *stylo-maxillary ligament*[6], extending from the styloid process to the angle of the jaw, is really a process of the deep fascia; *nerves*, the masseteric and auriculo-temporal branches of the inferior maxillary nerve; *arteries*, temporal, middle meningeal, ascending pharyngeal, posterior auricular, tympanic branch of the internal maxillary and ascending palatine.

Describe the costo-vertebral articulations.

These are each formed (1) between the head of a rib and the ~~bodies of~~ two adjoining vertebræ (except first, tenth, eleventh, and twelfth, which articulate with but one vertebra each), being *~~ginglymoid~~*; and (2) between the tubercle of each rib (except eleventh and twelfth) and the tran~~sverse process~~ of a vertebra, and is ~~arthrodial~~.

Describe the costo-central articulations.

Ginglymoid, the ligaments are (1):

A *capsular ligament*[4], attached all around each articular surface, the synovial lining being subdivided into two cavities by the

Inter-articular, or *costo-vertebral ligament*[4], passing between the ridge on the head of the rib to the intervertebral fibro-cartilage.

FIG. 45.



The *stellate ligament*[2], arising from the anterior surface of the neck of the rib, whence the fibres radiate to the vertebra next above and below to the one the rib belongs to, and to the intervertebral disk; *nerves*, anterior branches of spinal nerves; *arteries*, the intercostals.

Describe the costo-transverse articulations.

Arthrodial, ten in number, their ligaments are:

Capsular, attached beyond the margins of the articular facets.

Superior costo-transverse[3] (except first rib), passing between upper border of neck of rib to lower margin of transverse process of the vertebra above.

Middle costo-transverse, connecting the back of neck of rib with the contiguous portion of transverse process of vertebra to which rib belongs.

Posterior costo-transverse, stretching from the tip of transverse process to rough projection beyond facet on the tubercle of rib; *nerves*, posterior branches of dorsal nerves; *arteries*, intercostals and posterior spinal.

Describe the (1) costo-sternal articulations, (2) the connection between the ribs and costal cartilages, and (3) the interchondral joints.

(1) The first is *synarthrodial*, the other six joints are *ginglymoid*.

The *costal* (2) *cartilages* are firmly attached to the cup-like depression at the end of each rib, and by the periosteum continued over to form the perichondrium; in like manner, the cartilage of the first rib unites with the sternum. The cartilages of the remaining six ribs are bound to the sternum by a,

Complete capsular ligament, the anterior and posterior thicker segments sometimes described as,

Anterior and posterior costo-sternal ligaments. Occasionally there is an *interarticular ligament*, notably in the second costal joint, dividing the synovial cavity into two.

The (3) interchondral articulations, *arthrodial*, are usually, but not always, found between the edges of the fifth to tenth ribs, inclusive, having each an *interchondral ligament* and a *capsular*, lined with synovial membrane; they are commonly found only from the sixth to the ninth cartilages; in addition, the *costo-xiphoid ligament* binds the xiphoid cartilage and sixth and seventh cartilages together; *nerves*, intercostals; *arteries*, internal mammary or its branches.

Describe the ligaments of the sternum.

All three segments are bound together by the so-called,

Anterior and posterior ligaments, with a layer of cartilage between the manubrium and gladiolus (an amphiarthrodial joint, that between ensiform cartilage and gladiolus being synarthrodial).

Describe the sacro-vertebral articulation.

✓ Similar to other vertebral articulations, but has, in addition,

1. ANT The *sacro-lumbar ligament*, extending from the transverse processes, pedicles, and body of the fifth lumbar vertebra to the non-articular portions of the base of the sacrum and periosteum of the contiguous portions of ilium.

2. LAT. The *ilio-lumbar ligament*[1] (Fig. 46), extending from the front surface and back of the tip of the transverse processes of the fifth, and lower edge and front surface, the transverse processes and pedicles of the fourth lumbar vertebra, to the back part of the iliac crest; *nerves*, fourth and fifth lumbar, sympathetic; *arteries*, ilio-lumbar, last lumbar, lateral sacral.

Describe the sacro-iliac articulations.

They are *amphiarthrodial*, formed between the auricular surfaces of the ilium and sacrum, and have the following ligaments upon each side, in addition to the symphyseal cartilage:

1. The *anterior sacro-iliac*[3] (Fig. 47), from the first three pieces of the sacrum to the ilium above the great sacro-iliac foramen.

1. The *posterior sacro-iliac*[2] (Fig. 46), very strong, extending between back of the sacrum and posterior two inches of the iliac crest, including the posterior superior spine.

The *oblique*, stretching between the third transverse tubercle of the sacrum to the posterior superior iliac spine.

2. *Superior and inferior*, bands of fibrous tissue closing in the joint above and below.

The *interarticular* (interosseous), strongest of all, consists of numerous ligamentous bands passing between the contiguous rough surfaces of the sacrum and ilium; *nerves*, from posterior sacral and lumbo-sacral cords; *arteries*, gluteal, ilio-lumbar, and sacral spinal.

Describe the ligaments connecting the sacrum and ischiatic bones on each side.

They are the,

Greater sacro-sciatic[3], thin, triangular, passing from posterior inferior iliac spine, fourth and fifth transverse sacral tubercles and lower lateral margins of sacrum and coccyx, to the inner margin of the ischiatic tuberosity.

FIG. 46.



The *lesser sacro-sciatic*[4], posterior to former, of same shape, stretching between lateral margins of sacrum and coccyx, and the spine of the ischium.

These ligaments convert the two sacro-sciatic notches into the *greater* (by lesser ligament) and *lesser* (by greater ligament) sacro-sciatic foramina, described under iliac and ischiatic bones.

Describe the sacro-coccygeal articulation.

Amphiarthrodial, its ligaments are,

An *interarticular fibro-cartilage*, like that of a vertebra.

An *anterior sacro-coccygeal*, a continuation of the anterior common vertebral.

1· A *posterior sacro-coccygeal,* a continuation of the posterior common vertebral.

2. *Inter-cornual* and *inter-transverse* bands, connecting parts of same name; *nerves,* anterior and posterior branches of the fifth sacral and coccygeal, posterior division of fourth, and probably second and third sacral; *arteries,* lateral and median sacral.

Describe the pubic articulation.

Amphiarthrodial, between the two pubic bones; its ligaments are

1. An *interosseus fibro-cartilage.*

2· A *supra-pubic,* extending along crest of pubes on each side, blending with fibro-cartilage.

1. A *posterior,* little more than periosteum.

1· An *anterior,* thick, strong, decussating, attached to contiguous portions of body and rami.

2 An *inferior,* or *sub-pubic,* arched, filling up angle between pubic rami; *nerves* and *arteries* of no special moment.

The *obturator ligament*[5] is a fibrous membrane filling the same named foramen, except at upper outer part.

Describe the sterno-clavicular articulation.

An *arthrodial* joint formed between the sternal end of the clavicle, and the sternum and cartilage of the first rib, having an inter-articular fibro-cartilage usually dividing the joint into two distinct synovial sacs; its ligaments are,

A *capsular* (usually described as anterior, posterior, etc.) passing between the articular margins, and firmly connected with the fibro-cartilage.

The *inter-clavicular,* passing from the posterior superior angle of the inner extremity of each clavicle, and attached to the sternum between.

The *rhomboid,* or *costo-clavicular,* stretching from the upper border of the first costal cartilage to the rhomboid impression of the clavicle; *nerves,* from descendens noni; *arteries,* contiguous muscular branches.

Describe the acromio-clavicular articulation.

An arthrodial joint formed by the outer extremity of the clavicle and the acromion process of the scapula; it frequently has an inter-articular fibro-cartilage and two synovial sacs, otherwise there is only one sac; its ligaments are,

The *capsular*[1], attached to anterior and posterior borders, upper and lower surfaces of the acromion and clavicle.

The *coraco-clavicular*[2], composed of the *conoid*[2], conical, attached by apex to base of coracoid process, by base to conoid tubercle of clavicle and a line internal to it.

Trapezoid[2], broad and thin, quadrilateral, stretching between ridge on upper surface of coracoid to oblique line on under surface of clavicle; *nerves*, supra-scapular, anterior circumflex· *arteries*, supra-scapular, anterior circumflex, acromial thoracic.

What ligaments bind together the scapular processes?

The *coraco-acromial*[3], a triangular flat band, attached by apex to summit of acromion, by base all along outer border of coracoid process.

What other ligament of the scapula has not yet been described?

The *transverse*[4], bridging across the supra-scapular notch.

Describe the shoulder-joint. = Capsule.
Overlying tendons.
x accessory folds. (2)

An enarthrodial joint formed by head of the humerus and glenoid fossa of the scapula. Its synovial membrane is reflected upon the tendons of the biceps, subscapular, and infra-spinatus muscles, and the joint has numerous bursæ in its vicinity, with some of which it communicates, notably the subacromial and subscapular; the ligaments are,

Capsular[5], from the margin of the glenoid fossa above, below, and behind, in front extending up on the venter half an inch or more, to be attached by its upper half to the ~~anatomical neck~~ of the humerus, its lower some distance from the articular margin.

Coraco-humeral[6], a superadded band stretching from coracoid process of scapula to greater humeral tuberosity.

The *glenoid*, a dense fibro-cartilage, triangular in cross-section, attached to circumference of fossa, deepening the socket, and continuous above with the long head of biceps muscle[7], which really serves as one of the *chief ligaments*; *nerves*, supra-scapular, circumflex, subscapulars; *arteries*, supra-scapular, subscapular, dorsalis scapulæ, anterior and posterior circumflex.

Describe the elbow-joint.

Ginglymoid, formed by lower end of the humerus, greater sigmoid cavity of ulna and head of radius, its ligaments are,

FIG. 47.



FIG. 48.



3. A capsule[1], large, capacious, and usually described as *anterior, posterior, internal,* and *external ligaments.*

1. The *anterior segment*[1] extends from humerus, above articular surface and coronoid fossa, to front of coronoid process of ulna and neck of radius.

1. The *posterior segment* extends from the back of the humerus, from condyle to condyle, and above the olecranon fossa, to be attached all around the olecranon process, close to its articular margin, and to back of neck of the radius and coronary ligament of the same.

2· The *external lateral* radiates from lower part of condyle to outer side of neck of radius and coronary ligament. = orbicular.

2. The *internal lateral*[2], triangular, rises from antero-inferior aspect of inner condyle, and is attached to inner side of shaft of ulna and olecranon process; *nerves*, musculo-cutaneous, ulnar, median, musculo-spiral; *arteries*, the two profunda arteries, anastomotica magna, anterior and posterior ulnar recurrents, posterior interosseous recurrent, and radial recurrent.

Describe the superior radio-ulnar articulation.

It is a diarthrosis rotatoria, or trochoides, formed by head of radius and lesser sigmoid cavity of ulna, its synovial membrane being continuous with that of elbow-joint; its only ligament is the

Orbicular[3], surrounding the head of the radius, forming only three-fourths of a circle by most of its fibres, but some, continued below the sigmoid cavity, form a complete circle: *nerves* and *arteries*, those of elbow.

Describe the inferior radio-ulnar articulation.

A ~~lateral~~ ginglymus, formed by ulnar head and sigmoid cavity of radius; its synovial membrane is so loose as to be called the *membrana sacciformis*; the ligaments are,

f. The *triangular fibro-cartilage*, attached by apex to fossa at base of styloid process of ulna, by its base to margin of radius below sigmoid cavity.

1. The *anterior radio-ulnar* stretches between anterior edge of sigmoid cavity of radius to rough surface above articular surface of ulna.

1. The *posterior radio-ulnar* is similarly attached behind; *nerves*, median and posterior interosseous; *arteries*, anterior and posterior interosseous and carpal arch.

What other ligaments bind the radius and ulna together?

The *interosseous membrane*[5], passing obliquely downward from the interosseous ridge of the ulna to that of the radius.

The *oblique*[4], attached to the ~~tubercle~~ at base of coronoid process of ulna above, and below to the lower posterior edge of the ~~tubercle~~ of radius; *nerves* and *arteries* are from anterior interosseous nerve and artery.

Describe the wrist-joint.

Is ginglymoid, with a hinge-movement, not only of flexion and extension, but adduction and abduction, a combination of these producing circumduction, thus providing a joint with nearly every movement of a ball-and-socket joint, without its insecurity. It is formed between the radius and triangular fibro-cartilage above, and the first row of carpal bones below; its ligaments are,

An *anterior radio-carpal*, passing between the radius above the articular face and the fibro-cartilage, and the first and second rows of carpal bones.

A *posterior radio-carpal*, similarly disposed, except that, above, it is in addition attached to styloid process of ulna.

An *internal lateral*, fan-shaped, passing from styloid process of ulna to ~~pisiform~~, and side and back of cuneiform bones.

An *external lateral*, radiating from tip and front of styloid process of radius to scaphoid, os magnum, and trapezium; the *posterior annular ligament* also assists the wrist ligaments proper; *nerves*, ulnar, median, and posterior interosseous; *arteries*, anterior and posterior carpal arches, radial and ulnar.

Describe the carpal articulations.

Arthrodial, they consist of (1) the joints between the first row; (2) those between the second; and (3) those between the two rows, the *medio-carpal*.

(1) The *pisiform* has a separate capsular ligament, with two bands connecting it with the unciform and base of fifth metacarpal, and a separate synovial membrane; the other three bones of this rows are connected by two

Interosseous ligaments, between scaphoid and lunar, and lunar and cuneiform; and

Two dorsal and *two palmar* ligaments, binding together the same bones.

(2) The bones (4) of second row have

Three interosseous ligaments, connecting the os magnum with the trapezoid externally, the unciform internally; a third between the trapezium and trapezoid; and

Three dorsal and *three palmar ligaments*, passing between the contiguous bony surfaces.

(3) The two rows are united by

A *dorsal and palmar ligament* and *two laterals*, continuous with those of the wrist-joint; the *anterior annular ligament*, passing from the hook of the unciform and from pisiform to the trapezium and scaphoid bones, is an important carpal ligament. The synovial membrane is common to all the carpal joints, except *pisiform*; *nerves*, posterior interosseous, median and ulnar; *arteries*, anterior and posterior carpals of radial and ulnar, carpal of anterior interosseous, interosseous recurrent, carpal of deep palmar arch, and terminal twigs of anterior and posterior interosseous.

Describe the carpo-metacarpal articulations.

That of thumb is *arthrodial*, and enjoys all movements but rotation of the metacarpal on its own axis; it possesses a distinct synovial sac, and its only ligament is the

Capsule attached around the articular surfaces of trapezium and first metacarpal.

The other four metacarpals form *arthrodial* joints with the adjacent carpal bones, with three *dorsal ligaments*, passing from trapezium, trapezoid, and os magnum to second bone; *two ligaments* from os magnum to third; *two ligaments*—one from magnum, the other from cuneiform—to fourth; *one ligament* connecting the fifth metacarpal and unciform.

One palmar ligament passes from trapezium to second metacarpal; *one ligament* each from trapezium, magnum, and unciform to third; *one ligament* connects unciform and fourth bone; *one ligament* passes from unciform to fifth metacarpal.

An *interosseous ligament* connects the contiguous inferior angles of os magnum and unciform with adjacent surfaces of third and fourth metacarpal bones; the *synovial* membrane is that common to intercarpal joints, sometimes the joint formed between fourth and fifth metacarpals and unciform forms a separate synovial sac; *nerves* and *arteries* are same as medio-carpal joint.

Describe the union of the metacarpal bones with one another.

That of the thumb is isolated; the bases of the others are in contact, forming *arthrodial* joints, lined by prolongations of synovial sac of carpus, and are bound together by *palmar*, *dorsal*, and *interosseous ligaments*; a *transverse ligament* binds together their heads.

Describe the metacarpo-phalangeal and interphalangeal articulations.

Ginglymoid, they are connected by ~~*two lateral*~~ ligaments, and an *anterior fibro cartilage*, or ~~*glenoid ligament*~~, except for ~~thumb,~~ where this is replaced by two ~~*sesamoid bones*~~, while behind an expansion of the ~~extensor tendon~~ and some loose areolar tissue completes each little ~~synovial lined capsule;~~ *nerves* and *arteries* from digitals; the *inter-phalangeal* joints resemble the metacarpo-phalangeal in every respect.

Describe the hip-joint.

An *enarthrodial*, but not so freely moving joint as that of the shoulder, it is formed by the head of the femur and the acetabulum on the os innominatum; the ligaments are

3. The *capsular*[9], arising near to acetabular margin, and from outer surface of the transverse ligament, it is attached at the femur, in front to the trochanter major, the spiral line (chiefly formed by anterior intertrochanteric line), and behind, to the neck from one-half to two-thirds of an inch from the posterior intertrochanteric line; the capsule has a ~~thicker anterior segment~~, or superadded band, called

The *ilio-femoral*[8], or ~~*Y-ligament*~~, arising from the anterior inferior iliac spine to be attached to the anterior intertrochanteric line, the outer and inner margins so well marked as to seem like distinct ligamentous bands.

The *ligamentum teres*[10], arising by separate heads from each side of the cotyloid notch externally, passes beneath the transverse ligament covered by the synovial membrane to be attached to a depression on the head of the femur.

Cotyloid, a marginal fibro-cartilage attached all around the acetabular margin and upper surface of the transverse ligament, deepening the socket.

Transverse, a band of unyielding ligamentous fibres bridging the cotyloid notch, converting it into a foramen; *nerves*, from anterior crural, obturator, accessory obturator, nerve to quadratus femoris muscle, great sciatic or lower part of sacral plexus; *arteries*, internal and external circumflex, obturator, gluteal, and sciatic. The *synovial membrane* commonly communicates with the bursa beneath the ilio-psoas muscle.

FIG. 49.



Describe the knee-joint.

Ginglymoid, formed by condyles of femur, upper surface of tibial head, and patella, it has the ~~largest *synovial sac*~~ of any joint, sending a prolongation beneath the popliteus muscle and either another two or three inches up the front of the femur, or communicating with a bursa there placed beneath the quadriceps tendon; the bursa between the semi-membranosus and internal head of gastrocnemius, and inner condyle, usually communicates with the joint, and often one or more of the remaining bursæ near the articulation; its ligaments consist of,

1. An *anterior* or ~~*ligamentum patellæ*~~[8] (Fig. 51), a continuation of the quadriceps tendon attached to the tubercle of the tibia below, which, with the fibrous hood formed by the two vasti tendons attached to the patellar margins and the oblique lines on the head

of tibia extending upward from the tubercle, cover in the joint in front blending with the lateral ligaments.

1. *The posterior* or *ligamentum Winslowii*[8] (Fig. 50), formed of dense, interlacing fibres, having incorporated with it part of the ~~semimembranosus~~ tendon[7] (Fig. 50), arises close above the femoral condyles and intercondyloid notch from one lateral ligament to the other, and is fixed to the border of the outer ~~tibial tuberosity~~, the bone below the popliteal notch and the shaft below the inner tuberosity.

2. *The internal lateral*[6], a strong flat band passing from the depression on inner femoral condyle to the inner border and surface of shaft of tibia for an inch and a half below the head: it is attached to the internal semilunar fibro-cartilage.

FIG. 50.



FIG. 51.



3. *The external laterals*, two in number: < Long Short.

The long[4], a rounded cord attached above to external tuberosity of femur, below to outer part of head of fibula embraced by tendon of ~~biceps~~;

The short, behind and parallel to above, attached above to lower part of outer femoral tuberosity, below to summit of styloid process of fibula; it is intimately connected with capsule; all the preceding ligaments together form a complete

3. *Capsular ligament*, within which are,

The internal ligaments;

The ~~anterior~~ crucial[2], from ~~inner~~ side of depression in ~~front~~ of sp~~ine of tibia~~ to inner back part of outer condyle of femur;

The posterior crucial[3], from back part of depression behind tibial spine and popliteal notch to outer fore part of inner condyle—both crucial ligaments are attached to respective e~~xtremi~~ties of external semilunar fibro-cartilage.

4. *Two semilunar fibro-cartilages*[6-7] (Fig. 51), attached by extremities to the depressions in front and behind the s~~pine of ti~~bia; they serve to deepen the sockets for femoral condyles, and are attached to head of tibia by the *~~coronary~~ ligaments*, short bands passing between their outer margins and the tibial head, they are also connected to one another, in front, by the small band-like *transverse ligament*[4]; stretching from front of joint, below patella, to front of intercondyloid notch, is the triangular fold of synovial membrane called the

[illegible] *Ligamentum mucosum*, from whose sides extend upward and outward to the sides of the patella, the fringe-like folds termed *ligamenta alaria; nerves*, internal and external popliteal, anterior crural, obturator; *arteries*, five articular from popliteal, anastomotica magna from femoral, recurrent anterior tibial; in addition to bursa, already mentioned there is one over patella and its ligament, one beneath ligamentum patellæ, one between inner hamstrings and tibial head, and one or more, not always present, beneath contiguous tendons.

Describe the ~~inferior~~ Sup. tibio-fibular articulation.

Arthrodial, formed between facets on fibular head and outer tuberosity of tibia, its ligaments are really a *capsule*, described as,

Anterior superior tibio-fibular[10] (Fig. 51), directed obliquely upward and inward from fibula to tibia,

Posterior superior tibio-fibular[10] (Fig. 50), similarly disposed behind; the synovial membrane communicates with that of knee in twenty-five per cent. of specimens; *nerves*, from external popliteal; *arteries*, some of those of knee.

Describe the so-called middle tibio-fibular articulation.

The contiguous borders of the bones are connected by the *inter-*

osseous membrane[11] (Fig. 51), deficient above for passage of anterior tibial artery, below it is perforated by anterior peroneal vessels; its fibres pass from fibula upward to tibia.

Describe the inferior tibio-fibular joint.

Arthrodial, formed superiorly by the rough surfaces of the contiguous bones, below by small cartilage-coated surfaces; its synovial membrane is part of that of the ankle.

The ligaments are:

Inferior interosseous, really thicker part of interosseous membrane passing between the rough surfaces of the two bones.

Anterior and *posterior inferior tibio fibular*, oblique fibrous bands binding together the contiguous bony surfaces in front and behind.

Transverse, a narrow band passing from external malleolus to tibia behind the joint; *nerves*, those of ankle; *arteries*, some of those of ankle.

Describe the ankle-joint.

Ginglymoid, formed by lower end of tibia with its malleolus and the external malleolus, and the upper surface and sides of the astragalus; its ligaments are the,

FIG. 52.



FIG. 53.



Anterior[9] (Fig. 53), connecting the articular margins of the tibia and fibula with the rough upper surface of the astragalus.

Posterior[2] (Fig. 52), thin and imperfect, extends between the external malleolus, back of lower end of tibia and posterior tibio-fibular ligament above, and the posterior surface of astragalus from one lateral ligament to the other below.

Internal lateral[6] (Fig. 52), (deltoid) radiates from lower border of inner malleolus to astragalus behind and below, in front to scaphoid, and passes almost perpendicularly to the os calcis.

External lateral[6], [7], [8] (Fig. 53), composed of three fasciculi from anterior margin, apex, and back of external malleolus, it is attached to the astragalus and os calcis: *nerves*, internal saphenous and anterior tibial; *arteries*, anterior and posterior tibials, anterior and posterior peroneals.

Describe the tarsal joints.

Those of the first row are arthrodial: the astragalus and calcis have three ligaments;

External calcaneo astragaloid, from outer surface of astragalus to same of calcis;

The posterior calcaneo-astragaloid connects the posterior end of astragalus with upper surface of calcis;

The interosseous, the chief bond, filling up the grooves on the under surface of astragalus and upper of os calcis; there are two synovial sacs, one posterior, the other anterior continued between astragalus and scaphoid: *nerves*, posterior tibial or plantar; *arteries*, posterior tibial, tarsal, external malleolar (from ant. tibial), and terminal twigs of peroneal.

Those of second row, scaphoid, cuboid, and cuneiforms have,

Dorsal ligaments, small bands connecting the contiguous bones, and,

Plantar ligaments, similarly disposed; also four

Interosseous ligaments, connecting the sides of the scaphoid and cuboid, the internal and middle cuneiforms, the middle and external cuneiforms, the external cuneiform and cuboid; *nerves*, anterior tibial, internal and external plantars; *arteries*, metatarsal and plantars.

Those between the two rows, or medio-tarsal. (1) The *calcaneo-scaphoid*, arthrodial, has

The *superior calcaneo-cuboid ligament*, connecting the dorsal surfaces of calcis and cuboid.

The *internal* (*interosseous*) calcaneo-cuboid, connecting inner under part of front of calcis with inner postero-inferior angle and contiguous part of cuboid.

The *long calcaneo-cuboid* (*plantar*) arises from under surface of calcis between posterior tubercles and anterior tubercle, to be attached to oblique ridge of cuboid and bases of second, third, fourth, and fifth metatarsal bones, completing canal for the long peroneal tendon.

The *short plantar* arises from the anterior calcaneal tubercle and the bone in front, to be attached to the under surface of cuboid behind the ridge, except the outer angle; the synovial sac is distinct from that of other tarsal joints.

The (2) astragalo-scaphoid articulation is the only enarthrodial (ball-and-socket) joint of tarsus, and communicates with the anterior calcaneo-astragaloid; its ligaments are,

The *superior calcaneo-scaphoid*, dense and thick, extending from anterior internal extremity of calcis to under surface of scaphoid.

The *inferior calcaneo-scaphoid*, passing obliquely forward from neck of astragalus to upper surface of scaphoid; *nerves* of medio-tarsal joint are, external branch of anterior tibial, sometimes musculo-cutaneus or external plantar; *arteries*, anterior tibial, tarsal, metatarsal, or external plantar.

Describe the tarso-metatarsal articulations.

Arthrodial, between the three cuneiforms and the cuboid, and the bases of the five metatarsals; their ligaments are,

Dorsal, one for first metatarsal and internal cuneiform; one from each cuneiform to second; one from external cuneiform to third; one each from cuboid to fourth and fifth metatarsals.

Plantar, more irregularly disposed ligamentous fibres.

Interosseous, one connecting outer extremity of inner cuneiform to adjacent angle of second metatarsal; one between outer cuneiform and angle of second metatarsal; one connecting outer angle of external cuneiform with side of third metatarsal; *nerves*, anterior tibial and plantars; *arteries*, from dorsalis pedis, metatarsal, and deep plantar arch.

Describe the articulations of the metatarsal bones with one another.

Except the first, their bases are bound together by *dorsal*, *plantar*, and *interosseous* ligaments; the distal extremities are united by the *transverse metatarsal ligament.*

Describe the metatarso-phalangeal and inter-phalangeal articulations.

They resemble in all respects those of hand, which see.

How many distinct synovial sacs have the tarsus and metatarsus?

Six, viz.: (1) Between calcis and astragalus, posterior to interosseus ligament; (2) in front of same ligament, between calcis and astragalus, also between astragalus and scaphoid; (3) between calcis and scaphoid; (4) between scaphoid, cuboid, and cuneiforms, communicating between the middle and external cuneiforms with the joints formed between those bones and the bases of the second and third metatarsals; (5) between the sides and bases of the fourth and fifth metatarsals and the cuboid; (6) between the base of first metatarsal and internal cuneiform.

Muscles and Fasciæ.

Describe a muscle.

It is a structure connected with bones, cartilages, ligaments, or skin, directly or through the medium of a tendon or aponeurosis, whose structure is capable of shortening when irritated mechanically or by nerve stimulus, this property being called *muscular contractility* or *irritability.*

How are muscles divided?

According to their form into narrow, broad, penniform (when their fibres converge like the plumes of a pen to *one* side of the tendon); *bipenniform* when so arranged on both sides of a tendon, radiated, etc.

Is there only one variety of muscular tissue?

No, there are *two*, one variety consisting of prismatic fasciculi about $\frac{1}{400}$th of an inch in diameter, marked by transverse striæ, each bundle surrounded by a sheath or *perimysium*; each fasciculus is formed of a number of *fibrillæ*, $\frac{1}{18000}$th of an inch in diameter, surrounded by a tubular, transparent elastic membranous *sarcolemma;* every fibril is striated, and consists of a row of minute particles, Bowman's "*sarcous elements;*" this variety of tissue forms the *voluntary, striped muscles,* or those of *animal life,* being under the control of the will; the *involuntary, unstriped,* or the muscles of *organic life,* are not under the control of the will, and consist of elongated, spindle shaped, flattened, nucleated cells, $\frac{1}{600}$th to $\frac{1}{300}$th of an inch long, $\frac{1}{4500}$th to $\frac{1}{3500}$th broad, and are held together in bundles by a cement substance containing a few connective-tissue corpuscles, the bundles being further collected into larger fasciculi, or flattened bands, bound together by ordinary areolar tissue; this kind of muscle is that found forming the contractile coats of the stomach, intestines, bladder, arteries, veins, lymphatics, ureters, urethra, iris, ciliary body, etc.; when irritated a part contracts, slowly relaxes, while another contiguous portion is contracting, which action being continued produces the *vermicular* (worm-like), peristaltic movements of the intestines. The spontaneously coagulable albuminous substance composing muscle is called *myosin;* so-called *syntonin* is a modification produced by chemical agents.

What are tendons and aponeuroses?

Tendons are glistening white cords or bands, of white inelastic fibrous tissue, almost without blood- or nerve-supply, and connect the muscular tissue with the part to be moved; *aponeuroses* are membranous expansions of the same tissue serving similar purposes.

What are the fasciæ?

They are laminæ of fibrous or fibro-areolar tissue investing the soft tissues; the superficial fasciæ are usually fibro-areolar, lie beneath the skin, and are loaded with fat; the deep fasciæ resemble aponeuroses, are dense and inelastic, ensheathing muscles, or serving for their attachment, and tend to preserve the form of the part, since partitions descend between various muscles to become attached to the periosteum of the osseous framework.

To what are the tendons attached?

To the periosteum and perichondrium with which they become blended, to the subcutaneous tissue, and to ligaments.

What is meant by the origin and insertion of a muscle?

The *origin* is the most fixed and central point from which the muscle acts; the *insertion* the movable point to which the muscular force is directed; but, in many muscular acts the part usually described as the *insertion* becomes the fixed point and the *origin*, the movable one.

Do not some muscles have double origins, although single insertions?

Yes; as the occipito-frontal, digastric, and omo-hyoid.

356

Muscles of the Head.

Describe the origin, insertion, action, and nerve supply of the following muscles.

Occipito-frontalis[1]: *origin*, occipital portion, outer two-thirds of superior curved line of the occipital bone and mastoid process of the temporal; frontal portion, from pyramidalis nasi, corrugator supercilii, and orbicularis palpebrarum; *insertion*, into an aponeurosis covering the vertex of the skull; *action*, raises eyebrows and transversely wrinkles forehead; *nerves*, facial, posterior auricular branch of same, sometimes the small occipital.

Attolens aurem[4]: *origin*, occipito-frontalis aponeurosis; *insertion*, superior portion of pinna of ear; *action*, raises pinna; *nerve*, occipitalis minor.

696

Atrahens aurem: *origin*, lateral margin of occipito-frontalis tendon; *insertion*, a projection on front of helix; *action*, draws the pinna forward and upward; *nerve*, facial.

Retrahens aurem[5]: *origin*, mastoid portion of temporal; *insertion*, lower part of concha; *action*, draws ear back; *nerve*, posterior auricular branch of facial.

Orbicularis palpebrarum[6]: *origin*, internal angular process of the frontal, nasal process of superior maxillary bone, and the anterior surface and margins of the tendo oculi; *insertion*, skin of eyelids and contiguous portions of forehead, temple, and cheek, blending with the occipito-frontal and corrugator supercilii muscles; *action*, closes eyelids; *nerve*, facial.

Forehead.

Corrugator supercilii: *origin,* inner end of superciliary ridge; *insertion,* under surface of orbicularis palpebrarum; *action,* draws eyebrow downward and inward; *nerve,* facial.

lids.

Tensor tarsi (Horner's muscle): *origin,* crest of lachrymal bone; *insertion,* by two slips into tarsal cartilages near puncta; *action,* compresses lachrymal sac, and keeps puncta in contact with globe; *nerve,* facial.

Levator palpebræ superioris[8]: *origin,* lesser wing of sphenoid; *insertion,* upper border of superior tarsal cartilage; *action,* elevates upper lid; *nerve,* third cranial (motor oculi).

FIG. 54.



Nose

Pyramidalis nasi[3]: *origin,* occipito-frontalis; *insertion,* into compressor naris; *action,* depresses inner angle of the eyebrow; *nerve,* facial.

Levator labii superioris alæque nasi[7]: *origin,* upper part of nasal process of superior maxillary bone; *insertion,* the cartilage of ala of nose, and into upper lip blending with orbicularis and levator labii; *action*, draws upward the upper lip, and dilates nostril; *nerve*, facial.

✓ ***Dilator naris anterior:*** *origin*, cartilage of ala; *insertion*, integument near its margin; *action*, dilates nostril; *nerve*, facial.

✓ ***Dilator naris posterior:*** *origin*, margin of nasal notch of superior maxilla and the sesamoid cartilages; *insertion*, skin near margin of nostril; *action*, dilates nostril; *nerve*, facial.

Compressor nasi[9]: *origin*, above and external to incisive fossa of superior maxilla; *insertion*, fibro-cartilage of nose, and is continuous with its fellow and the aponeurosis of pyramidalis nasi; *action*, dilates nostril, by increasing breadth of nose; *nerve*, facial.

✓ ***Compressor narium minor***: *origin*, alar cartilage; *insertion*, skin of tip of nose; *action*, dilates nostril; *nerve*, facial.

Depressor alæ nasi: *origin*, incisive fossa of superior maxilla; *insertion*, septum and back of ala; *action*, narrows nostril; *nerve*, facial.

[illegible]

Levator labii superioris[8]: *origin*, lower margin of orbit above infra-orbital foramen; *insertion*, muscular substance of upper lip; *action*, elevates lip; *nerve* facial.

Levator anguli oris[16]: *origin*, canine fossa of superior maxilla; *insertion*, angle of mouth; *action*, elevates angle of mouth; *nerve*, facial.

Zygomaticus major[11]: *origin*, malar bone; *insertion*, angle of mouth; *action*, elevates angle of lip; *nerve*, facial.

Zygomaticus minor[10]: *origin*, malar bone anteriorly; *insertion*, angle of mouth; *action*, same as zygomaticus major; *nerve*, facial.

Levator labii inferioris (***levator menti***): *origin*, incisive fossa of lower jaw; *insertion*, skin of lower lip; *action*, elevates lower lip; *nerve*, facial.

Depressor labii inferioris[17-18] (***quadratus menti***): *origin*, external oblique line of lower jaw; *insertion*, skin of lower lip; *action*, depresses lower lip; *nerve*, facial.

Depressor anguli oris[14]: *origin*, external oblique line of lower jaw; *insertion*, angle of mouth; *action*, depresses angle of mouth; *nerve*, facial.

Orbicularis oris[15]: *origin*, nasal septum and superior and inferior maxillary borders by accessory fibres called accessorii orbicularis superioris and inferioris, and naso-labialis; *insertion*, the buccinator, and other muscles converging to the mouth; *action*, closes mouth; *nerve*, facial.

Buccinator[13]: *origin*, behind, from the pterygo-maxillary ligament, above and below, the external surfaces of alveolar processes of the three molar teeth; *insertion*, orbicularis oris; *action*, compresses cheek; *nerve*, facial and buccal branch of inferior maxillary.

Risorius (***Santorini***): *origin*, fascia over masseter muscle; *insertion*, angle of mouth; *action*, laughing muscle; *nerve*, facial.

MASTICATION,

Masseter[12]: *origin*, the superficial portion, from the malar process of the superior maxilla and the anterior two-thirds of the lower border of the zygomatic arch, the deep portion, from the posterior third of the lower border and the inner surface of the zygomatic arch; *insertion*, upper half of ramus and outer surface of coronoid process of lower jaw; *action*, raises, protracts, and retracts lower jaw, a masticatory muscle; *nerve*, inferior maxillary.

Temporal: *origin*, temporal fossa and fascia; *insertion*, coronoid process of lower jaw; *action*, raises and retracts lower jaw, a muscle of mastication; *nerve*, inferior maxillary.

Internal pterygoid: *origin*, inner surface of external pterygoid plate and grooved surface of palate bone, the tuberosity of the palate and superior maxillary bones; *insertion*, lower back part of inner side of ramus as high as dental foramen, and angle of jaw; *action*, draws lower jaw forward, a muscle of mastication; *nerve*, inferior maxillary.

External pterygoid: *origin*, pterygoid ridge on greater wing of sphenoid, the bone included between it and base of pterygoid process, the outer surface of external pterygoid plate; *insertion*, depression in front of neck of condyle of lower jaw and inter-articular fibro-cartilage; *action*, draws jaw forward, a triturating masticatory muscle; *nerve*, inferior maxillary. Sometimes the internal maxillary artery passes between the two heads of this muscle; sometimes pierces it; often lies below it.

P. 351.

Muscles of the Neck.

Platysma myoides: *origin*, clavicle, acromion, and fascia of pectoral, deltoid, and trapezius muscles; *insertion*, lower jaw beneath oblique line, angle of mouth, cellular tissue of face; *action*, wrinkles skin of neck, depresses jaw and angle of lip; *nerves*, facial, superficial branches cervical plexus.

Sterno-cleido-mastoid[11] (Fig. 55): *origin*, upper front part of sternum, inner third upper border of clavicle; *insertion*, mastoid process of temporal and outer two-thirds superior curved line of occipital bone; *action*, acting *singly*, flexes head sideways, and rotates chin to opposite side, *both* acting flex head on neck, and then on chest; *nerves*, spinal accessory, deep branches of cervical plexus.

FIG. 55.



Sterno-hyoid[14]: *origin*, riband like, from posterior surface of sternum and inner end of clavicle; *insertion*, body of hyoid bone; *action*, depresses hyoid bone; *nerve*, filaments from loop between descendens and communicans noni.

Sterno-thyroid[15]: *origin*, posterior surface of sternum and cartilage of first rib; *insertion*, oblique line on ala of thyroid cartilage; *action*, depresses larynx; *nerve* same as sterno-hyoid.

Thyro-hyoid[16]: *origin*, oblique line of thyroid cartilage; *insertion*, lower border of body and greater cornu of hyoid bone; *action*, may elevate larynx when hyoid is fixed; *nerve*, hypoglossal.

Omo-hyoid[18]: *origin*, from upper border of scapula near suprascapular notch, perhaps, also, from transverse ligament; *insertion*, body of hyoid bone, its *central tendinous portion* being held down by a process of deep cervical fascia prolonged down to be attached to cartilage of first rib; *action*, depresses hyoid bone, larynx, and draws them backward to one or other side; *nerve*, from loop between descendens and communicans rami.

Digastric[12]: *origin*, has two fleshy bellies with an intermediate rounded tendon, the posterior[1] is attached to digastric groove of mastoid process of temporal, the anterior[2] to depression on inner side of lower border of jaw near symphysis; *insertion*, tendon perforates stylo-hyoid, and is held to side of body of hyoid bone by aponeurotic loop lined with synovial membrane; *action*, raises the hyoid bone; if this be fixed by its muscles, depresses lower jaw; *nerves*, posterior belly by facial, anterior by mylo-hyoid branch of inferior dental.

Stylo-hyoid[4]: *origin*, middle of outer surface of styloid process of temporal; *insertion*, body of hyoid at junction with greater cornu, it is perforated by digastric tendon; *action*, same as digastric; *nerve*, facial. Wharteth

Mylo-hyoid[5]: *origin*, mylo-hyoid ridge of lower jaw; *insertion*, body of hyoid bone and median raphé; *action*, same as digastric and stylo-hyoid; *nerve*, mylo-hyoid branch of inferior dental; it forms the muscular floor of the mouth.

Genio-hyoid[6]: *origin*, inferior genial tubercle of lower jaw; *insertion*, body of hyoid bone; *action*, same as mylo-hyoid; *nerve*, hypoglossal. → VERTEBRAL · p122.

p.727 GERRISH

Genio-hyo-glossus[7]: *origin*, superior genial tubercle of lower jaw; *insertion*, body of hyoid bone, side of pharynx and tongue from base to apex; *action*, protrudes tongue by posterior and inferior fibres, retracts it by anterior fibres, both muscles acting render tongue concave from side to side, as in ~~sucking~~; *nerve*, hypoglossal.

Hyo-glossus[8]: *origin*, body, lesser, and whole length of greater cornu of hyoid bone; *insertion*, side of tongue; *action*, draws down sides of tongue, making it ~~convex~~; *nerve*, hypoglossal.

Stylo glossus*: origin*, near centre of antero-external surface of styloid process and stylo-maxillary ligament; *insertion*, side of tongue and hyo-glossus muscle; *action*, draws tongue upward and backward; *nerve*, hypoglossal.

Lingualis*: origin*, forms bulk of tongue lying between hyo-glossus and genio-hyo glossus on each side, running from base to apex, some fibres being attached to hyoid bone; *action*, renders tongue convex from before backward; *nerve*, chorda tympani.

1ERRISN 718-19

Palatal Region.

Levator palati*: origin*, under surface of apex of petrous portion of temporal bone and contiguous portion of cartilaginous Eustachian tube; *insertion*, into posterior surface of soft palate; *action*, elevates soft palate; *nerve* facial through petrosal nerve of Vidian.

Tensor palati*: origin*, scaphoid fossa, outer side of Eustachian tube, spine of sphenoid, edge of tympanic plate; *insertion*, after reflection around hamular process, into forepart of aponeurosis of soft palate and under surface of palate bone; *action*, renders soft palate tense; *nerve*, branch from otic ganglion.

Azygos uvulæ*: origin*, posterior nasal spine and palatal aponeurosis; *insertion*, uvula; *action*, raises uvula; *nerve*, facial through petrosal of Vidian; it is not a *single* muscle as the name implies.

Palato glossus (anterior pillar of fauces): *origin*, anterior surface of palate external to uvula; *insertion*, side and dorsum of tongue; *action*, as one of its names implies, constrictor isthmii faucium; *nerve*, palatine branches Meckel's ganglion.

Palato pharyngeus (posterior pillar of fauces): *origin*, soft palate; *insertion*, posterior border of thyroid cartilage and side of pharynx; *action*, closes posterior fauces; *nerve*, same as for palato-glossus.

Vertebral Region.

Rectus capitis anticus major*: origin*, by four tendinous slips from anterior tubercles of transverse processes of third, fourth, fifth, and sixth cervical vertebræ; *insertion*, basilar process of occipital bone; *action*, flexes head; *nerves*, suboccipital, deep branches of cervical plexus

Rectus capitis anticus minor: *origin*, front of lateral mass of atlas and root of its transverse process; *insertion*, basilar process of occipital bone behind preceding muscle; *action*, flexes head; *nerves*, suboccipital, deep branches of cervical plexus.

Rectus lateralis: *origin*, upper surface transverse process of atlas; *insertion*, jugular process of occipital bone; *action*, draws head laterally when one acts, flexes when both act; *nerves*, suboccipital, deep branches of cervical plexus.

Longus colli: *origin*, *superior oblique portion* from anterior transverse tubercles of third, fourth, and fifth cervical vertebræ; *insertion*, tubercle on anterior arch of atlas; the *inferior oblique portion arises* from bodies of first two or three dorsal vertebræ; *insertion*, transverse processes of fifth and sixth cervical vertebræ; *vertical portion* extends between bodies of upper three dorsal and lower three cervical, and those of second, third, and fourth cervical vertebræ; *action*, flexes and slightly rotates cervical spine; *nerves*, anterior branches from lower cervical nerves.

Scalenus anticus[20] (Fig. 55): *origin*, tubercle on upper surface of first rib; *insertion*, anterior tubercles of transverse processes of third, fourth, fifth, and sixth cervical vertebræ; *action*, lateral flexion of cervical spine or elevation of ribs; *nerves*, anterior branches lower cervical nerves.

Scalenus medius[21]: *origin*, upper surface of first rib behind subclavian groove; *insertion*, posterior tubercles of transverse processes of lower six cervical vertebræ; *action*, similar to preceding; *nerves*, anterior branches from lower cervical.

Scalenus posticus: *origin*, outer surface of second rib; *insertion*, posterior tubercles of lower two or three cervical vertebræ; *action*, flexes neck laterally, or elevates second rib; *nerves*, anterior branches from lower cervical.

Muscles of the Back

First layer:

1° 3 layers = 3 ead...

Trapezius[1]: *origin*, inner third of superior curved line of occipital bone, ligamentum nuchæ, spinous processes of seventh cervical and all dorsal vertebræ, and corresponding portion of supra-spinous ligament; *insertion*, outer third posterior border of clavicle, inner margin of acromion process, the superior lip of crest of spine of scapula, and tubercle at its inner extremity; *action*, one acting draws head to its own side, both acting, draw head backward, and with head fixed, elevate point of shoulder; middle and lower fibres partially rotate scapula on chest; *nerves*, spinal accessory, deep branches of cervical plexus.

Ligamentum ① *nuchæ* (a thin band of condensed cellulo-fibrous membrane): *origin*, external occipital protuberance; *insertion*, spinous processes of all cervical vertebræ except atlas; a rudiment of strong elastic ligament sustaining head in herbivora, etc.

Latissimus dorsi[4]: *origin*, by an aponeurosis from spinous processes of lower six dorsal vertebræ, those of lumbar and sacral vertebræ and supra-spinous ligament, also external lip of iliac crest behind external oblique, and from three or four lower ribs; *insertion*, inner lip and bottom of bicipital groove of humerus; *action*, acting on humerus, draws it downward and backward while rotating it inward, with fixed arms, raises lower ribs in forcible inspiration, assists greater pectorals and abdominal muscles in drawing trunk forward as when climbing, using crutches, etc.; *nerve*, long subscapular nerve.

Second layer: (1-5)

Levator anguli scapulæ[10]: *origin*, by three or four tendinous slips from posterior tubercles of transverse processes of three or four upper cervical vertebræ; *insertion*, posterior border of scapula between superior angle and triangular surface at root of spine; *action*, raises scapular angle; *nerves*, filaments from fifth cervical and from deep branches cervical plexus.

Rhomboideus minor[11]: *origin*, ligamentum nuchæ, and spinous processes of seventh cervical and first dorsal vertebræ; *insertion*, margin of triangular surface at root of spine of scapula; *action*, draws inferior angle backward and upward; *nerve*, fifth cervical.

Rhomboideus major[12]: *origin*, spinous processes of upper four or 1-5) five dorsal vertebræ and supra-spinous ligament; *insertion*, tendinous arch stretched from triangular surface at root of spine of scapula to inferior angle, the arch being connected with scapular border by thin membrane; *action*, draws inferior angle upward and backward; *nerve*, fifth cervical.

① (Nape of neck

Third layer:

Serratus posticus superior: *origin*, by thin aponeurosis from ligamentum nuchæ and spinous processes of seventh cervical and

FIG. 56.



two or three upper dorsal vertebræ; *insertion*, by fleshy digitations into upper borders of second, third, fourth, and fifth ribs a little beyond their angles; *action*, elevates ribs during inspiration; *nerves*, external posterior branches of cervical nerves

2-5 Ribs

Serratus posticus inferior[16]: *origin*, by thin aponeurosis from spinous processes of last two dorsal and upper two or three lumbar

2 - 3 = 5

vertebræ and interspinous ligaments; *insertion*, by four digitations into lower borders of four lower ribs, beyond their angles; *action*, depresses ribs in expiration; *nerves*, external branches of dorsal nerves.

— 1 — 6

① ***Splenius capitis et colli***[13] [14]: *origin*, tendinous from lower half of ligamentum nuchæ, the spinous processes of last cervical and of upper six dorsal vertebræ, and supraspinous ligament; *insertion*, (*S. capitis*[13]) mastoid process and rough surface beneath superior curved line of occipital bone (*S. colli*[14]), posterior tubercles of transverse processes of upper three or four cervical vertebræ; *action*, together, draw head directly backward, singly, flex head laterally and slightly rotate to same side; *nerves*, external posterior branches of cervical nerves.

Fourth layer: 3 div.

Erector spinæ[1], [2], [3], [4], [5], [6] (Fig. 57): *origin*, sacro-iliac groove, the anterior surface of lumbo-sacral tendon, which is attached internally to spines of sacrum, the spinous processes of lumbar and three lower dorsal vertebræ and supraspinous ligament externally, the back part of inner lip of iliac crest, with the rudimentary transverse processes of the sacrum and great sacro-sciatic ligament; this muscle opposite last rib divides into ~~sacro-lumbalis~~ ilio costalis and longissimus dorsi muscles; *action*, maintains spine erect and bends body backward; *nerves*, external posterior branches of lumbar and dorsal nerves.

Sacro-lumbalis[1], [2]: *origin*, is part of erector spinæ; *insertion*, by six or seven tendons into angles of six lower ribs; *action*, same as erector spinæ; *nerves*, same as erector spinæ.

Musculus accessorius ad sacro-lumbalem: *origin*, by flattened tendons from angles of six lower ribs; *insertion*, angles of six upper ribs; *action*, same as erector spinæ; *nerves*, those of erector spinæ.

Cervicalis ascendens: *origin*, angles of four or five upper ribs; *insertion*, posterior tubercles of transverse processes of fourth, fifth, and sixth cervical vertebræ; *action*, keeps neck erect; *nerves*, external posterior branches of cervical nerves.

Longissimus dorsi[3]: *origin*, is part of erector spinæ; *insertion*, posterior surface of lumbar transverse processes, tubercles at back of articular processes, tips of transverse processes of all the dorsal

① = Bandage

vertebræ, and from six to eleven ribs between their tubercles and angles; *action* and *nerves* same as erector spinæ.

iddo 3 ***Transversalis colli***[6],[9]: *origin*, tendinous from summits of six upper dorsal transverse processes; *insertion*, posterior tubercles of transverse processes from second to sixth cervical vertebræ; *action*, keeps neck erect; *nerves*, external posterior branches of cervical nerves.

FIG 57.



Trachelo-mastoid[7]: *origin*, by tendons from transverse processes of third, fourth, fifth, and sixth dorsal vertebræ and articular processes of the three or four lower cervical; *insertion*, posterior margin of mastoid process; *action*, flexes head laterally; *nerves*, external posterior branches of cervical nerves.

Spinalis dorsi: *origin*, by tendons from spinous processes of last two dorsal and first two lumbar vertebræ; *insertion*, by separate tendons into spinous processes of from four to eight dorsal vertebræ; *action*, erects spine; *nerves*, external posterior branches of dorsal nerves.

Spinalis colli: *origin*, fifth and sixth cervical spinous processes, also sometimes those of first and second dorsal (this muscle is sometimes absent); *insertion*, spinous process of axis, occasionally those of the two vertebræ below; *action*, extends cervical spine; *nerves*, external posterior branches of cervical nerves.

5° Layer. ***Complexus***[8]: *origin*, 6 by seven tendons from tips of transverse processes of upper three dorsal and seventh cervical vertebræ, and articular processes of the three cervical vertebræ above this; *insertion*, internally between curved lines of occipital bone; *action*, both draw head backward, singly, Semisp Capiti lateral flexion with rotation turning face to opposite side; *nerves*, internal posterior branches of cervicals, sub-occipital and great occipital: (the biventer cervicis being usually blended with this muscle will receive no separate description).

Fifth layer:

Semispinalis dorsi[10]: *origin*, by small tendons from transverse 3 processes of from tenth or eleventh to fifth or sixth dorsal vertebræ; *insertion*, by five or six tendons, into spinous processes of upper four dorsal and lower two cervical vertebræ; *action*, erects spinal column; *nerves*, internal posterior branches of dorsal nerves.

Semispinalis colli[11]: *origin*, tendinous and fleshy from upper four dorsal transverse processes and articular processes of lower four cervicals; *insertion*, second, third, fourth, and fifth cervical spinous processes; *action*, erects cervical spine; *nerves*, internal posterior branches of cervical nerves.

Multifidus spinæ[16]: *origin*, back of sacrum, aponeurosis of erector spinæ (sacral region); posterior superior iliac spine, posterior sacro-iliac ligaments (iliac regions); articulating processes (lumbar and cervical regions); transverse processes (dorsal region); *insertion*, lamina and spinous process of third or fourth vertebra above (most superficial), second or third above (middle layer); two contiguous vertebræ (deepest layer); keeps spine erect and rotates it; *nerves*, internal posterior branches of cervical, dorsal, lumbar, and sacral nerves. (Cont. below of Semi-spinalis)

Rotatores spinæ (eleven pairs in dorsal region only): *origin*, each v from upper back part of transverse process; *insertion*, lower outer part of lamina of vertebra above; *action*, as name implies; *nerves*, internal posterior dorsal branches.

✓ ***Supra-spinales:*** *origin* and *insertion*, cervical spinous processes; *action*, extend cervical spine; *nerves*, branches of cervical.

Inter-spinales: *origin*, variable in number, there are usually six cervical pairs commencing between apices of second and third vertebral spines; two, occasionally three dorsal pairs, between first and second, eleventh and twelfth, sometimes second and third vertebræ; four lumbar pairs; sometimes a pair between last dorsal and first lumbar, and fifth lumbar and sacrum; *action*, extend spine; *nerves*, internal posterior nerves of regions.

Extensor coccygis: *origin,* last bone of sacrum, or first of coccyx; *insertion,* lower part of coccyx, behind; *action,* as named; *nerve,* posterior sacral.

Intertransversales[18]: *origin* and *insertion,* between anterior and posterior tubercles of transverse processes in cervical[7], occupy intertransverse spaces in other regions (twelve dorsal and four lumbar); *action,* lateral flexion, singly, steady spine when both sides act; *nerves,* posterior spinal branches of each region.

Rectus capitis posticus major[13]: *origin,* spinous process of axis; *insertion,* inferior curved line of occipital bone and surface below; rotates head; *nerves,* suboccipital and great occipital.

Ir. occ.
3.

Rectus capitis posticus minor[12]: *origin,* tubercle on posterior arch of atlas; *insertion,* rough surface beneath inferior curved line, nearly as far as foramen magnum; *action,* extends head; *nerves,* suboccipital and great occipital.

Obliquus capitis inferior[15]: *origin,* apex of spinous process of axis; *insertion,* apex of transverse process of atlas; *action,* rotates atlas and head; *nerves,* suboccipital and great occipital.

Obliquus capitis superior[14]: *origin,* tendinous from upper part of transverse process of atlas; *insertion,* between curved lines of occipital bone external to complexus; *action,* extends head; *nerves,* suboccipital and great occipital.

Muscles of the Abdomen.

External oblique[12]: *origin,* by eight fleshy digitations from external surface and lower borders of the eight inferior ribs; *insertion,* anterior half of outer lip of iliac crest, by an aponeurosis joining its fellow of opposite side it forms the *linea alba,* a median tendinosis raphé, is connected with ensiform cartilage, symphysis, spine, and pectineal line of pubes; it is continuous above with aponeurosis of pectoralis major, below it forms a broad infolded band, continuous with fascia lata, stretching from anterior iliac spine to pubic spine, *i. e., Poupart's ligament,* reflected from which into the pectineal line, is a portion called *Gimbernat's ligament;* just above crest of pubis is the triangular external abdominal ring, its external border being Poupart's ligament, its inner, aponeurotic fibres, both called *pillars of the ring;* *action,* compresses viscera,

upper [illegible]
S. [illegible]

flexes thorax on pelvis, or flexes pelvis on thorax, according to which is the fixed point, also assists expiration; *nerves*, lower intercostal, ilio-hypogastric, ilio-inguinal.

Internal oblique[18]: *origin*, fleshy from outer half of Poupart's ligament, anterior two-thirds of middle lip of iliac crest, and posterior lamella of lumbar fascia; *insertion*, crest and pectineal line of pubis with transversalis muscle—forming *conjoined tendon* part of inner boundary of internal abdominal ring—cartilages of lower four ribs, and by an aponeurosis, which splits for its upper

Fig. 58.



three-fourths to enclose the rectus muscle, into linea alba, the anterior layer blending with aponeurosis of external oblique, the posterior with that of transversalis muscle, the undivided lower fourth passing in front of rectus; *action* and *nerves*, same as external oblique.

Transversalis: *origin*, fleshy from outer third of Poupart's ligament, anterior three-fourths of inner lip of iliac crest, inner surface of six lower costal cartilages, and by a broad aponeurosis from lumbar spinous and transverse processes; *insertion*, with internal oblique as the *conjoined tendon* into pubic crest and pectineal line, by its aponeurosis into linea alba, the upper three-fourths passing *behind* rectus abdominis, the remainder *in front* of that muscle; *action* and *nerves* similar to external oblique.

Lumbar fascia: the vertebral aponeurosis of the transversalis divides into an anterior thin layer attached to front of lumbar transverse processes and to lower margin of last rib, thus forming the *ligamentum arcuatum externum*; a middle, stronger layer, attached to apices of transverse processes; and a posterior layer attached to apices of spinous processes; the quadratus lumborum lies between anterior and middle layers, the erector spinæ between the middle and posterior; the last receives the attachment of the internal oblique, and blended with the aponeuroses of the serratus posticus inferior and latissimus dorsi, forms the lumbar fascia.

Rectus abdominis[16]: *origin*, by two tendons from pubic crest and ligaments over symphysis pubis; *insertion*, cartilages of fifth, sixth, and seventh ribs; this muscle is enclosed in a sheath, the upper three-fourths in front formed by the aponeurosis of the external oblique and half of that of the internal oblique, behind for the same extent of the transversalis aponeurosis with the posterior division of that of the internal oblique; at the lower fourth the aponeuroses of all these muscles pass *in front* of the rectus; from two to five tendinous intersections traverse the muscle called *lineæ transversæ*, while the *lineæ semilunares*, two curved tendinous lines corresponding to the outer border of the recti muscles, extend on either side from the eighth costal cartilage to the pubes; *action*, flexes thorax on pelvis, and *vice versa*, also compresses abdominal viscera; *nerves*, same as external oblique.

INT.
EXT.
TRAN

Pyramidalis: *origin*, tendinous from pubic bone and anterior pubic ligament, lying in front of, but in same sheath with rectus; *insertion,* linea alba midway between pubes and umbilicus; *action,* a tensor of linea alba; *nerves,* same as those of external oblique.

Quadratus lumborum[19] (Fig. 57): *origin,* ilio-lumbar ligament, adjacent two inches of iliac crest, and upper borders of transverse processes of third, fourth, and fifth lumbar vertebræ; *insertion,* one-half of lower border of last rib, apices of transverse processes of upper four lumbar vertebræ; *action,* flexes trunk—both acting; flexes laterally—one acting; depresses ribs with fixed pelvis and is then an expiratory muscle; *nerves,* anterior branches of lumbar nerves.

Muscles of Thorax.

External intercostals[11] (Fig. 58) (eleven pairs): *origin,* each from outer lip of groove on lower border of each rib from tubercle to costal cartilage; *insertion,* pass obliquely downward and forward to upper border of rib below for same distance; *action,* raise and evert ribs in inspiration; *nerves,* intercostals.

Internal intercostals (eleven pairs): *origin,* each from inner lip of groove on lower border of rib reaching from angle to sternum; *insertion,* passing downward and backward to upper border of rib below; *action,* depress and invert the ribs in expiration; *nerves,* intercostals.

Infracostales: vary in number, most common between lower ribs; *origin,* from inner surface of one rib; *insertion,* inner surface of first, second, or third rib below, passing obliquely downward and backward; *action,* inspiratory muscles; *nerves,* intercostals.

Triangularis sterni: origin, lower part of side of sternum, inner surface of ensiform cartilage, and sternal ends of costal cartilages of three or four lower true ribs; *insertion,* by fleshy digitations into lower border and inner surfaces of costal cartilages of second, third, fourth, and fifth ribs; *action,* draws down costal cartilages in expiration; *nerves,* intercostals.

Levatores costarum: origin (twelve pairs), extremities of dorsal transverse processes; *insertion,* rough surface between tubercle and angle of rib below; *action,* assist external intercostals in raising ribs; *nerves,* intercostals.

Diaphragm: *origin,* arched and convex toward chest, from ensiform cartilage in front, on either side, inner surfaces of cartilages and bony portions of lower six or seven ribs, behind, from the ligamentum arcuatum externum[7] and internum[6], and by its crura[4],[5], the left from bodies of second and third, the right, from those of second, third, and fourth lumbar vertebræ; *insertion,* into central cordiform tendon[1],[2],[3]; *action,* chief respiratory muscle, increasing capacity of chest by becoming flattened when contracted; also aids all efforts of expulsion as of feces, etc.; *nerves,* phrenic and phrenic plexus of sympathetic.

Fig. 59.



V Fig 505 (Gerrish)

Openings of the Diaphragm: the *aortic*[8], median, in front of vertebral bodies—really behind diaphragm—for aorta, vena azygos major, thoracic duct, sometimes left sympathetic nerve; the *œsophageal*[9], elliptical, muscular, formed by crura, for œsophagus and pneumogastric nerves; the *opening for vena cava*[10], highest, quadrilateral bounded by four tendinous bundles meeting at right angles; the right *crus*[5], transmits the sympathetic, greater and lesser splanchnic nerves; the *left crus*[4], the left greater and lesser splanchnic nerves, and vena azygos minor.

Muscles of shoulder and arm.

Pectoralis major[1] (Fig. 58): *origin*, anterior surface inner half of clavicle, half the breadth of anterior surface of sternum as low as sixth or seventh rib, cartilages of all true ribs except first or seventh, or both, and from aponeurosis of external oblique; *insertion*, fibres converge and overlap, the lower being folded back upon themselves, to be attached by a flat tendon to anterior lip of bicipital ridge of humerus; *action*, draws arm forward and inward; with fixed arm, an accessory muscle of forced inspiration; *nerve*, anterior thoracic.

Pectoralis minor[6] (Fig. 58): *origin*, tendinous from upper margin and outer surface of third, fourth, and fifth ribs near cartilages, and aponeurosis over intercostal muscles; *insertion*, anterior border coracoid process of scapula; *action*, depresses point of shoulder, elevates ribs during inspiration when scapula is fixed; *nerves*, anterior thoracic.

Subclavius[5]: *origin*, tendinous from first costal cartilage; *insertion*, groove on middle third of under surface of clavicle; *action*, draws clavicle downward and inward toward the thorax; *nerve*, branch from cord formed by fifth and sixth cervical.

Serratus magnus[4] (Fig. 58): *origin*, by nine fleshy digitations from outer surface and upper border of upper eight ribs (second rib has two) and aponeurosis covering upper intercostal spaces; *insertion*, whole length of anterior aspect of posterior border of scapula; *action*, elevates ribs in inspiration with fixed scapula, also raises point of shoulder by rotating bone on chest wall; *nerve*, posterior thoracic.

Deltoid[2] (Fig. 58): *origin*, outer third of anterior border and upper surface of clavicle, outer margin and upper surface of acromion process, and the whole length of lower border of spine of scapula; *insertion*, tendinous into rough prominence on middle of outer side of shaft of humerus; *action*, raises arm to right angle, also draws it forward by anterior fibres, or slightly backward by posterior portion; *nerve*, circumflex.

Subscapularis[4] (Fig. 60): *origin*, inner two-thirds of subscapular fossa; *insertion*, lesser tuberosity of humerus; *action*, rotates head of humerus inward; *nerve*, subscapular.

Supraspinatus[17] (Fig. 56): *origin*, internal two-thirds of same-named fossa and the fascia covering it; *insertion,* highest facet of greater tuberosity of humerus; *action,* assists deltoid, fixes head of humerus in socket; *nerve,* suprascapular.

Infraspinatus[18] (Fig. 56): *origin,* internal two-thirds of infraspinous fossa; *insertion,* tendinous, middle facet of greater tuberosity of humerus; *action,* rotates head of humerus outward; *nerve,* suprascapular.

Teres minor[19] (Fig. 56): *origin,* upper two-thirds of dorsal surface of axillary border of scapula and intermuscular septa; *insertion,* lowest facet on greater tuberosity of humerus; *action,* rotates head of humerus outward; *nerve,* circumflex.

Teres major[20] (Fig. 56): *origin,* dorsal aspect of inferior angle of scapula and intermuscular septum separating from teres minor and infraspinatus; *insertion,* internal lip of bicipital groove; *action,* draws humerus downward and backward when raised, also rotates it inward; *nerve,* lower subscapular.

FIG. 60.



Coraco-brachialis[6] (Fig. 60): *origin,* fleshy from apex of coracoid process of scapula; *insertion,* tendinous, into rough ridge at middle of inner side of shaft of humerus; *action,* draws humerus forward and inward, also elevates it; *nerve,* musculo-cutaneous, which perforates the muscle.

Biceps flexor cubiti[7] (Fig. 60): *origin, short head,* tendinous, in common with coraco-brachialis from tip of coracoid process of scapula, *long head,* upper margin of glenoid cavity by rounded tendon continuous with glenoid ligament; *insertion,* tendinous into back part of tuberosity of radius (a bursa is interposed in front), and by a broad aponeurosis opposite elbow, into fascia of forearm; *action,* flexes and supinates forearm, renders forearm fascia tense; *nerve,* musculo-cutaneous.

Brachialis anticus[9]: *origin,* lower half of outer and inner surfaces of shaft of humerus, embracing insertion of deltoid; *insertion,* tendinous into anterior surface of coronoid process of ulna; *action*, flexes forearm; *nerves*, musculo-cutaneous and musculo-spiral.

Triceps extensor cubiti[10]: *origin*, *scapular or long head*, from rough triangular depression below glenoid fossa, the *external head* from posterior surface of shaft between upper part of musculo-spiral groove and insertion of teres major, the *internal head* from posterior surface below musculo-spiral groove; *insertion*, tendinous, back, upper surface of olecranon process of ulna; *action*, extends forearm and arm; *nerve*, musculo-spiral.

Subanconeus: *origin*, humerus above olecranon fossa; *insertion*, posterior ligament of elbow; *action*, prevents loose capsule from being pinched during extension; *nerve*, musculo-spiral.

Muscles of the Forearm.

Anterior, or flexor and pronator group: All take origin from internal condyle, except pronator quadratus.

Superficial layer.

Pronator radii teres[4] (Fig. 61): *origin*, double, above, from internal condyle and common tendon, fascia of forearm and intermuscular septum, below, from inner side of coronoid process of ulna; *insertion*, tendinous, into rough ridge on middle of outer surface of shaft of radius; *action*, pronates forearm, aids flexion; *nerve*, median.

Flexor carpii radialis[5]: *origin*, internal condyle by common tendon, forearm fascia, and intermuscular septa; *insertion*, tendinous, into base of index metacarpal; *action*, flexes wrist, then forearm; *nerve*, median.

Palmaris longus[6]: *origin*, inner condyle by common tendon, fascia, and intermuscular septa; *insertion*, tendinous, into anterior annular ligament and palmar fascia; *action*, tightens palmar fascia; *nerve*, median.

Flexor carpi ulnaris[8]: *origin*, double, by common tendon from inner condyle and from inner margin of olecranon by aponeurosis, and from upper two-thirds of posterior border of ulna and intermuscular septum; *insertion*, tendinous, into pisiform bone, annular ligament and base of little finger metacarpal; *action*, flexes hand on forearm; then forearm on arm; *nerve*, ulnar.

Flexor sublimis digitorum[7] (***perforatus***): *origin*, by three heads, one from internal condyle by common tendon and from internal lateral ligament and intermuscular septa, a second from inner side of the coronoid process of ulna, the third from oblique line of radius; *insertion*, by four tendons, each of which splits for the passage of the deep flexor tendons, each half becoming attached to middle of lateral margins of second phalanges; *action*, flexes fingers, then hand on forearm, then forearm on arm; *nerve*, median.

FIG. 61.



Deep layer.

Flexor profundus digitorum (***perforans***): *origin*, upper two-thirds of antero-internal surface of shaft of ulna, inner side of coronoid process, by an aponeurosis from upper two-thirds of posterior border of ulna and ulnar half of interosseous membrane; *insertion*, into bases of last phalanges by four tendons, which pass between the two slips of flexor sublimis tendons; *action*, flexes fingers, then wrist on forearm; *nerves*, ulnar and anterior interosseous.

Flexor longus pollicis: *origin*, upper two-thirds of front of shaft of radius, adjacent interosseous membrane, sometimes from base of coronoid process; *insertion*, base of last phalanx of thumb; *action*, flexes last thumb phalanx; *nerve*, anterior interosseous.

Pronator quadratus: *origin*, oblique line on lower fourth of anterior surface of ulna, the bone below, the anterior border of ulna and aponeurosis over muscles; *insertion*, lower fourth of anterior surface and external border of radius; *action*, pronates hand; *nerve*, anterior interosseous.

Radial Region: all the extensors and supinators arise, at least partially, from the external condyle and ridge of the humerus.

Supinator longus[13]: *origin*, upper two-thirds of external condyloid ridge and external intermuscular septum; *insertion*, base of styloid process of radius; *action*, supinates forearm after it has been pronated, flexes forearm on arm, second in power only to biceps; *nerve*, musculo-spiral.

Extensor carpi radialis longior[5] (Fig. 62): *origin*, lower third of external condyloid ridge and external intermuscular septum; *insertion*, radial side of base of metacarpal bone of index finger; *action*, extends wrist; *nerve*, musculo-spiral.

FIG. 62.



Extensor carpi radialis brevior[6]: *origin*, external condyle by common tendon, external lateral ligament, the covering aponeurosis and the intermuscular septa; *insertion*, radial side of base of metacarpal of middle finger; *action*, extends wrist; *nerve*, posterior interosseous.

Extensor communis digitorum[8]: *origin*, external condyle by common tendon, deep fascia and intermuscular septa; *insertion*, second and third phalanges of each finger by three main tendons, one of which subdivides into two; *action*, extends fingers, then hand, finally forearm on arm; *nerve*, posterior interosseous.

Extensor minimi digiti[9]: *origin*, common tendon and intermuscular septa; *insertion*, with tendon of common extensor into second and third phalanges of little finger; *action*, extends little finger; *nerve*, posterior interosseous.

Extensor carpi ulnaris[10]: *origin*, by common tendon from external condyle, middle third of posterior border of ulna below aconeus and forearm fascia; *insertion*, ulnar side of base of fifth metacarpal; *action*, extends wrist; *nerve*, posterior interosseous.

Anconeus: *origin*, back of outer condyle of humerus; *insertion*, side of olecranon and upper fourth of posterior surface of shaft of ulna; *action*, extends forearm; *nerve*, musculo-spiral.

Supinator brevis[11]: *origin*, external condyle of humerus, external lateral ligament and orbicular ligament of radius, oblique line of ulna, triangular depression in front of it and aponeurosis covering muscle; *insertion*, back inner part of neck of radius, bicipital tuberosity, and oblique line; *action*, supinates forearm; *nerve*, posterior interosseous which pierces it.

Extensor ossis metacarpi pollicis[13]: *origin*, posterior surface of shaft of ulna below anconeus, interosseous ligament, and middle third of posterior surface of shaft of radius; *insertion*, base of thumb-metacarpal; *action*, extends thumb-metacarpal; *nerve*, posterior interosseous.

Extensor primi internodii pollicis[13]: *origin*, posterior surface of shaft of radius and interosseous membrane; *insertion*, base of first thumb-phalanx; *action*, extends phalanx on metacarpal; *nerve*, posterior interosseous.

Extensor secundi internodii pollicis[14]: *origin*, posterior surface of shaft of ulna and interosseous membrane below extensor ossis metacarpi; *insertion*, last phalanx of thumb; *action*, extends thumb; *nerve*, posterior interosseous.

Extensor indicis: *origin*, posterior surface of shaft of ulna below extensor secundi; *insertion*, with tendon of common extensor into second and third phalanges of index finger; *action*, extends index finger; *nerve*, posterior interosseous.

Fasciæ of the Hand.

The ***anterior annular ligament*** is stretched between pisiform bone and unciform process of unciform bone on one side, and the tuberosity of scaphoid and ridge of trapezium on the other; it is continuous with forearm and palmar fascia, and receives the insertion of the palmaris longus muscle and partially that of flexor carpi radialis, which pierces it; beneath it pass the median nerve and, in one synovial sheath, the tendons of the flexor sublimis and profundus digitorum and the flexor longus pollicis.

The ***post rior annular ligament***, continuous with forearm fascia, is attached on dorsum internally to ulna, cuneiform, and pisiform bones and palmar fascia, externally to ridges on back of and margin of radius. Six separate synovial compartments exist for the extensor tendons; from without inward, they are: (1) outer side of styloid process, for extensor ossis metacarpi and extensor primi internodii pollicis; (2) behind same process, for extensor carpi radialis longior and brevior; (3) next for extensor secundi internodii pollicis; (4) more internally still, for extensor communis digitorum and extensor indicis; (5) opposite interval between radius and ulna, for extensor minimi digiti; and (6) back of ulna, for extensor carpi ulnaris.

The *palmar fascia*[9] (Fig. 61), consisting of a central and two lateral portions, invests muscles of hand forming their common sheath; the central segment is triangular, attached to anterior annular ligament above, below dividing into four slips opposite metacarpal heads, each slip dividing again to enclose flexor tendons and be inserted on sides of first phalanges; strong transverse fibres connect the separate processes, and vertical septa, continuous on each side with interosseous aponeurosis, separate the middle from lateral palmar groups of muscles; beneath is the superficial palmar arch, median and ulnar nerves; the lateral portions of this fascia thinly invest the muscles of the thenar and hypothenar eminences.

Muscles of the Hand.

Abductor pollicis[11] (Fig. 61): *origin*, ridge of trapezium and annular ligament; *insertion*, tendinous into radial side of base of first thumb-phalanx; *action*, draws thumb from median line—*i. e.*, abducts it; *nerve*, median.

Opponens pollicis: *origin*, palmar surface of trapezium and annular ligament; *insertion*, whole length of radial side of thumb-metacarpal; *action*, flexes metacarpal; *nerve*, median.

Flexor brevis pollicis[12] (Fig. 61): *origin*, by two heads between which lies flexor longus pollicis tendon, the external from trapezium and outer two-thirds of annular ligament, the internal from trapezoid, os magnum, base of third metacarpal, and sheath of tendon of flexor carpi radialis; *insertion*, sides of base of first thumb-phalanx by two tendons each containing a sesamoid bone; *action*, as name implies; *nerves*, median and ulnar.

Adductor pollicis: *origin*, whole palmar length of metacarpal of middle finger; *insertion*, ulnar side of base of first thumb-phalanx and internal sesamoid bone; *action,* draws thumb toward median line; *nerve,* ulnar.

The preceding muscles form the *thenar eminence.*

Palmaris brevis[10]: *origin,* tendinous from annular ligament and palmar fascia; *insertion,* skin on inner border of palm; *action,* wrinkles skin on inner palmar side; *nerve,* ulnar.

Abductor minimi digiti: *origin,* pisiform bone and expansion of tendon of flexor carpi ulnaris; *insertion,* ulnar side of base of first little finger phalanx; *action,* draws little finger from median line; *nerve,* ulnar.

Flexor brevis minimi digiti: *origin,* tip of unciform process and annular ligament; *insertion,* base of first little finger phalanx with the abductor; *action,* flexes little finger; *nerve,* ulnar.

Opponens minimi digiti: *origin,* beneath preceding, from unciform process and annular ligament; *insertion,* whole length of little finger metacarpal; *action,* flexes fifth metacarpal; *nerve,* ulnar.

The four preceding muscles form the *hypothenar eminence.*

Lumbricales: *origin,* tendons of deep flexor; *insertion,* tendinous expansion of extensor communis tendon over dorsum of each finger; they are four in number; *action,* aid in extending second and third phalanges (Duchenne, Hutchinson); *nerves,* two outer by median, two inner by ulnar.

Dorsal interossei (four in number): *origin,* by two heads from adjacent sides of the metacarpal bones; *insertion,* into bases of first phalanges and aponeurosis of common extensor; *action,* extend second and third phalanges, abduct fingers from middle line passing through centre of middle finger; *nerve,* ulnar.

Palmar interossei (three in number): *origin,* from entire length of palmar surfaces of second, fourth, and fifth metacarpal bones; *insertion,* into bases of first phalanges of same fingers; *action,* adduct (draw) fingers toward middle finger; *nerve,* ulnar.

Muscles of Lower Extremity.

Psoas magnus[11] (Fig. 63): *origin,* sides of bodies, intervertebral substances and bases of transverse processes of last dorsal and all the lumbar vertebræ; *insertion,* tendinous, with iliacus, into lesser trochanter; *action*, flexes thigh on pelvis, or *vice versâ*, also rotates femur outward; *nerves*, anterior branches of lumbar nerves.

Psoas parvus (often absent): *origin*, sides of bodies and intervertebral substance between last dorsal and first lumbar vertebræ; *insertion*, tendinous into ilio-pectineal eminence and iliac fascia; *action*, tensor of iliac fascia; *nerves*, anterior branches of lumbar nerves.

Iliacus[10]: *origin*, iliac fossa and inner margin of iliac crest, ilio-lumbar ligament and base of sacrum, anterior superior and inferior iliac spines, the notch between them, and capsule of hip-joint; *insertion*, with tendon of psoas magnus into lesser trochanter; *action*, same as psoas magnus; *nerve*, anterior crural.

The ***fascia lata***, or ***deep fascia***, is attached above to Poupart's ligament and iliac crest, behind to margins of sacrum and coccyx, internally to pubic arch and linea ilio-pectinea, and below to condyles of femur, tuberosities of tibia, and head of fibula; two strong intermuscular septa pass from the inner surface to whole length of linea aspera; numerous smaller septa provide separate sheaths for each muscle; just below Poupart's ligament, at the upper inner aspect of thigh, is the large oval *saphenous opening*, formed by the margins of the pubic and iliac portions of the fascia lata; *Poupart's ligament* is an artificial production, being in reality the thickened lower portion of the external oblique aponeurosis where the fascia lata becomes continuous with it.

Tensor vaginæ femoris[4]: *origin*, anterior part of outer lip of iliac crest and anterior superior iliac spine; *insertion*, fascia lata one-fourth down thigh, whence fascia is continued down to tibial head as the *ilio-tibial band*; *action*, tensor of fascia lata; *nerve*, superior gluteal.

Sartorius[5]: *origin*, anterior infe ~~sup~~erior iliac spine and upper half of notch below it; *insertion*, aponeurotic, into upper inner surface of shaft of tibia; *action*, flexes leg upon thigh, thigh upon pelvis, at same time drawing limb inward, thus crossing one leg over the other; *nerve*, anterior crural.

Rectus femoris[6]: *origin*, by two tendons, the *straight* from anterior inferior iliac spine, the *reflected* from groove above acetabulum; *insertion*, patella in common with three next muscles; *action*, extends leg; *nerve*, anterior crural.

Vastus externus[7]: *origin,* tubercle of femur, great trochanter, rough line leading thence to linea aspera and whole outer lip of linea aspera; *insertion,* aponeurotic into outer border of patella; *action,* extends leg; *nerve,* anterior crural.

Vastus internus and **crureus**[8]: *origin,* line leading from inner side of neck of femur to linea aspera and its whole inner lip, internal intermuscular septum, internal, anterior, and external surfaces of the shaft of femur between the anterior intertrochanteric line and the lower fourth of the bone; *insertion,* aponeurotic, into inner side

FIG. 63.



FIG. 64.



of patella, blending with other portions of quadriceps tendon; *action*, extends leg; *nerve*, anterior crural. The preceding four muscles are called the quadriceps extensor[6],[7],[8], whose tendon contains the patella[9], and which is *inserted* into tubercle of tibia by ligamentum patellæ.

Subcrureus: *origin*, by two heads from lower part of shaft of femur; *insertion*, upper part of synovial pouch of knee; *action*, draws up synovial sac: *nerve*, anterior crural.

Gracilis[15]: *origin*, aponeurotic from inner margin of ramus of pubes and ischium; *insertion*, upper inner surface of shaft of tibia above and behind sartorius; *action*, flexes leg and adducts thigh; *nerve*, obturator.

Pectineus[12]: *origin*, linea ilio-pectinea, bone in front of it, and tendinous prolongation of Gimbernat's ligament; *insertion*, rough line passing from lesser trochanter to linea aspera; *action*, flexes, adducts, and rotates thigh outward; *nerves*, obturator, accessory obturator, anterior crural.

Adductor longus[13]: *origin*, tendinous from angle of pubes; *insertion*, middle third of linea aspera; *action*, adducts and flexes thigh; *nerve*, obturator.

Adductor brevis: *origin*, outer surface of descending ramus of pubes; *insertion*, upper part of linea aspera; *action*, adducts and flexes thigh; *nerve*, obturator.

Adductor magnus[14]: *origin*, descending ramus of pubes, ascending ramus, outer margin, and under surface of tuberosity of ischium; *insertion*, rough line running from great trochanter to linea aspera, whole length of linea aspera, and by a tendon into tubercle above inner condyle of femur, between these portions of muscle is an angular interval for passage of femoral vessels, the lower orifice of Hunter's canal; *action*, adducts and rotates thigh outward; *nerves*, obturator and great sciatic; the external portion of the muscle presents four apertures, the three upper for the perforating arteries, the lowest for termination of profunda artery.

Gluteus maximus[2] (Fig. 64): *origin*, superior curved line of ilium and portion of bone and crest just behind it, posterior surface of last piece of sacrum, side of coccyx, aponeurosis over multifidus spinæ muscle and the great sacro-sciatic ligament; *insertion*, fascia lata and rough line running from great trochanter to linea aspera between vastus externus and adductor magnus; *action*, maintains trunk erect upon thigh, extends, abducts, and rotates thigh outward; *nerves*, small sciatic, and branch from sacral plexus.

Gluteus medius[1] (Fig. 64): *origin*, outer surface of ilium between superior and middle curved lines, outer lip of crest between, and gluteal aponeurosis; *insertion*, tendinous into oblique line on great trochanter; *action*, supports trunk, the posterior fibres rotate thigh *outward*, the *anterior* fibres rotate thigh *inward*, it also abducts and draws thigh forward; *nerve*, superior gluteal.

Gluteus minimus[8] (Fig. 65): *origin*, outer surface of ilium between middle and inferior curved lines and margin of sciatic notch; *insertion*, anterior border of great trochanter; *action*, draws forward, abducts and rotates thigh inward, aids in maintaining trunk erect; *nerve*, superior gluteal.

Pyriformis[9]: *origin*, by three fleshy digitations from bone between first, second, third, and fourth anterior sacral foramina, likewise grooves leading from them, from margin of great sacro-sciatic foramen, and great sacro-sciatic ligament; *insertion*, passing out of pelvis by great sacro-sciatic foramen is attached to upper border of great trochanter; *action*, an external rotator of thigh, which, when flexed, is abducted by this muscle, also draws pelvis forward with fixed femur; *nerves*, branches of sacral plexus.

FIG. 65.



Obturator internus[11]: *origin* internal surface of obturator membrane and tendinous arch for passage of vessels, and inner side of obturator foramen; *insertion*, upper border of great trochanter after leaving pelvis by lesser sacro-sciatic foramen; four or five tendinous bands pass in the substance of the muscle (Bigelow) from origin to insertion; *action*, similar to pyriformis; *nerves*, branches of sacral plexus.

Gemellus superior[10]: *origin*, spine of ischium; *insertion*, with tendon of obturator internus into upper border of great trochanter; *action*, rotates thigh outward; *nerves*, branches of sacral plexus.

Gemellus inferior[12]: *origin*, tuberosity of ichium; *insertion*, great trochanter with obturator internus; *action*, an external rotator of thigh; *nerves*, branches of sacral plexus.

Obturator externus: *origin*, inner margin of obturator foramen externally, and inner two thirds of outer surface of obturator membrane; *insertion*, digital fossa of femur; *action*, an external rotator of thigh; *nerve*, obturator.

Quadratus femoris[13]: *origin*, outer border of tuberosity of ischium; *insertion*, linea quadrati on back of trochanter major; *action*, external rotator of thigh; *nerves*, branches of sacral plexus.

Biceps[16] (Fig. 65); [4],[5] (Fig. 64): *origin*, by two heads, the *long*[16] (Fig 65) from lower inner facet on tuberosity of ischium by tendon common to semitendinosus, the *short head*[5] (Fig. 64), from linea aspera between adductor magnus and vastus externus, and from intermuscular septum; *insertion*, outer side of head of fibula, the tendon splitting to embrace external lateral ligament of knee, one band extending down as far as outer tuberosity of tibia; this muscle forms the *outer hamstring*; *action*, flexes leg, after which it slightly rotates it outward; *nerve*, great sciatic.

Semitendinosus[6] (Fig. 64), [18] (Fig. 65): *origin*, by common tendon with preceding muscle from tuberosity of ischium and adjacent aponeurosis; *insertion*, upper inner surface of shaft of tibia, behind sartorius and below gracilis; *action*, flexes leg on thigh; *nerve*, great sciatic.

Semimembranosus[7] (Fig. 64): *origin*, tendinous from upper outer facet on tuberosity of ischium; *insertion*, groove on inner tuberosity of tibia; *action*, flexes leg on thigh, after which it assists popliteus in rotating leg inward; *nerve*, great sciatic; the tendons of sartorius[10] (Fig. 64), gracilis[8] (Fig. 64), semimembranosus[7] (Fig 64), and semitendinosus[6] (Fig. 66) (enumerated from before backward), form the *inner hamstring*.

Muscles of the Leg.

Tibialis anticus[3] (Fig. 66): *origin*, outer tuberosity and upper two-thirds of external surface of shaft of tibia, adjacent interosseous membrane, deep surface of fascia, and intermuscular septum; *insertion*, inner under surface of internal cuneiform bone, base of first metatarsal; *action*, flexes and adducts tarsus; *nerve*, anterior tibial.

Extensor proprius pollicis[5]: *origin*, middle two-fourths of anterior surface of fibula, and to same extent from interosseous membrane; *insertion*, base of last phalanx of great toe; *action*, extends great toe and, continuing action, flexes tarsus on leg; *nerve*, anterior tibial.

Extensor longus digitorum[4]: *origin*, outer tuberosity of tibia, upper three-fourths of anterior surface of fibula, interosseous membrane, deep surface of fascia, and intermuscular septa; *insertion*, divides into three tendons, and one subdivides making four tendons, which are attached to second and third phalanges of four lesser toes; *action*, extends toes, then flexes tarsus on leg; *nerve*, anterior tibial.

Peroneus tertius[6] (part of preceding muscle): *origin*, lower fourth of anterior surface of fibula, interosseous membrane, and intermuscular septum; *insertion*, upper inner part of base of little toe metatarsal; *action*, same as long extensor; *nerve*, anterior tibial.

Gastrocnemius[10]: *origin*, by two heads from upper back part of condyles of femur and supra-condyloid ridges; *insertion*, with tendons of soleus and plantaris longus forms tendo-Achillis, which is inserted into posterior tuberosity of os calcis; *action*, extends foot; *nerve*, internal popliteal.

Soleus[9],[9]: *origin*, tendinous, from back of head of fibula and upper third of postero-internal surface of shaft, from oblique line of tibia and from middle third of its inner border; *insertion*, tuberosity of os calcis by tendo-Achillis; *action*, extends foot; *nerve*, external popliteal.

Plantaris: *origin*, lower part of outer bifurcation of linea aspera and posterior ligament of knee-joint; *insertion*, posterior surface of os calcis with tendo-Achillis; *action*, extends foot; *nerve*, internal popliteal.

Popliteus[6] (Fig. 67): *origin,* depression on outer side of external condyle of femur and posterior ligament of knee-joint; *insertion,* inner two-thirds of triangular surface above oblique line of tibia posteriorly, and tendinous expansion covering muscle; *action,* flexes leg upon thigh, then rotates tibia inward; *nerve,* internal popliteal.

FIG. 66.



FIG. 67.



Flexor longus pollicis[9] (Fig. 67): *origin,* lower two-thirds of postero-internal surface of shaft of fibula, except its lowest inch, lower part of interosseous membrane, intermuscular septum, and fascia covering tibialis posticus; *insertion,* base of last phalanx of great toe; *action*, flexes great toe, then extends foot; *nerve*, posterior tibial.

Flexor longus digitorum[7] (Fig. 67) (***perforans***): *origin*, posterior surface of tibia below oblique line internal to tibialis posticus, except lower three inches, also intermuscular septum, tendon passes behind malleolus in groove with tibialis posticus, but in separate synovial sheath; *insertion*, bases of phalanges of lesser toes by four tendons, each passing through a fissure in the tendon of the flexor brevis; *action*, flexes phalanges, then extends foot; *nerve*, posterior tibial.

Tibialis posticus[8] (Fig. 67): *origin*, by two pointed processes, between which pass the anterior tibial vessels, from whole of posterior surface of interosseus membrane except lowest part, posterior surface of tibia, external to flexor longus, between oblique line above and middle of external border below, and from upper two-thirds of internal surface of fibula; *insertion*, tuberosity of scaphoid and internal cuneiform bones; *action*, extends tarsus on leg, adducts foot; *nerve*, posterior tibial.

Peroneus longus[7] (Fig. 66): *origin*, head and upper two-thirds of outer surface of shaft of fibula, deep surface of fascia, and intermuscular septa, passes behind outer malleolus in groove with peroneus brevis[8] (Fig. 64), then traverses groove of cuboid; *insertion*, outer side of base of great toe metatarsal and internal cuneiform; *action*, extends foot, then everts it; *nerve*, musculo-cutaneous.

Peroneus brevis[8] (Fig. 66): *origin*, lower two-thirds of outer surface of shaft of fibula, intermuscular septa; *insertion*, passes with preceding muscle behind external malleolus in a groove lined with a synovial sheath common to both tendons, to be attached to dorsum of base of fifth metatarsal; *action*, same as peroneus longus; *nerves*, musculo-cutaneous.

Fasciæ of Foot.

The ***anterior annular ligament*** consists of (1) an upper vertical portion binding down the extensor tendons, attached to lower ends of tibia and fibula, continuous above with leg fascia, and containing only one synovial sheath situated internally for tendon of anterior tibial, the extensor longus digitorum, peroneus tertius, and extensor proprius pollicis tendons, and anterior tibial vessels and nerve passing beneath it without a distinct sheath; and (2) a horizontal portion attached externally to upper surface of os calcis, and internally to inner malleolus and plantar fascia, containing three sheaths, that internal for anterior tibial tendon, the next for extensor proprius pollicis tendon, and that most external for extensor longus digitorum and peroneus tertius.

The ***internal annular ligament*** is a strong band extending from inner malleolus to inner margin of calcis, converting the bony grooves on its surface into canals for the flexor tendons and plantar vessels; it is continuous with deep fascia of leg, plantar fascia, and origin of abductor pollicis muscle. Its three fibro-osseous canals transmit, enumerated from within outward, the tendons of posterior tibial, flexor longus digitorum, posterior tibial vessels and nerve—through broad space beneath ligament—and tendon of flexor longus pollicis.

The ***external annular ligament*** stretches between the extremity of the outer malleolus to outer surface of calcis, forming a common sheath for peroneal tendons.

The ***plantar fascia,*** the densest of all fibrous membranes, consists of a central and lateral portions; the former, thick and narrow behind, arising from inner tubercle of os calcis, divides into a process for every toe, each again splitting opposite metatarso-phalangeal joints into two slips—between which pass the flexor tendons—to be attached to sides of metatarsal bones and transverse metatarsal ligament; two chief and several subordinate intermuscular septa are given off, separating the muscular layers; the outer and inner segments are thin, and of no special interest.

Muscles of the Foot.

Dorsal region.

Extensor brevis digitorum: *origin,* outer surface of os calcis, external calcaneo-astragaloid ligament, and annular ligament; *insertion,* by four tendons, the first into first phalanx of great toe, the other three into outer sides of long extensor tendons of second, third, and fourth toes; *action,* accessory to long flexor on four inner toes; *nerve,* anterior tibial.

Plantar region. First layer.

Abductor pollicis[3] (Fig. 68): *origin*, inner plantar tubercle of os calcis, internal annular ligament, plantar fascia, and intermuscular septum; *insertion*, with innermost tendon of flexor brevis pollicis into inner side of base of first great toe phalanx; *action*, abducts great toe; *nerve*, internal plantar.

FIG. 68.



Flexor brevis digitorum[5]: *origin*, tendinous from inner plantar tubercle of calcis, central part of plantar fascia and intermuscular septa; *insertion*, by four tendons for four lesser toes, perforated opposite middle of first phalanges by tendon of long flexor, beyond the perforation the tendon splitting to be inserted into sides of second phalanges; *action*, flexes lesser toes; *nerve*, internal plantar.

Abductor minimi digiti[4]: *origin*, outer plantar tubercle of os calcis, bone in front of both tubercles, plantar fascia, and intermuscular septum; *insertion*, with short flexor of little toe into outer side of base of first little toe phalanx; *action*, abducts little toe; *nerve*, external plantar.

Second layer.

Flexor accessorius : *origin*, by two heads, from concave surface of calcis and calcaneo-scaphoid ligament and os calcis in front of outer tubercle, and from plantar ligament; *insertion*, outer margin and upper and under surfaces of tendon of long flexor of toes; *action*, aids long flexor; *nerve*, external plantar.

Lumbricales[7]: *origin*, tendons of long flexor; *insertion*, expansion of long extensor and bases of first phalanges; *action*, accessory to flexors; *nerves*, the two inner lumbricales by internal, the two outer by external plantar nerve.

Third layer.

Flexor brevis pollicis : *origin*, inner border of cuboid and contiguous portion of external cuneiform, and prolongation of posterior tibial tendon ; *insertion,* inner and outer side of base of first great toe phalanx by two tendons, each containing a sesamoid bone, the inner tendon blending with that of the abductor pollicis, the outer with that of adductor pollicis; *action,* flexes great toe; *nerves,* internal plantar, sometimes external plantar.

Adductor pollicis: *origin,* tarsal extremities of second, third, and fourth metatarsal bones and sheath of tendon of peroneus longus; *insertion,* with outer portion of the short flexor into outer side of base of first phalanx of great toe; *action,* adducts great toe; *nerve,* external plantar.

Flexor brevis minimi digiti: *origin,* base of little toe metatarsal and sheath of peroneus longus tendon; *insertion,* outer side of base of first phalanx of little toe; *action,* flexes first, extends second phalanx; *nerve,* external plantar.

Transversus pedis: *origin,* under surface of head of fifth metatarsal bone and transverse metatarsal ligament; *insertion,* with adductor pollicis into outer side of first phalanges of great toe; *action,* adducts great toe; *nerve,* external plantor.

Fourth layer.

Dorsal interossei (four): *origin,* each by two heads from adjacent sides metatarsal bones; *insertion,* tendinous into bases of first phalanges of corresponding toes and aponeurosis of common extensor; *action,* abducts toes from imaginary line passing through centre of second metatarsal; *nerve,* external plantar.

Plantar interossei (three): *origin,* base and inner sides of shaft of third, fourth, and fifth metatarsal bones; *insertion,* inner sides of bases of phalanges of same toes; *action,* adduct toes toward a line passing through middle of second toe; *nerve,* external plantar.

Vascular System.

What is the pericardium?

A closed fibro-serous sac of conical form, its apex surrounding the great vessels for about two inches above their origin, its base downward, attached to the central tendon of diaphragm. The outer fibrous coat is continued as tubular prolongations, lost upon the external coats of the roots of all the great vessels except the inferior vena cava, and is finally traceable as continuous with the deep layer of the cervical fascia; the serous coat lines the sac, forming a *parietal layer*, is reflected over the heart and great vessels, forming a *visceral layer*; its function is the secretion of a thin fluid in sufficient amount to moisten the surfaces, thus lessening friction during the heart's movements.

Describe the heart.

It is a hollow, conical muscle, with four distinct cavities, situated obliquely between the lungs, the base upward, backward, and to the right, the apex[11] (Fig. 69) downward, forward, and to the left, corresponding to fifth intercostal space, one and a half inches below and three-fourths of an inch to the right of the left nipple.

Give the points upon the exterior of the thorax corresponding to space occupied by the heart.

The upper border corresponds to a line drawn across the sternum on a level with the lower borders of the second costal cartilages; its lower border to a line crossing gladiolus from right side of costo-xiphoid joint to apex point above described; the right border extending from median line of sternum three inches, its left, one and one-half inches; as the lungs partially cover the heart in front, the "area of heart's dulness" can be roughly indicated, according to Holden, by "a circle one inch in radius, the centre of which is midway between the left nipple and the end of the sternum."

Give the size and weight of the heart.

In adults it measures five inches long, three and one-half broad, and two and one-half thick; weighs in male ten to twelve ounces, in proportion to body 1 to 169; in females, eight to ten ounces, 1 to 149.

What are the four cavities of the heart called?

The right auricle[3] and right ventricle[4], the left auricle and left ventricle, separated by a longitudinal septum—indicated externally in front and behind by the *interventricular grooves*—dividing the heart into lateral halves, the right, or venous heart, the left, or arterial heart; a transverse *auriculo-ventricular groove* on the exterior corresponds to the division between auricles and ventricles.

Describe the right auricle and the chief points requiring study.

Larger than the left auricle, with a capacity of about ~~two fluid~~ ounces, its walls are about one line thick; the venous blood is poured in by the superior[1] and inferior venæ cavæ[2] and the coronory sinus; note the following points:

The *sinus*, the large quadrangular cavity between the two venæ cavæ; the *appendix auriculæ*, a conical pouch with dentated edge, projecting forward to the left over the root of the aorta[8]; the *openings* of the two venæ cavæ, also that of the coronary sinus[3''], its orifice guarded by a semicircular fold of the auricular lining called the *coronory valve*—sometimes at its junction with the great coronary vein a valve with two unequal segments is found; the *tubercle of Lower*[3], a small projection on the right wall directing blood from the superior cava toward auriculo-ventricular opening; the *foramina Thebesii*, the mouth of numerous minute veins returning blood from the heart muscle; the semilunar *Eustachian valve*[3'] between the anterior margin of the inferior vena cava and auriculo-ventricular opening—it is larger in the fœtus, serving to direct the blood of the inferior cava through the foramen ovale; the *fossa ovalis*[3'], an oval depression at the lower part of the septum auricularum, the site of the oval foramen in the fœtus; the *annulus ovalis*, the prominent margin of fossa ovalis; the *musculi pectinati*, small prominent muscular columns, running across the inner surface of the appendix and adjoining wall of the sinus; the oval *auriculo-ventricular orifice*, about one inch in diameter, communicating with the right ventricle, its margins formed by a fibrous ring covered by the lining membrane, and guarded by the tricuspid valve.

Describe the right ventricle.

It is triangular, with apex downward, not quite reaching that of the heart, forming with the right auricle the anterior, rather than the right side of the heart; its capacity has been variously estimated at from ~~two to six fluid ounces~~; it presents the following points for examination:

Above is the conical prolongation called the *conus arteriosus*, from which opens the orifice of the *pulmonary artery* placed to the left of the auriculo-ventricular orifice anteriorly, corresponding to junction of third left costal cartilage with sternum, and is surrounded by a fibrous ring; the pulmonary *semilunar valves* guard this orifice; the *tricuspid valve*[5],[5'],[5''] formed of three triangular segments—the largest on the left side—by a reduplication of the lining membrane with interposed fibrous tissue, which are prevented from being forced into the auricle by the,

Chordæ tendineæ[5''], fine tendinous cords stretching from (1) the columnæ carneæ to the attached margin, (2) to the centres of the leaflets, and (3) to their free margins.

The *columnæ carneæ*[4'], three varieties of muscular columns, (1) simple ridges, (2) bands attached by both extremities, and (3) three or four *musculi papillares*, muscular eminences from which arise the chordæ tendineæ.

The three *pulmonary semilunar valves*[4], ~~two anterior~~ and one posterior, formed by a reduplication of the lining membrane with interposed fibrous tissue, with their free margins strengthened by a bundle of fibrous tissue, whose fibres radiate from a fibro-cartilaginous nodule—the *corpus Arantii*—to all parts, except to two narrow lunated portions on either side of the nodule, which are forced in contact when the valves are closed; above and behind each valve is a dilatation, the *sinus of Valsalva.*

Describe the left auricle.

Resembles the right in having a principal cavity or sinus and an appendix, but its walls are thicker, measuring about one and one-half lines. The following points should be studied:

The *openings of the pulmonary veins*, usually four in number, two emptying into the right and two into the left side; frequently the latter terminate by a common opening.

The *oval auriculo-ventricular opening*, surrounded by a fibrous ring, smaller than the right; the *musculi pectinati* fewer and smaller than on the right side; a *depression* over site of the fossa ovalis of the right auricle.

Describe the left ventricle.

Larger and more conical than the right it forms but little of the left side, most of the posterior surface and all the apex[11] of the heart, its walls being three times as thick as those of the right ventricle; its inner surface presents for examination, The *circular aortic opening*, with the usual fibrous ring, in front and to the right of the auriculo-ventricular opening, between them intervenes one of the segments of the mitral valve; the aortic orifice is opposite the left half of the sternum, on a line with the lower border of the third costal cartilage, its mouth being guarded by three semilunar valves; the *mitral valve*, composed of two irregular segments—the ~~larger in front~~—similar to those of the tricuspid valve, but larger and thicker, the leaflets being provided with chordæ tendineæ, whose mode of attachment is identical with those of the right side. The *semilunar aortic valves* resemble those of the pulmonary artery, but are larger and stronger; *sinuses*, like the sinuses of Valsalva of the pulmonary artery, are found behind the aortic valves.

FIG. 69.



The *columnæ carneæ* are more numerous, but smaller than on the right side, while there are only two *musculi papillares.*

What is the endocardium?

The delicate lining membrane of the heart continuous with the intima of the great vessels, by its reduplications forming the various valves.

Describe the heart structure.

It consists of *striated anastomosing* muscular fibres taking origin from the fibrous rings surrounding the aortic, pulmonary, and auriculo-ventricular openings, disposed in the *auricles* in a *deep layer* composed of looped and annular fibres, and a *superficial* transverse layer; in the *ventricles* numerous layers have been described, the deepest of which run circularly, the more superficial spirally, curving round the apex to form the *whorl* or *vortex,* those fibres from in front curving around to enter posteriorly, and *vice versa*; the most superficial fibres, especially those behind, pass across the septum from one ventricle to the other.

The Arteries.

What are the arteries?

Cylindrical tubular vessels conveying blood ~~from both~~ ventricles *to* all parts of the body; the blood they carry is called *arterial*; the pulmonary artery arising from the right ventricle conveying blood to the lungs, and the four pulmonary veins returning it to the left auricle, constitute the *lesser or pulmonic circulation,* while the aorta arising from the left ventricle carrying blood to the body generally, and the veins returning it, finally emptying by the two cavæ into the right auricle, form the *greater or systemic circulation.* In their distribution the arteries freely communicate with one another, the large branches as well as the small forming what are called *anastomoses* or *inosculations,* permitting the establishment of the collateral circulation after obliteration of a main artery.

Describe the structure of the arteries.

They possess three coats, an (1) *internal* (serous) or *intima*; (2)

a *middle* (*media*), composed in small vessels almost purely of circular muscular fibres, in the larger chiefly of yellow elastic tissue; this prevents the arteries from collapsing when cut across; (3) an *external* (*adventitia*) composed of connective tissue. Every vessel, except the intracranial vessels, is included with its vein or veins in a fibro-areolar *sheath*.

FIG. 70.



How are the arteries nourished and their calibre regulated?

The larger vessels are supplied by minute *vasa vasorum,* bloodvessels distributed in a fine network to their external and middle coats; while intricate nerve networks (plexuses), chiefly derived from the sympathetic with branches of the spinal system, supply the same coats of the large vessels—the smaller are usually supplied only with single filaments.

What are capillaries?

The intermediate vessels between the arteries and veins, disposed in the form of a network, of an average diameter of one-three-thousandths of an inch, formed of a fine transparent layer of endothelial cells united by a cement substance.

Describe the aorta[1], [2], [3], (Fig. 70).

This, the main trunk of the systemic arteries, arises from the upper part of the left ventricle, ascends and then arches backward to the left over the root of the left lung to descend within the thorax on the left side of the vertebral column, entering the abdominal cavity by the aortic opening[a] through the diaphragm, where it terminates opposite the ~~fourth lumbar~~ vertebræ in the two common iliac arteries[23]; the aorta is divided for convenience of study into the arch[1], thoracic[2], and abdominal aorta[3], while the arch is described as consisting of an *ascending, transverse,* and *descending portion.* The branches of each subdivision are,

From the arch { Two coronary, Left common carotid[7],
Innominate[4], Left subclavian[8].

From the thoracic aorta { Pericardiac, Posterior mediastinal,
Bronchial[9], Intercostals[11], [12],
Œsophageal[10].

From the abdominal aorta, Two phrenic[13], Two spermatic[x20].
Cœliac axis[14] { Gastric[15], Inferior mesenteric[21],
Hepatic[17],
Splenic[16], Eight lumbar[22].
Superior mesenteric[18],
Two supra-renal[19], Sacra media[24],
Two renal[20].

4f.C H.

Describe the coronary arteries.

The right arises from the aorta above the free margin of the right semilunar valve, passes in groove between right auricle and ventricle to ~~posterior interventr~~icular groove, where, dividing into two branches, one ~~continues~~ onward anastomosing with the ~~left coro~~nary, the other descends along the interventricular groove to apex of heart anastomosing with descending branch of left coronary; the ~~left coronary~~, the smaller, arises above the free edge of left semilunar valve, passes forward between pulmonary artery and left appendix, thence obliquely to anterior interventricular groove, where it divides, one branch passing around to join the right vessel, the other descending in the groove to anastomose at apex with descending branch of right vessel.

Describe the innominate artery[4].

Arising from the commencement of transverse portion of arch, it is from one and a half to ~~two~~ inches long and bifurcates at upper border of right sterno-clavicular articulation into right common carotid[5] and subclavian[6], sometimes it sends off a *middle thyroid* branch—occasionally no innominate exists, the right carotid and subclavian springing directly from the aorta.

Describe the common carotid^{5},^{7}.

The right arises opposite right ~~sterno-clavicular~~ joint from innominate, the left from summit of aortic arch. In the neck their course corresponds to a line passing from the sterno-clavicular joint to a point midway between the mastoid process and angle of lower jaw; below, the trachea only separates them, above there is a wide interval; a common sheath of deep fascia encloses the internal ~~jugular~~ vein, the p~~neumog~~astric nerve (posterior to both), and a~~rte~~ry, enumerated from *without inward*, while upon the sheath lies the descending branch of the ninth ~~crani~~al nerve (descendens noni); the vessel is overlapped for most of its course by the anterior margin of the sterno-cleido-mastoid muscle, while about its middle it is crossed by the omo-hyoid muscle, the anterior jugular, and middle thyroid veins, while above the omo-hyoid muscle, the sterno-mastoid artery, and the superior thyroid vein cross it; at the lower part of the neck the right internal jugular vein diverges from the artery, but the left often crosses the lower part of corresponding artery; opposite the upper border of the thyroid cartilage, each vessel divides into the internal and external

FIG. 71.



Name the branches of the external carotid artery with their subdivisions.

Superior thyroid[4], arising below greater cornu of hyoid bone; its branches are

Muscular,
Hyoid,
Superior laryngeal,
Superficial descending (sterno-mastoid),
Crico-thyroid.

Lingual arises between thyroid and facial running beneath hyoglossus muscle to under surface of tongue; its branches are

Hyoid, Dorsalis Linguæ, Sublingual, Ranine.

Facial[6] arises just above lingual to cross lower jaw just anterior to masseter muscle; its branches are

Inferior or ascending palatine,
Tonsillar,
Submaxillary, CERVICAL
Submental[7],
Muscular[10],
Inferior labial[8],
Inferior coronary[8],
Superior coronary[9].
Lateralis nasi[11],
Angular[12],
FACIAL

Occipital[13] arises posteriorly opposite facial, lies in occipital groove of temporal bone; its branches are

Muscular,
Auricular,
Sterno-mastoid.
Meningeal,
Arteria princeps cervicis.
Cranial (distributed over occiput).

Posterior auricular[16] arises opposite styloid process, ascends beneath parotid gland to groove between cartilage of ear and mastoid process; its branches are

Stylo-mastoid, Auricular, Muscular, Parotid.

Ascending pharyngeal, running between carotid and side of pharynx; its branches being

External (muscular and nervous), Pharyngeal, Meningeal.

Temporal, the smaller terminal branch, commences in ~~substance of parotid gland~~, crosses root of zygoma, two inches above which it divides into anterior[24] and posterior[25] temporal; its branches are

Transverse facial[20] (Fig. 71), Middle temporal[23] (Fig. 71),
Anterior auricular. Orbital.

Internal maxillary[1], the larger terminal branch passes inward at right angles to the vessel at the inner side of neck of condyle of the lower jaw; it is divided into three portions, (1) maxillary, (2) pterygoid, and (3) spheno-maxillary.

(1)	Anterior tympanic,	Small meningeal,
	Middle meningeal,	Inferior dental { Incisor, Mental.
(2)	Deep temporal anterior and posterior,	Masseteric, } Mastication.
	Pterygoid,	Buccal. (Buccinator)
(3)	Alveolar,	Vidian, ([illegible]
	Infra-orbital,	Pterygo-palatine,
	Posterior or descending palatine.	Nasal or spheno-palatine. [illegible]

FIG. 72.



Describe the internal carotid artery.

It runs in ~~front of the transverse~~ processes of the ~~three~~ upper cervical vertebræ in contact with the pharynx and tonsil to the carotid canal of temporal bone, enters the skull, passes through the cavernous sinus and pierces the dura mater near the anterior clinoid processes when it divides into its terminal branches.

Give its branches.

Tympanic (deep), from artery in carotid canal, anastomosing with tympanic branch of internal maxillary, stylo-mastoid, and Vidian arteries.

Arteriæ receptaculi, small vessels supplying cavernous and inferior petrosal sinuses, pituitary body, Gasserian ganglion; one to the dura mater anastomosing with the middle meningeal is called the *anterior meningeal*,

Ophthalmic, from cavernous portion, enters orbit by optic foramen, dividing at the inner angle of the eye into two terminal divisions, the ~~*frontal* and *nasal*~~; its branches are

.X v. 9. p4

Lachrymal,
Supraorbital,
Posterior ethmoidal,
Anterior ethmoidal,
Palpebral,
~~Frontal~~,
~~Nasal~~,
Muscular,
Anterior ciliary,
Short ciliary,
Long ciliary,
Arteria centralis retinæ.

1. *Anterior cerebral* passes forward in the great longitudinal cerebral fissure, and communicates with its fellow by the anterior communicating artery, a vessel about two lines long.

2. *Middle cerebral*, the largest branch is lodged in the Sylvian fissure and divides into an anterior branch to pia mater; a middle branch to small lobe at outer extremity of fissure; and a posterior branch, supplying middle cerebral lobe.

3. *Anterior choroid*, to hippocampus major, corpus fimbriatum, velum interpositum, and choroid plexus.

4. *Posterior communicating* runs back to anastomose with posterior cerebral, a branch of basilar artery.

Describe the circle of Willis.

It consists of a vascular anastomosis at the base of the brain, between the branches of the carotids and basilar artery, ~~where~~by pressure on, or ~~blocking of any~~ two of the main trunks, will not prevent both sides of the brain from receiving a supply of blood. Its formation is as follows: The two vertebrals by their junction form the *basilar*, which divides into two *posterior cerebral*, these latter being connected with the back part of the internal carotids on each side by a *posterior communicating*, while the anterior terminals of the carotid, viz., the *anterior cerebral* on each side, is connected in front with its fellow by the short *anterior communicating artery* before mentioned.

Describe the subclavian.

On the right side it springs from the innominate, on the left directly from the aortic arch. It is divided into three portions, viz., that internal to the anterior scalene muscle, that behind the muscle, and that external to the scalene, this muscle intervening between the artery and subclavian vein; it ceases to be called subclavian at the lower border of the first rib, being termed axillary. All the branches arise from the first portion, except the superior intercostal on the right side, which comes from the second portion; its branches are,

The *vertebral*, which passes through the foramina in cervical transverse processes except that of seventh, enters skull through foramen magnum, joining its fellow at the lower border of the pons Varolii to form the *basilar artery*; its branches are,

[illegible] Lateral spinal,
Muscular,
Posterior meningeal,
Anterior spinal,
Posterior spinal,
Posterior inferior cerebellar.

The *basilar*, formed by the vertebrals, gives off the following branches ; [illegible]

Transverse, Anterior inferior cerebellar, Superior cerebellar, Posterior cerebral.

Thyroid axis divides almost at once into

Inferior thyroid, to same named gland, giving off

Laryngeal,
Tracheal,
Muscular,
Œsophageal,
Ascending cervical.

~~*Suprascapular*~~, chiefly to shoulder-joint and supra-spinous fossa; it anastomoses with acromial thoracic, posterior circumflex, the posterior and subscapular arteries.

Transversalis colli, larger than preceding, passes transversely outward to trapezius, beneath which it divides into the

Superficial cervical,
~~Posterior sca~~pular.

Internal mammary arises from under surface of first portion, ~~oppo~~site ~~thyroid~~ axis, descends upon costal cartilages a short distance from the sternum to the sixth interspace, where it divides into *musculo-phrenic* and *superior epigastric*, the latter anastomosing with the deep epigastric of external iliac; the branches are,

Superior phrenic (comes nervi phrenici),
Mediastinal,
Pericardiac,
Sternal.
Anterior intercostal,
Perforating,
Musculo-phrenic,
Superior epigastric.

Superior intercostal, its branches are,

Intercostals, [illegible] Posterior muscular, Spinal, Profunda cervicis, supplying posterior cervical muscles and anastomosing with arteria princeps cervicis from occipital.

Describe the axillary.

It extends from lower border of first rib to lower border of tendons of latissimus dorsi and teres major muscles, when it takes the name of brachial; its branches are,

Superior thoracic, supplying pectoral muscles and thoracic wall.

Acromial thoracic, supplying the deltoid by acromial branches, the serratus magnus, and pectorals by thoracic and descending branches.

Long thoracic, to chest muscles and mammary gland.

Alar thoracic, supplies axillary glands.

~~*Subscapular*~~, anastomosing with supra- and posterior scapular arteries, and giving off the do~~rsalis scapulæ~~. , infra sp., –

Posterior circumflex, to deltoid muscle and shoulder-joint; it anastomoses with the anterior circumflex, supra-scapular, acromial thoracic, and superior profunda arteries.

Anterior circumflex, to joint and deltoid; it anastomoses with posterior circumflex and acromial thoracic arteries.

Describe the brachial. < Radial / Ulnar.

A continuation of axillary, extending from lower border of teres major and latissimus dorsi tendons to its bifurcation into radial and ulnar, usually just below bend of elbow; the ~~median~~ nerve ~~crosses~~ it from without inward about midway in its course; its branches enumerated from above downward are,

5 br. *Superior profunda*, descending arm in musculo-spiral groove, giving off posterior articular artery, which anastomoses with the interosseous recurrent, posterior ulnar recurrent, and anastomotica magna, or inferior profunda, while the terminal twigs of the main trunk inosculate with the recurrent radial.

Nutrient artery, to humerus.

Inferior profunda, anastomosing with anterior and posterior ulnar recurrents and anastomotica magna.

Anastomotica magna, anastomosing with posterior articular, inferior profunda, anterior and posterior ulnar recurrents.

Muscular, to arm muscles.

Describe the radial artery. (V. [illegible]

It appears to be a ~~continuation of brach~~ial, but much smaller, extending from bifurcation at elbow along radial side of forearm to wrist, where, winding be~~neath the thum~~b, it passes ~~between~~ the two heads of the first ~~dorsal interosseous~~ muscle to ~~palm~~, there inosculating with the deep or communicating branch of the ulnar to form

The *deep palmar arch*, having as branches three to four palmar interossei; sometimes described as also giving off the radialis indicis, perforating, and recurrent branches, here credited to main vessel.

The branches of radial are,

Forearm,		Wrist,
Radial recurrent,	Arch →	*Posterior carpal,* → arch 2-3 D. Int
Muscular,		*Metacarpal,* [illegible]
Superficialis volæ,		*Dorsalis pollicis,*
Anterior carpal.		*Dorsalis indicis.* = [illegible]

Hand,
Princeps pollicis, (Int side thumb, 1st [illegible]
Radialis indicis, Dors. Indic.
Perforating, (Post. Communicating → D. I. [illegible]
Palm. *Interosseous.* (3)

Describe the ulnar.

Larger than the preceding, it passes from bifurcation obliquely inward to middle of forearm, thence runs along its ulnar border, across annular ligament to radial side of pisiform bone, curving across palm where, either with or without anastomosing with the superficialis volæ of the radial, it forms the

Superficial palmar arch, whose branches are the four digitals.

The branches of the ulnar are, in the

Forearm,
Anterior ulnar recurrent,
Posterior ulnar recurrent,
Interosseous { *Anterior interosseous,* nutrient, m.,
Interosseous { *Posterior interosseous,* recurrent [illegible]
Muscular.

Wrist,
Anterior carpal, arch
Posterior carpal. "

Hand,
Deep, or communicating branch, → palmar arch " "
Superf. [illegible] → *Digitals.* < Collateral dig. → D. br. anast.
Deep " "

Describe the thoracic aorta[2] (Fig. 73).

Fig. 73.



Commencing at left side of ~~low~~er border of fifth dorsal vertebra, it terminates at the ~~aortic o~~pening[a], in diaphragm in front of body of ~~last dorsal~~ vertebra; its branches are,

Pericardiac, variable in number and origin, for pericardium.

Bronchial[9], irregular in number and origin, usually one right and two left, for lung-tissue proper.

Œsophageal[10], commonly numbering four to five, anastomosing around the gullet with branches of inferior thyroid, phrenic, and gastric arteries.

Posterior mediastinal, numerous and small to glands and ~~areolar tissu~~e.

Intercostals[11], [12], usually ~~ten~~ on either side, each dividing into an anterior and posterior branch, the former subdividing into two at the angle of the ribs, one branch running in groove at lower border of rib above, the other running along the upper border of the rib below, in front both anastomosing with ~~anterior inter~~costal branches of the inter~~nal ma~~mmary, with thoracic branches of axillary, with epigastric, phrenic, and lu~~mbar arte~~ries; the posterior division supplies the vertebræ, spinal cord, dorsal muscles, and skin.

Describe the abdominal aorta[3].

It commences at the aortic opening[a], of the diaphragm in front of body of last dorsal vertebræ, whence descending a little to left of vertebral column it terminates on the body of the ~~fourth~~ lumbar vertebra by dividing into the common iliacs[23]; its branches are,

Cœliac axis[14], arises opposite margin of diaphragm, passes for half an inch forward to divide into the

7 *Gastric*[15], passing [illegible] ~~along~~ lesser ~~greater~~ curvature of stomach, anastomosing with aortic œsophageal, and branches of splenic and hepatic arteries.

Hepatic[17], dividing in transverse hepatic fissure into right and left branches for same lobes of liver; its branches are,

Pyloric, Stomach
Gastro-duodenalis, { Gastro-epiploica dextra, (Splenic
Pancreatico-duodenalis superior, (S [illegible]
Cystic,

whereby it supplies the parts indicated by names of vessels, and anastomoses with splenic, gastric, and superior mesenteric arteries.

Splenic[16], the largest branch, passes behind upper border of pancreas to spleen, giving off the

Pancreaticæ parvæ, Gastric (vasa brevis),
Pancreatica magna, Gastro-epiploica sinistra.

(One of the phrenics may arise from the cœliac axis.)

Phrenic[13], one on each side (sometimes one from cœliac axis instead of aorta) to under surface of diaphragm.

Superior mesenteric[18] comes off about ~~one-quarter inch~~ below cœliac axis, arching forward and downward to the left, supplying all of small intestine, except first part of duodenum, also cœcum, ascending, and transverse colon, giving off,

Inferior pancreatico-duodenal, Ileo-colic,
Vasa intestini tenuis, Colica-dextra,
Colica media.

Inferior mesenteric[21], arising from left side of aorta two inches above bifurcation, passes down into left iliac fossa and pelvis, supplying descending colon, sigmoid flexure, and greater part of rectum, anastomosing above with the middle colic of superior mesenteric; its branches are, Colica sinistra, Sigmoid, Superior hemorrhoidal.

Suprarenal, each arises opposite superior mesenteric to supply suprarenal bodies.

Renal[20], spring nearly at right angles from sides of aorta below superior mesenteric, the right longer than left; each divides into four or five branches before entering hilum of kidney, intervening between the renal vein in *front*, and the ureter *behind*.

Spermaticx[20] (ovarian in female), spring from front of aorta on each side, a little below renals, run behind peritoneum to pass in *male* through a~~bdominal~~ ring ~~to testes~~, in *female* between the laminæ of the broad ligaments to the ovaries, Fallopian tubes, uterus, and integument of labium and groin.

Lumbar[22], commonly four on each side corresponding to inter~~costals~~, and like them dividing into

Dorsal branches to vertebræ, spinal cord and back muscles, and abdominal branches, passing forward to anastomose with twigs from epigastric, internal mammary, intercostals, ilio-lumbar, and circumflex iliac.

Middle sacral[24], springs from bifurcation, descends along middle of sacrum and coccyx, sending branches to rectum, lateral sacral arteries, etc.

Describe the common iliac arteries.

They extend from the aortic bifurcation to left of umbilicus—corresponding to a line touching the highest point of iliac crests—to divide opposite the i~~ntervertebral disk~~ between the ~~last lumbar~~ vertebræ and the sacrum, into the *internal* and *external iliac*; the right vessel is somewhat the longer, both being about two inches long, and each at its bifurcation is crossed by the ureter. (naturally

Describe the internal iliac.

It measures about one and a half inches, arising at point of bifurcation of common iliac, to divide at upper margin of ~~great sacr~~o-sciatic foramen into the ~~anterior~~ and *posterior trunk*; its branches are

Middle inf [illegible] Anterior division.

1. (3 *Superior vesical*, part of fœtal hypogastric artery; it supplies ~~vas deferens~~ and ~~ureter~~, as well as ~~bladder~~. , 4. Middle [illegible]

Middle vesical, usually branch of former to bladder and vesiculæ seminales.

Inferior vesical (vaginal in female), arising in ~~common~~ with middle hemorrhoidal, and is distributed to base of bladder, prostate gland, and seminal vesicles.

Middle hemorrhoidal, supplies rectum.

Uterine (in female), anastomosing with ovarian.

4. *Obturator*, passes through obturator canal to thigh, there dividing into an internal and external branch, anastomosing with twigs of internal circumflex; inside the pelvis its branches are

Iliac, to same named bone and muscle anastomosing with iliolumbar; a *vesical*, to bladder; and a *pubic*, inosculating back of pubes with epigastric; in two out of every three cases the obturator springs from internal iliac, in one case in three and a half from epigastric.

5 *Internal pudic*, the smaller terminal of anterior division, is distributed to the external organs of generation; giving off the following branches

Inferior hemorrhoidal,
Superficial perineal,
Transverse perineal.
Artery of the bulb,
Artery of the corpus cavernosum,
Dorsal artery of the penis.

Sciatic, the larger terminal, supplies ~~muscles~~ on ~~back of~~ pelvis; its branches are,

Muscular (internal),
Hemorrhoidal,
Vesical,
Coccygeal,
Inferior gluteal,
Comes nervi ischiadici.
Muscular (external),
Articular (hip),

Posterior division.

1. *Ilio-lumbar*, dividing into an iliac and lumbar branch supplying muscular, spinal, and bone branches, anastomosing with gluteal, circumflex iliac, external circumflex, and epigastric arteries, etc.

2. *Lateral sacral*, superior and inferior on each side anastomosing with gluteal.

3. Sup *~~Gluteal~~*, the ~~termination of posterior~~ division, divides into a superficial and deep branch, giving off

Muscular, *cutaneous*, *nutrient* (to ilium), and *articular branches*, anastomosing with circumflex iliac and external circumflex arteries.

Describe the external iliac.

Passes along inner border of psoas muscle from bifurcation of common iliac to Poupart's ligament; a line drawn from left side of umbilicus to the midpoint between symphysis pubis and anterior superior iliac spine (in females a little nearer the former), indicates its course; its branches are

Muscular, *Lymphatic* (to glands).

~~Deep~~ epigastric, usually coming off just above Poupart's ligament, passing ~~between peritoneu~~m and transversalis fascia to pierce the lower third of sheath of the rectus abdominis muscle, continuing back of which it anastomoses with internal mammary and inferior intercostal arteries; its branches are

Cremasteric, *Pubic,* (→ in. side Fem. ring *Muscular.*

~~Deep~~ circumflex iliac, arises externally nearly opposite epigastric, running along i~~nner side of~~ iliac ~~cre~~st to its middle, there to pierce the transversalis muscle; it anastomoses with ilio-lumbar, gluteal, lumbar, and epigastric arteries.

Describe the femoral artery.

Extending from Poupart's ligament, where the letters N. A. V. indicate its relation with the anterior crural nerve and femoral vein—it terminates at the opening in the adductor magnus (commencement of Hunter's canal); the upper two thirds of a line drawn from the midpoint between the anterior superior iliac spine and symphysis pubis to inner side of internal condyle of the femur indicates its course. The artery and vein are enclosed in a strong fibrous sheath, but separated by a partition from each other, and lie very superficially above, in "Scarpa's triangle," bisecting it.

What is Scarpa's triangle?

It is a space bounded above by Poupart's ligament, internally by the adductor longus, and externally by the sartorius, the floor being formed chiefly by the iliac, psoas, pectineus, and long adductor muscles, the inner margin of the latter intervening between the artery and capsule of the hip-joint.

The femoral gives off the

Superficial epigastric[4], to inguinal glands, fascia, and skin, anastomosing with deep epigastric and internal mammary.

Superficial circumflex iliac[5], to skin of groin, glands, etc., anastomosing with ~~deep~~ circumflex iliac, g~~lutea~~l, and e~~xterna~~l circumflex.

FIG. 74.



FIG. 75.



Simple Simon Si
Dam Profunda Pi (3 Perf) } Femoral
Anus Circum Mag.

Superficial external pudic[6], to skin of genitals, anastomosing with ~~internal~~ pudic.

Deep external pudic[6], to skin of genitals and perineum, anastomosing with ~~superficial perineal~~.

Profunda femoris[7], springing from outer back part of femoral from one to two inches below Poupart's ligament, terminating at lower third of thigh by the small fourth perforating artery; it gives off

(*External*[8] and *Internal circumflex*[9]. *Three perforating*[10].)

Muscular[11], chiefly to sartorius and vastus internus.

Anastomotica magna[12], arises just above Hunter's canal, and divides into a superficial (cutaneous), and a deep branch; anastomosing with the superior[13] internal and external articular, and recurrent tibial.[14]

Describe the popliteal artery.

It commences at the opening in the adductor magnus, and passing obliquely downward and outward behind the knee-joint, divides opposite the lower border of the popliteus muscle into the *anterior* and *posterior tibial arteries*. Its branches are

Superior muscular,
Cutaneus,
Superior external articular,
Superior internal articular,
Azygos articular,
Inferior internal articular,
Inferior external articular,
Inferior muscular or sural.

These anastomose with the inferior perforating terminal branches of profunda, comes nervi ischiadici, anastomotica magna, recurrent tibial and with each other, except the muscular; the azygos articular enters back of joint to supply the synovial membrane and intra-articular ligaments.

Ext (like Radial).

Describe the anterior tibial artery.

Commencing at lower border of popliteus muscle, passing between the two heads of the posterior tibial muscle by the defect at the upper part of the interosseous membrane, lying upon its anterior surface and that of the lower third of the tibia, it terminates at the front of bend of the ankle in the *dorsalis pedis artery*; the anterior tibial nerve lies in close contact externally; it gives off the

Recurrent tibial[2], *Muscular*, *Internal*[5] *and External*[4] *malleolar.*

" Ant. Post. Carpal.

Describe the dorsalis pedis artery[3].

It extends from front of bend of ankle to back part of ~~first interosseous~~ space, where it divides into the

Dorsalis hallucis[7], supplying the great toe and adjoining side of second toe, and the (Rad. = Princeps indicis

Communicating[7], dipping down between heads of first dorsal interosseous muscle to reach the sole of foot, and form with external plantar the plantar arch; the other branches are the (= Rad.)

Tarsal, arching outward across tarsus. (= Carpal)

Metatarsal[6], runs anterior to preceding over bases of metacarpal bones, giving off (= Deep arch

Three interossei, from which arise *seven digitals*. (= ditto)

Describe the posterior tibial artery. (~~of Ulnar (Dist.~~)

It extends as a large vessel obliquely downward from lower border of popliteus muscle along tibial side of leg to the midpoint of the depression between the inner ankle and heel, where, ~~beneath the adductor~~ pollicis muscle it divides into the internal and external plantar; the posterior tibial nerve lies first to its inner side, but soon crosses it, to remain close to the outer side; it gives off the

Peroneal, along back of fibular side of leg, giving off (~~of Ant.~~ Inteross.

Anterior peroneal, piercing interosseous ligament two inches above outer malleolus, anastomosing with external malleolar and tarsal arteries; it also gives muscular branches and ~~nutrient~~ artery to fibula: the *posterior tibial* also gives off the EXT. CALC; POST. [illegible]

Nutrient (of tibia), largest to any bone.

Muscular, to posterior leg muscles.

Communicating, to peroneal, passing transversely across back of tibia about two inches above ankle.

Internal calcanean, several large branches to inner muscles of sole, fat and skin ~~of~~ heel, anastomosing with peroneal and internal malleolar.

Describe the internal plantar.

The smaller terminal of posterior tibial, it runs along inner side of foot and great toe.

Describe the external plantar artery.

It passes obliquely outward and forward to base of fifth metatarsal, whence it curves inward to interval between bases of first and second metatarsal bones, there anastomosing with the communicating branch of the dorsalis pedis, completing the *plantar arch*, whose branches, in addition to numerous *muscular* ones, are,

Posterior perforating, passing up the three outer interosseous spaces to anastomose with the interossei from the metatarsal.

Four digitals, by division supplying both sides of the three outer toes, and the outer side of the second—~~both~~ sides of the ~~great~~ and ~~inner~~ side of the ~~second~~ toe being supplied by the ~~communi~~cating branch of ~~dorsalis~~ pedis. Supplies A. & P. 4th toe, inn. 2d toe.
like Radial.

Describe the pulmonary artery.

It is a wide vessel conveying venous blood from the right ventricle to the lungs, about ~~two inches~~ long, and springs, in ~~front of~~ the ~~aorta~~, from left side of the base of the right ventricle, lying for most of its course within the pericardium, passing obliquely to the left, upward and backward, dividing at the ~~under~~ surface of the ~~aortic~~ arch into a ~~right and~~ *left pulmonary artery*, the former the longer; each vessel passes horizontally outward to its respective lung, where it divides into two main branches, one of those of the right subdividing to supply the third lobe; these vessels subdivide to terminate in the pulmonary capillaries.

Comm. [illegible]

Describe the collateral circulation after ligature of the common carotid.

The chief communications are between superior and inferior thyroids, the profunda cervicis and princeps cervicis of the occipital, the vertebral taking the place of the internal carotid within the cranium. (The student must remember that while all the anastomosing vessels coming off above and below the site of ligature enlarge, and should, therefore, be carefully impressed upon the mind where enumerated in the preceding pages, yet the blood, by dissection, has been found to pass chiefly by the vessels mentioned under each caption.)

After ligature of the subclavian.

Between the supra-scapular and posterior scapular with subscapular, the internal mammary and the long and short thoracic and subscapular.

After ligature of axillary.

If below acromial thoracic, chiefly between the subscapular and other scapular arteries of subclavian, and long thoracic, through intercostals with internal mammary; if below subscapular, the posterior circumflex anastomosing with supra-scapular and acromial thoracic, and inosculations between the subscapular and superior profunda.

After ligature of brachial.

(1) Upper third, by anastomoses between circumflex and subscapular and superior profunda; (2) below the profunda arteries, by branches of the profundæ anastomosing with recurrent ulnar, radial, and interosseous.

When thoracic aorta is obliterated.

The internal mammary anastomosing with intercostals; phrenic, by musculo-phrenic and comes nervi phrenici and deep epigastric, superior intercostal and first aortic intercostal; inferior thyroid with first aortic intercostal; transversalis colli with posterior branches of intercostals; axillary and subclavian branches to side of chest with lateral branches of intercostals.

When abdominal aorta is tied.

The deep epigastric communicating with the internal mammary, the intercommunications of superior and inferior mesenteric, or the latter with the internal pudic, and the anastomoses of the lumbar with branches of internal iliac.

After common iliac is tied.

The anastomosis of hemorrhoidal branches of internal iliac with superior hemorrhoidal from inferior mesenteric, the inosculations of the uterine and ovarian, and of vesicals of opposite sides, that of lateral sacral with sacra media, of epigastric with internal mammary, intercostal and lumbar, of ilio-lumbar with last lumbar, of obturators with one another and deep epigastric, of gluteal with posterior sacral branches.

After internal iliac is tied.

Same as above, except obturator also communicates with internal circumflex, the circumflex and perforating branches of femoral with sciatic, and the circumflex iliac with ilio-lumbar and gluteal.

After external iliac is tied.

The anastomoses between the ilio-lumbar and circumflex iliac; the gluteal and external circumflex; the obturator and internal circumflex; the sciatic with superior perforating and circumflexes; the internal pudic with external pudic and internal circumflex.

After common femoral is tied.

Anastomoses of gluteal and circumflex iliac with external circumflex; obturator and sciatic with internal circumflex; ilio-lumbar with external circumflex; comes nervi ischiadici with surals.

After superficial femoral is tied.

Branches from profunda anastomosing with superior and inferior articulars of knee, anastomotica magna, surals, and origins of anterior and posterior tibial.

(The popliteal is never tied except for wound, when the articular arteries, anastomotica magna, recurrent tibial, perforating branches, and surals would convey the blood.)

The Veins. V. Dict.

Describe the veins.

They are vessels returning venous blood—*i. e.*, blood surcharged with carbonic acid, to the heart, and have the same coats as arteries, but not so thick, especially the middle, in consequence collapsing when divided. One set of veins only carries other than venous blood, viz., the pulmonary, conveying arterial blood from the lungs to the left auricle. Certain of the veins, usually the superficial, have their lining membrane forming semilunar reduplications or valves, arranged in pairs opposite one another, which prevent any reflux of blood. The large arteries are accompanied by deep veins of the same name, generally included in the same sheath, while such vessels as the brachial, radial, ulnar, etc., have two veins each, called *venæ comites.* The superficial veins run between the layers of the superficial fascia, terminating in the deep veins, and are not usually accompanied by arteries. All veins freely anastomose, and even those dignified with special names have very irregular origins and distributions.

What are sinuses?

Venous channels, found only within the skull, formed by a separation of the layers of the dura mater, lined with endothelium continuous with that of the veins.

How are the veins classed?

As belonging to the *pulmonary*, *systemic*, or *portal systems*, the latter an appendage of the systemic.

Name the principal veins which have no valves.

The venæ cavæ, hepatic, portal, renal, uterine, ovarian, cerebral, spinal, and pulmonary.

Name the veins of the head and neck.

(1) Those of the exterior:
Facial, Internal maxillary, Posterior auricular,
Temporal, Temporo-maxillary, Occipital.

(2) Those which return the blood from head and face:
External jugular, Anterior jugular,
Posterior external jugular, Internal jugular,
Vertebral.

(3) Veins of diploë and cranium:
Veins of diploë, Cerebral,
Cerebellar, Ventricular, or venæ galenæ.

Name the cerebral sinuses.

Superior longitudinal,
Inferior longitudinal,
Occipital (2),
Circular,
Superior petrosal,
Straight sinus,
Lateral sinuses (2),
Cavernous (2),
Transverse,
Inferior petrosal.

How is the internal jugular[7] (Fig. 76) formed?

By the junction of the lateral and inferior petrosal sinuses just outside jugular foramen; at base of neck the internal jugular unites with subclavian vein to form the innominate vein—at, or above the junction is a pair of valves; into the jugular empty the facial, lingual, pharyngeal, superior and middle thyroid, and sometimes occipital veins.

Where do the following veins empty?

The *external jugular*[8]; into subclavian vein external to internal jugular; the *posterior external jugular,* into internal jugular; the *anterior jugular,* into termination of external jugular or the subclavian vein; the *vertebral,* descends through vertebral foramina in transverse processes of cervical vertebræ to empty into back of innominate vein, valves guarding the orifice.

Describe the veins of the upper extremity.

They consist of superficial and deep, the latter being the ***venæ comites*** of the arteries; commencing as ***digitals*** these empty into the *interossei,* these into ***palmar*** until the ***deep radial*** and ***ulnar*** are formed, which, uniting, form the so-called ***brachial vein,*** really consisting of two ***venæ comites*** with transverse anastomoses. The superficial veins running in the superficial fascia are called

Radial		*Cephalic*		
	Median cephalic			
Median				*Axillary.*
	Median basilic		*Brachial*	
Anterior and *Posterior ulnar*		*Basilic*		

Describe the axillary vein.

It is a continuation of brachial and basilic, receiving veins of corresponding names with the arteries of the axilla, and is continued beneath the clavicle under the name of the ***subclavian vein***[9]; it has valves opposite lower border of subscapular muscle, also at mouths of cephalic and subscapular veins.

Describe the subclavian vein[9].

It extends from lower border of first rib to back of sternoclavicular joint, there uniting with internal jugular to form the vena innominata—the subclavian vein lies in *front* of the artery with the anterior scalene muscle interposed at its second part; its branches are, the external and anterior jugular and branch of cephalic; on right side the *right lymphatic duct* empties at the junction of the axillary and internal jugular veins, and at the same point on the left side the *thoracic duct.*

How are the venæ innominatæ[5], [6] formed?

Each by the subclavian and internal jugular, which unite just below the first costal cartilage to form the superior vena cava[4]; the *right innominate* is about one and a half inches long, receiving blood by right vertebral, internal mammary, inferior thyroid, and superior intercostal veins; the *left vein* is larger and about three inches long; the left vertebral, internal mammary, inferior thyroid, superior intercostal, and occasionally some thymic and pericardiac veins empty into it.

Fig. 76.



Describe the superior vena cava[4].

It measures from two and a half to three inches and is formed by the union of the two innominate veins[5], [6], is half covered by the pericardium and enters upper part of right auricle; it returns the blood of the upper half of body and receives the vena azygos major and small mediastinal and pericardiac veins.

Describe the azygos veins.

They connect the superior and inferior venæ cavæ, taking the place of those vessels in that part of the chest occupied by the heart.

The *right azygos*[10] commences opposite first and second lumbar vertebræ by a branch from right lumbar or renal vein, or from inferior vena cava, enters thorax by aortic opening, arches over root of right lung to empty into superior cava, receiving in its course the nine or ten right lower intercostal veins, the vena azygos minor, œsophageal, mediastinal, vertebral, and right bronchial veins, also at times it is connected with right superior intercostal vein.

The *left lower azygos*[11] (vena azygos minor) commences by a branch from left lumbar or renal vein, enters thorax through left crus of diaphragm, passes across from the left side of vertebral column at sixth or seventh dorsal vertebra to enter the right azygos vein, receiving veins from four or five lower intercostal spaces; also mediastinal and œsophageal branches.

The *left upper azygos*[12] is formed by veins, usually two to three, from the intercostal spaces between left superior intercostal and highest branch of left lower azygos, and empties into right azygos or left lower azygos; it is sometimes absent, its place being taken by left superior intercostal.

Name the other principal veins of base of the neck and of the thorax.

Internal mammary,	*Mediastinal,*
Inferior thyroid,	*Pericardiac,*
Intercostals,	*Bronchial.*

Briefly describe the spinal veins.

They are the *dorsi-spinal* on the exterior of the spinal column, forming plexuses around vertebral spines, laminæ, and processes, emptying into intercostal, lumbar, and sacral veins respectively.

The *meningo-rachidian,* lying between vertebræ and theca spinalis forming plexuses, one running along the posterior surfaces of vertebral bodies forming the *anterior longitudinal veins* receiving the venæ basis vertebrarum, the other on the inner surface of the laminæ, the *posterior longitudinal veins,* both extending the whole length of the spinal canal; the posterior emptying into dorsi-spinal, the anterior into vertebral, intercostal, and sacral veins, respectively.

The *venæ basis vertebrarum,* in vertebral bodies, empty into anterior longitudinal.

The *medulli-spinal,* those of the cord itself, forming a minute plexus over cord between the pia mater and arachnoid, near base of skull converging to form two or three trunks terminating in the inferior cerebellar veins or petrosal sinuses.

Describe the chief veins of the lower extremity.

They are deep and superficial, the former commence as venæ comites of *digitals*, which form the *interossei*, these the *anterior* and *posterior tibial* and *peroneal* accompanying same named arteries, which uniting, form the *popliteal*, in the thigh to be called the *femoral*, being joined by *profunda femoris* and *internal saphenous veins*, which again changes its name to *external iliac* above Poupart's ligament; into external iliac empty the epigastric and circumflex iliac veins.

The superficial veins are the

Internal or long saphenous, commencing on inner side of dorsum of foot, running up on inside of leg and thigh to enter femoral vein after passing through the saphenous opening of fascia lata; its branches are

Cutaneous, *Superficial circumflex iliac*,
Superficial epigastric, *Communicating*,
Pudic.

External or short saphenous, commences at outer side of dorsum of foot, passes behind the external malleolus, up the middle of leg posteriorly to empty into popliteal vein between heads of gastrocnemius.

Describe the internal iliac vein.

It is formed by the venæ comites of all branches of external iliac artery except the umbilical, and unites opposite sacro-iliac articulation with external iliac vein to form the common iliac vein; it receives the following veins,

Gluteal, *Internal pudic*, *Hemorrhoidal*, *Uterine and vaginal*,
Sciatic, *Obturator*, *Vesico-prostatic* *Plexuses*, in female,
Plexuses, in male,
Dorsal vein of penis.

Describe the common iliac veins.

Formed by the junction of the external and internal iliac veins, and receiving the ilio-lumbar, occasionally the lateral sacral, and the *left iliac* the middle sacral vein, they unite at an angle upon the intervertebral substance between the fourth and fifth lumbar vertebræ to form the inferior vena cava.

Describe the inferior vena cava[1].

It runs upward from junction of the two common iliacs, along the right side of the aorta, pierces the central tendon of diaphragm, and terminates at back of right auricle, being partially covered by serous layer of the pericardium; it returns the blood from all parts below the diaphragm, and receives the following branches,

Lumbar,	*Suprarenal,*	sometimes *Middle sacral,*
Right spermatic,	*Phrenic,*	
Renal[2], [3],	*Hepatic.*	

Describe the portal system of veins.

Formed by the union behind the head of the pancreas of the *superior and inferior mesenteric, splenic, and gastric veins,* collecting the blood from the viscera of digestion, the resulting *portal vein* divides in the transverse fissure of the liver into a branch each for the right and left lobe, which ramify to form a *venous plexus* in the liver tissue; the hepatic artery sends branches within the liver to the portal vein, and external to the organ the vein receives the smaller gastric and the cystic vein; the portal blood is returned to the inferior vena cava by the *hepatic veins.*

What veins return the blood from the substance of the heart?

Great cardiac vein,	*Anterior cardiac veins,*
Middle cardiac vein,	*Right or small cardiac vein,*
Posterior cardiac veins,	*Venæ Thebesii.*

What is the coronary sinus?

A dilatation of about one inch of the great cardiac vein in posterior part of left auriculo-ventricular groove, covered by the muscular tissue of the left auricle and receiving the posterior cardiac veins, and an oblique vein from back of left auricle; its orifice is guarded by the coronary valve.

Describe the pulmonary veins.

Commencing in the lung capillaries they form a main vein for each lobule, which unite into two trunks for each lung, opening separately into the left auricle; at times there are three veins on the right side, or the two left terminate by a common opening.

The Lymphatics.

What are lymphatics?

Delicate vessels with transparent walls formed of same three coats as arteries and found in all parts of the body probably, except the nails, cuticle, hair, and cartilage; they have numerous valves producing their characteristic beaded appearance; they are supplied with nutrient arteries but not with nerves.

What are the lacteals?

The lymphatics of the small intestine, conveying *chyle during digestion*, lymph at other times.

Describe the lymphatic glands.

They are small solid, round, or ovoid glandular bodies, situated in the course of the absorbent vessels which previous to entering a gland break up into several *afferent vessels*, form a plexus within, and emerge by several *efferent vessels* which soon unite to form a single trunk; each gland is surrounded by a fibrous capsule which sends partitions inward, forming alveoli in which lies the gland-pulp or *lymphoid tissue* consisting of a rete whose meshes are filled with lymph-cells. The glands are chiefly found in the mesentery, along great vessels, in the mediastinum, axilla, neck, at front of elbow, groin, and popliteal space, being usually named from their locality, as axillary, etc.

Describe the thoracic duct.

This conveys the bulk of the lymph and chyle into the blood, being the common lymph-trunk, except for right upper extremity, right side of head, neck, and thorax, right lung, same side of heart, and convexity of liver. It commences by the triangular *receptaculum chyli*, on the front of body of second lumbar vertebra, enters = Beg. the thorax by aortic opening, and opposite the upper border of [illegible] seventh cervical vertebra it curves downward to empty at the junction of the left internal jugular and subclavian veins.

Describe the right lymphatic duct.

It is about one inch long, receiving lymph from those parts *excepted* in the account of the thoracic duct, and empties at the junc-

9 152-4.

tion of the right internal jugular and subclavian veins; both ducts have double semilunar valves at their orifices, preventing regurgitation of blood.

Nervous System.

What are the two divisions of the nervous system?

The *cerebro-spinal* or that presiding over animal life, and the *sympathetic*, that regulating organic life.

Describe the structure of the nervous tissue.

It is composed chiefly of two structures, the *gray or vesicular* originating impulses and receiving impressions; and *white or fibrous*, conducting impressions; in the sympathetic system is found a third structure, *gelatinous nerve-tissue*; seventy-five per cent. of nerve-tissues is composed of water, the remainder being albumen, phosphorized-fat, and salts.

Describe the microscopic structure of the white nerve-tissue.

It is formed of tubular fibres each consisting of a central *axis-cylinder*, surrounded by the *white substance of Schwann*, the whole enclosed by the *tubular membrane* or *primitive sheath*. A bundle of these fibres invested by a fibro-areolar membrane, the *perineurium* (neurilemma), constitutes a *nerve*, receiving a special blood-supply by the *vasa nervorum*; the *gelatinous* variety consists of finely granular fibrillæ enclosed in a sheath—by some these are not considered to be nerves.

Describe the gray or vesicular nerve-tissue.

This consists of large granular cells containing nuclei and nucleoli, ovoid, or with one or many processes (*unipolar, multipolar*), some of which become continuous with an axis-cylinder.

How do nerves terminate?

Peripherally sensory nerves end in minute plexuses, end-bulbs, tactile corpuscles, and Pacinian corpuscles; in the special organs they end in cells and in other not well ascertained ways; motor nerves end peripherally in plexuses or by "*motorial end plates.*" The central terminations are not well understood.

What organs compose the cerebro-spinal system?

The brain, spinal ganglia, and the cranial and spinal nerves.

Name the membranes of the brain.

The dura mater, the arachnoid, and pia mater.

Describe the cerebral dura mater with its processes.

It is a dense fibrous membrane lining the interior of skull, constituting the internal periosteum, is continuous with that of spinal cord, and is prolonged to outer surface of the skull through the various foramina; by separation of its layers the cerebral sinuses are formed; its smooth under surface is covered with endothelial cells; its processes are the,

Falx cerebri, a sickle-shaped layer occupying the longitudinal fissure of the brain; along the upper and lower border respectively run the superior and inferior longitudinal sinuses.

Tentorium cerebelli, covering the upper surface of the cerebellum; it supports the weight of the posterior lobes of the cerebrum; it is attached to the horizontal arms of the occipital cross, enclosing the lateral sinuses, to the upper margin of the petrous bone, including the superior petrosal sinuses, whence it extends to anterior and posterior clinoid processes; to the mid-line, above, the falx cerebri is attached antero-posteriorly, and below medianly the base of the

Falx cerebelli, a small triangular process passing between cerebellar lobes behind.

What are the Pacchionian bodies?

Numerous aggregations of small whitish granulations of unknown function formed upon outer surface of dura mater near superior longitudinal sinus, lying in depressions in the bone, in the superior longitudinal sinus, on the inner surface of the dura mater, and on the pia mater; unknown in infancy, rare before third year; they are usually found after tenth year.

Describe the arachnoid.

It is a delicate membrane lying between the pia mater and dura mater, being separated from the latter by the *subdural space*; it bridges over the convolutions, forming *part of subarachnoidean space*, and at the base, by being stretched between the middle lobes, the *anterior subarachnoidean space* is left; while between the cerebellar hemispheres and medulla oblongata lies the *posterior subarachnoidean space*, the two communicating across the crura cerebelli, and by an opening in its lower boundary with the fourth ventricle; these spaces ~~contain the cerebro-spinal fluid~~, forming an elastic water-cushion for the encephalon: it consists of interwoven bundles of fibrous and yellow elastic tissue covered with a layer of endothelium.

What is the pia mater?

It consists of a minute ~~plexus of bloodvessels~~ derived from the internal carotid and vertebral arteries, held together by fine areolar tissue; it dips between the convolutions, forms the velum interpositum and choroid plexuses of the fourth ventricle, and contains nerves and lymphatics.

What are the divisions of the brain called?

The cerebrum, cerebellum, pons Varolii, and medulla oblongata.

What is the weight of the brain?

~~Forty~~-nine and a half ounces, on the average, in males, and ~~forty-four~~ ounces in females; heaviest male brain recorded sixty-eight and three-eighths ounces, lightest thirty-four ounces; female brain, heaviest fifty-six ounces, lightest thirty-one ounces; idiots' brains seldom weigh more than twenty-three ounces.

Name the fissures and lobes of the cerebrum.

Each half, or *hemisphere*, has its external surface subdivided into five lobes by the

Fissure of Sylvius, f. Sy. p., beginning at anterior perforated space, it passes to external surface of hemisphere and subdivides, one arm (*f. Sy. a.*) ascends toward longitudinal fissure, one passes nearly horizontally backward.

The *fissure of Rolando, f. Ro.*, commences at or near middle of the longitudinal fissure, running downward and forward to a little above the horizontal branch of the Sylvian fissure.

Parieto-occipital fissure, par. oc. f., commences about midway between posterior extremity of brain and fissure of Rolando, running downward and forward for a variable distance.

The *frontal lobe* lies in front of the fissure of Rolando, and above the horizontal part of the Sylvian fissure; its under surface is called the *orbital lobe*.

FIG. 77.



Convolutions of outer surface of brain.

The *parietal lobe* is bounded in front by fissure of Rolando, behind by parieto-occipital fissure, and below by horizontal limb of Sylvian fisure.

The *occipital lobe* lies behind parieto-occipital fissure.

The *temporo-sphenoidal lobe* occupies middle cerebral fossa of skull, and is limited above and in front by Sylvian fissure.

The *central lobe*, or island of Reil, lies in the fissure of Sylvius at the base of brain.

The inner or median surface of each hemisphere presents five fissures:

The *calloso-marginal fissure, c. m. f.* (Fig. 78), separating the marginal convolution from the gyrus fornicatus, *g. f.* (= limbic [illegible].

The *parieto-occipital fissure, par. oc. f.*, a continuation of that of same name on outer surface of hemisphere.

The *calcarine fissure, calc. f.*, runs from back of hemisphere horizontally forward to posterior inferior extremity of gyrus fornicatus, *g. f.*; it is joined by the parieto-occipital fissure.

The *collateral fissure, coll. f.*, runs below and nearly parallel to preceding, separated by the uncinate gyrus.

[illegible]

The *dentate fissure, d. f.*, commences below posterior extremity of corpus callosum, running forward to end at recurved part of uncinate gyrus.

The *transverse fissure,* between the middle lobe and crus cerebri, admitting pia mater into the lateral ventricles.

Mention the principal convolutions of the brain.

The convolutions, or *gyri,* are elevated ridges covered with gray matter, separated by deep furrows, or *sulci,* thus securing a great extent of gray matter; while not uniform in all brains, nor symmetrical, certain principal convolutions are constant, such as

The *gyrus fornicatus, g. forn.*, that lying over corpus callosum.

The *marginal, marg. c.*, forming anterior superior margin of the great longitudinal fissure.

The *ascending frontal, asc. fr. c.* (Fig. 77), forming anterior boundary of fissure of Rolando.

FIG. 78.



Convolutions of outer surface of brain.

The *ascending parietal, asc. par. c.*, the posterior boundary of same fissure.

Angular gyrus, ang. c., lying between back of horizontal limb of Sylvian fissure, and a short fissure running upward from it.

Other convolutions are indicated by contractions of names on the diagrams, so that with the preceding explanations of points of reference no further details are requisite.

What is the quadrate lobe (Fig. 78), and what is the cuneus (Fig. 78)?

The former is the marginal convolution between the calloso-marginal fissure in front and the parieto-occipital behind; the latter lies between the parieto-occipital and calcarine fissures.

Mention the various objects seen on the under surface of cerebrum, naming them from before backward.

Longitudinal fissure[4] (Fig. 79), antero-posterior.

FIG. 79.



Base of brain.

Corpus callosum, the transverse commissure, connecting cerebral hemispheres.

Lamina cinerea, a thin layer of gray substance, extending from end of corpus callosum back to tuber cinereum above optic tracts.

Fissure of Sylvius[6], separating by its horizontal limb the frontal and parietal lobes from temporo-sphenoidal; (*old nomenclature*) separates anterior and middle lobes chiefly by the ascending or precentral limb.

Anterior perforated spaces[7], between roots of olfactory nerve on each side for passage of vessels into ~~corpus striatum~~.

Optic commissure[24], union of optic tracts.

Tuber cinereum, an eminence of gray matter, part of floor of third ventricle.

Infundibulum[8], a hollow conical process of gray matter projecting from middle of under surface of tuber cinereum, communicating with ~~third ventricle~~.

Pituitary body, a vascular, two-lobed body projecting from apex of infundibulum into sella turcica of sphenoid.

Corpora albicantia[9], the two small rounded, infolded terminations of the anterior crura of ~~fornix~~.

Posterior perforated space[10], between corpora albicantia in front, pons Varolii behind, and cerebral crura on either side, forms part of floor of third ventricle, and gives passage for vessels to optic thalami.

Crura cerebri[11] (*peduncles of cerebrum*) connect the cerebrum with cerebellum, medulla oblongata, and spinal cord.

Pons Varolii[12] (see page 195).

Name the five great ganglia of the brain, other than the gray matter of the cerebral and cerebellar hemispheres and medulla oblongata.

Olfactory bulbs[23],
Corpora striata,
Optic thalami,
Tubercula quadrigemina.
Pons Varolii[12],

Describe these ganglia.

Olfactory bulbs[23], the ganglia of the sense of smell, lie in olfactory grooves of cribriform plate of the ethmoid, and arise by two white roots, one crossing Sylvian fissure from a nucleus of gray matter in middle lobe, the other from inner back part of frontal (*orbital*) lobe, and a gray root from under surface of same lobe.

Corpora striata, ~~chief part of~~ ~~motor~~ tract, lying in lateral ventricles, the intra-ventricular gray portion called *caudate nucleus*, the extra-ventricular, the *lenticular nucleus*, the *internal capsule* dividing the two.

Optic thalami, they are white externally, the ganglia of general [redacted], forming the lateral boundaries of the third ventricle, each presenting an anterior tubercle in lateral ventricle, and posteriorly two small *internal* and *external geniculate bodies*.

Corpora quadrigemina (optic lobes) are four rounded eminences situated behind third ventricle and posterior commissure; the anterior pair are the *nates*, the posterior pair the *testes*; they are the centre of vision.

The *pons Varolii*, containing much gray matter and one special mass called the *superior olivary* body, might be considered a fifth ganglion.

What are the commissures?

Connecting bands of white or gray matter.

Enumerate them.

Those pursuing an antero-posterior course, are

Olfactory tracts,
Tœnia semicircularis,
Crura cerebri,
Peduncles of pineal gland,
Fornix,
Gyrus fornicatus,
Fasciculus unciformis,
Processus e cerebello ad testes.
Striæ longitudinales.

Those passing transversely, are

Anterior,
Middle,
Posterior,
Corpus callosum,
Optic chiasm,
Pons Varolii,
Fornix (also a longitudinal commissure),
Posterior medullary velum,
Valve of Vieussens.

What are the ventricles of the brain?

Five serous cavities in the brain, four of which intercommunicate; they are *two lateral*, a *third*, a *fourth*, and a *fifth*.

Describe the lateral ventricles.

The *corpus callosum* roofs them in; each has an *anterior cornu*, curving outward and forward into the anterior lobe, a *middle cornu* descending into the middle lobe downward and backward, then turning downward and forward, containing the curved *hippocampus major*, and a *posterior cornu* curving downward and inward into the posterior lobe, containing the hippocampus minor; the floor from before backward is formed by the *corpus striatum*, *tænia semicircularis*, *optic thalamus*, *choroid plexus*, *corpus fimbriatum*, and *fornix*; the septum lucidum forms the inner wall.

Describe the following parts:

The *corpus callosum*; it is a thick arched layer of transverse fibres at the bottom of the longitudinal fissure, anteriorly curving upon itself, giving off two peduncles to the entrance of the Sylvian fissure, posteriorly it becomes continuous with the fornix; a ~~median~~ linear depression on its upper surface is called the ~~*raphé*,~~ parallel to which on each side run two or more elevated longitudinal bands, the ~~*striæ longitudinales*~~ or *nerves of Lancisi*.

The *tænia semicircularis*; lies in a depression between corpus striatum ~~and optic tha~~lamus and is a commissure between these bodies.

Choroid plexus; the vascular margin of the velum interpositum, communicating with that of opposite side just behind anterior pillars of fornix, through the oval *foramen of Monro*; posteriorly, it descends into the middle horn, becoming there continuous with pia mater through transverse fissure.

The *corpus fimbriatum* (*tænia hippocampi*); the lateral edge of the posterior pillar of the fornix, forming a white band just behind choroid plexus.

The *fornix*; continuous with corpus callosum behind, consisting of a ~~triangular~~ body with ~~apex forward~~, two *anterior crura*, curving downward to base of brain, there to form the *corpora albicantia* and terminate in the optic thalami, and two ~~*posterior crura*~~, running down the middle cornua of the lateral ventricles, as the ~~*hippocampi majores*~~; on the back under surface, between posterior crura, certain transverse longitudinal and oblique lines have been termed the *lyra*.

The *septum lucidum*; is a vertical septum, attached above to under surface of corpus callosum, below to anterior part of fornix and prolonged portion of corpus callosum; it is triangular in form, and consists of two lamina of ~~white nerve~~ matter lined internally with gray matter, the space between ~~forming the *fifth ventricle*~~, not communicating with other ventricles.

What is the pes hippocampi ?

The rounded elevations of lower extremity of hippocampus major.

What is the pes accessorius ?

A white projection at the junction of the two hippocampi.

What is the fascia dentata ?

The gray substance of the dentate convolution, seen by raising the edge of corpus fimbriatum.

What is the internal capsule ?

A large tract of white matter belonging to pyramidal tract, consisting of two limbs, the *anterior* lying between anterior part of lenticular nucleus and caudate nucleus, the *posterior* between posterior part of lenticular nucleus and thalamus, which, by their junction, form a projection inward, the *knee* or *genus.*

What is the external capsule ?

A small tract of white matter lying between the outer part of the lenticular nucleus and cerebral cortex.

Describe the third ventricle.

Narrow and oblong, the lateral walls being the optic thalami and peduncles of pineal gland; above the under surface of the velum interpositum roofs it in, containing the choroid plexuses of this ventricle; the lamina cinerea, tuber cinereum, infundibulum, corpora albicantia, and posterior perforated space form its floor; it is limited in front by the anterior crura of the fornix and part of anterior commissure; behind, by the posterior commissure, beneath which opens the *iter e tertio ad quartum ventriculum* ; while in front by the foramen of Monro it communicates with the lateral ventricles; the cavity is crossed by an anterior and posterior *white commissure* and a middle *gray one.*

Describe the fourth ventricle.

It is the space between posterior surface of the medulla oblongata and pons in front, and the cerebellum behind. Its *roof* is the valve of Vieussens and that portion of the cerebellum called the *nodulus, uvula, and amygdalæ*; the *lateral boundaries* on each side are the processus e cerebello ad testes, and posterior pyramids and restiform bodies of the medulla oblongata; the *floor* consists of the posterior surface of the medulla oblongata and the pons, upon the former of which will be seen the posterior median fissure obliterated above, terminating below in front of *calamus scriptorius* at the orifice of a short blind canal, the remains of the fœtal central canal of cord; on each side of median fissure are two convex longitudinal eminences, the *fasciculi teretes*; external to these, opposite crus cerebri on each side, is the *locus cœruleus*—a ganglionic mass; a thin streak of this embedded gray matter continued to top of ventricle is called the *tænia violaceæ*, and certain white lines, the *lineæ transversæ*: this cavity communicates with that of the third by iter e tertio, etc., and with the subarachnoid space, through an opening in the pia mater extending between the medulla oblongata and cerebellum.

The fifth ventricle has been described with septum lucidum, p. 194.

Describe the pons Varolii.

It is the bond of union between the different segments of the encephalon: bridging the medulla oblongata is a broad transverse band forming the *middle peduncles* or *crura cerebelli*: longitudinal fibres pass up from the medulla oblongata to form the *two crura cerebri*, which become connected with the corpora striata and optic thalami, through which they pass to reach the gray matter of the hemispheres forming the *corona radiata*: the *locus niger*, a mass of gray matter, is found in the substance of each crus; the third nerve emerges from inner side of each crus, the fourth nerve around the outer side from above, while the optic tract is adherent by its upper border.

Describe the corpora quadrigemina.

These are four rounded projections placed just behind third ventricle beneath posterior border of corpus callosum: they are the centre of vision. Two white bands on each side connect them with optic thalamus and optic tracts, those passing from the anterior pair, or *nates*, to thalamus are the *brachia anteriora*, those running from the posterior pair, the *testes*, to the thalamus are the *brachia posteriora*; passing from the testes to the cerebellum on each side is the *processus e cerebello ad testes* or *inferior cerebellar peduncle.*

What is the valve of Vieussens?

A thin lamina of nerve tissue, stretching from the vermiform process of cerebellum from one processus e cerebello ad testes to the other, forming the ~~roof of the iter e tertio~~, etc.: a little ridge descending upon the upper part from the corpora quadrigemina is the *frœnulum*, and on either side are the transverse fibres connecting the fourth nerves.

What is the pineal gland?

A reddish conical body lying beneath the nates, connected by its base by two *peduncles* to the anterior crura of fornix: the gland has a small cavity—said by some to open into that of the third ventricle—containing a viscid fluid and a sandy substance, the *acervulus cerebri*, composed of calcium carbonate and phosphate, magnesium and ammonium phosphate, with some animal matter.

Describe the medulla oblongata.

It is the upper enlarged part of spinal cord, extending from upper border of atlas to lower border of pons Varolii; its posterior surface forms the floor of the fourth ventricle, its anterior rests on the basilar groove of the occiput. It contains the vaso-motor, cardiac, and respiratory centres, also those of deglutition, mastication, etc. Divided into lateral halves by the *anterior* (*d*, Fig. 80) and,

Posterior median fissures, it presents on each side of anterior fissure,

The *anterior pyramid* (*p*, *a*, Fig. 80) formed by antero-lateral columns of the cord, these latter fibres *decussating* above *d* (crossing from one pyramid to that of the other) at the lower part; behind the pyramid is the *olivary body*, containing in its interior a capsule of gray matter, the *corpus dentatum*; behind the olivary body is, The *lateral tract* continuous with lateral tract of the cord; back of this is the *restiform body*, continuous with posterior columns of cord below, above passing into corresponding hemisphere of cerebellum forming its *inferior peduncle*; running along the posterior median fissure on either side are the

Posterior pyramids continuous with posterior median. columns of cord; diverging above, they form the lateral boundaries of the *calamus scriptorius.*

Describe the cerebellum.

It is that portion of the encephalon contained in the inferior occipital fossæ composed of *laminæ* covered with gray matter; on the upper surface the two *hemispheres* are seen connected by a median elevated lobe, the *superior vermiform process*, while they are separated in front and behind by notches respectively called the *incisuræ cerebelli anterior and posterior*; the superior vermiform process consists of a *lobulus centralis* in incisura anterior, the *monticulus cerebelli* the central projecting portion, and the *commissura simplex*, near the incisura posterior.

Describe the under surface of the cerebellum.

It is divided by a central longitudinal depression, the *valley*, into two hemispheres. Projecting from the bottom of the valley is the *inferior vermiform process*, consisting of the *commissura brevis* in the posterior notch, in front of the conical *pyramid*, more anteriorly, the *uvula*, lying between two rounded lobes the *amygdalæ* or *tonsils*, and which projects into fourth ventricle, and finally in front of uvula the *nodule*; attached to each side of nodule and also to flocculus. is a thin white layer which together form the *posterior medullary velum.*

Name the lobes of the cerebellum.

Below, from before backward, they are, on each side,

The *flocculus, or pneumogastric lobule,* a prominent tuft below and behind middle peduncle.

The *amygdala*, just described.

The *digastric*, on outside of tonsils, partially connected with pyramid.

The *slender*, behind former, connected with commissura brevis and back of pyramid.

The *inferior posterior*, joining the commissura brevis in the valley.

Only one *fissure*—the *great horizontal*—pertains to the cerebellum, commencing in front at pons, and passing horizontally round free margin of each hemisphere to median line; from this numerous secondary fissures proceed, marking out lobes, as two on the upper cerebellar surface on each side, viz., the *anterior*, or *square globe*, extending back to posterior edge of vermiform process, and the *posterior*, or *semilunar*, passing from preceding to great horizontal fissure.

How many peduncles has the cerebellum?

Three; the transverse fibres of pons Varolii, or *middle peduncle*; on each side the restiform bodies of medulla oblongata, or *inferior peduncles*; and the processus e cerebello ~~ad~~ testis, forming the *~~supe~~rior peduncles*.

Describe the arrangement of the gray matter of the cerebellum.

A vertical section reveals a central stem of white matter containing a capsule of gray matter, the *corpus dentatum*. From the central white stem ten or twelve plates, or *laminæ*, spring, giving origin to smaller secondary and tertiary laminæ, covered externally by a layer of gray matter, so that the cut surface presents the foliated appearance giving origin to the name *arbor vitæ*.

The Spinal Cord.

What is the spinal cord?

The elongated cylindrical part of the cerebro-spinal axis contained in the vertebral canal, measuring about ~~seventeen inches~~ in length, extending from upper border of atlas to lower border of body of first lumbar vertebra, presenting a *~~cervical~~ enlargement* from third ~~cervical to first or second dorsal~~ vertebra, and a *lumbar enlargement* opposite last ~~two or three dorsal~~ vertebræ; it terminates by a slender filament of gray substance, the *filum terminale*. The white matter is disposed externally, the gray internally in the form of two crescents joined by a transverse commissure, the anterior thicker extremities, forming the *anterior cornua*, or *horns*, the posterior the *posterior cornua.*

How many membranes has the cord?

Three; the *dura mater*, continuous with that of brain, but separated from bony walls by loose areolar tissue, containing a plexus of veins; the *arachnoid*, continuous with cerebral arachnoid, also having a *subdural* and ~~*subarachnoidean* space~~, the latter communicating with general ~~ventricular~~ cavity of brain by the *foramen of* ~~*Magendie*~~, an aperture in pia mater of fourth ventricle—this space contains an abundant serous secretion, the *cerebro-spinal fluid*; and the *pia mater*, sending processes down into anterior and posterior median fissures, having medianly in front a fibrous band, the *linea splendens*, and on each side another, the *ligamentum denticulatum*, whose outer border presents about twenty serrations, the apices of each attached to inner surface of dura mater, serving to support the cord.

Describe the fissures.

They are the *anterior and posterior median*, dividing cord into two lateral halves, joined medianly by a white *commissure*; on each side of anterior fissure is a series of foramina for the exit of anterior roots of nerves; this line of openings is called the *antero-lateral fissure*; two *postero-lateral fissures* run parallel to posterior median fissure, giving exit to posterior roots; finally, a delicate groove on each side between the postero-lateral and posterior median fissures exists, marking off the *posterior pyramids.*

Name the columns of the cord marked off by these fissures.

On each side the *anterior*, continuous with the anterior pyramid of medulla oblongata.

The *lateral*, continuous with lateral tract of medulla oblongata.

The *posterior*, continuous with restiform body of medulla oblongata.

The *posterior median*, continuous with posterior pyramid of medulla oblongata.

What is the ventricle of the cord?

The fœtal central canal, usually obliterated, except for a few lines below floor of fourth ventricle of brain.

The Cranial Nerves.

How many pairs of cranial nerves are there?

Nine, according to most anatomists; twelve, according to others.

Describe each of the following nerves, giving their number according to each classification.

First, Olfactory $^{1'}$ (Fig. 80), nerve of smell; *superficial origin*, by one root from middle lobe, two from anterior lobe (see p. 191); *deep origin*, gray nucleus in temporo-sphenoidal lobe, corpus striatum, and gyrus fornicatus; *escapes*, by foramina of cribriform plate of ethmoid; *distribution*, nasal (Schneiderian) mucous membrane; may be regarded as lobe of cerebrum.

→ optic F

FIG. 80.



Second, Optic ^{11}, nerve of sight; *deep origin*, from optic tracts $^{11'}$, which arise from optic thalami, corpora geniculata, and nates of corpora quadrigemina; is also attached to crus cerebri and receives fibres from tuber cinereum, lamina cinerea, and, according to some, the tænia semicircularis and gyrus fornicatus; in front of tuber cinereum lies the optic *chiasm* or *commissure*, formed of six sets of fibres, viz., a set crossing from right side of brain to left eye, a second pursuing the same course from the opposite side, *decussating*

1 2 3 etc

On Old Olympus Towering Top a Fat-
man Picked Some Hops.

fibres; a third, anterior, connecting the two retinæ—*inter-retinal fibres*; a fourth and fifth, lateral, *cerebro-retinal*, connecting the hemisphere of one side with retina of eye of same side; and a sixth, posterior—*inter-cerebral*—connecting one optic tract with the other; *distribution*, retina.

Third^{iii}, Motor oculi; *superficial origin*, inner surface of crus cerebri; *deep origin*, a nucleus for each in floor of iter e tertio ad quartum ventriculum; *escapes*, by sphenoidal fissure; *distribution*, to all eye muscles including iris, except external rectus and superior oblique. 4° 6°

Fourth^{iv} (Fig. 80), Patheticus, motor, *superficial origin*, outer side crus cerebri; *deep origin*, floor of aqueduct of Sylvius; *escapes* through sphenoidal fissure; *distribution* to superior oblique of eye.

FIG. 81.



Fifth^{v} (Fig. 80), Trigeminus or Trifacial, nerve of general sensation, motion, and taste; *superficial origin*, by a motor and a sensory root from side of pons Varolii; *deep origin*, sensory root from upper expanded posterior gray horns of medulla at junction with resti-

Tenth, Pneumogastric^{x} (Fig. 80) or *par vagum of the eighth nerve;* both motor and sensory; *superficial origin,* from lateral tract of medulla oblongata behind olivary body and below the glosso-pharyngeal; *deep origin,* gray nucleus lower part of floor of fourth ventricle—the motor filaments probably come from spinal accessory; *escapes,* by jugular foramen presenting a gangliform enlargement, *ganglion of the root,* and lower another, *ganglion of the trunk*; *distribution,* to organs of voice and respiration, the pharynx, œsophagus, stomach, and heart; the branches are

Auricular, (Arnold.)
Pharyngeal, ([illegible] Plexus)
Superior laryngeal,
Recurrent laryngeal,
Cervical cardiac,
Thoracic cardiac,
Anterior pulmonary,
Posterior pulmonary,
Œsophageal,
Gastric,
Hepatic (to hepatic sympathetic plexus).

Eleventh, Spinal accessory^{xi} (Fig. 80), or *third division of the eighth nerve,* a motor nerve; *superficial origin,* lateral tracts of medulla oblongata below roots of vagus, and from same part of spinal cord as low as sixth cervical vertebra; *deep origin,* gray matter below nucleus of vagus and anterior horn of gray matter of cord; it sends filaments to ganglion of root of vagus and pharyngeal and superior laryngeal branches of same nerve; *escapes,* by jugular foramen, the spinal portion first entering skull through foramen magnus; *distribution,* to sterno-cleido-mastoid and trapezius muscles, communicating with second, third, and fourth cervical nerves.

Twelfth, Hypoglossal^{xii} (Fig. 80), or ninth nerve, motor nerve of tongue; *superficial origin,* groove between pyramidal and olivary bodies by numerous filaments; *deep origin,* special nucleus at lowest point of fourth ventricle; *escapes,* by anterior condyloid foramen; *distribution,* to thyro-hyoid, genio-hyoid, stylo-glossus, hyo-glossus, genio-hyo-glossus, and by descendens noni to sterno-hyoid, sterno-thyroid, and omo-hyoid muscles; it communicates with the pneumogastric, gustatory of fifth, sympathetic, and first and second cervical nerves; its branches are

Descendens ~~noni~~ Hypoglossi, *Muscular,* *Thyro-hyoid,* *Meningeal.*

oblongata, from groove between olivary and restiform bodies; *deep origin*, from nucleus of fasciculus teres in floor of fourth ventricle and nucleus of motor root of fifth nerve; *escapes*, by internal auditory meatus to aquæductus Fallopii, and thence by stylo-mastoid foramen; *distribution*, to muscles already mentioned; its branches are

Sens. to tongue. → LINQUALIS N.

Tympanic,		*Temporal,*
Chorda tympani,	*Temporo-facial*	*Malar,*
Posterior auricular,		*Infra-orbital,*
		Buccal,
Digastric,	*Cervico-facial*	*Supra-maxillary,*
Stylo-hyoid,		*Infra-maxillary.*

The communications of the facial are

With auditory nerve; with Meckel's ganglion by large petrosal nerve; with otic ganglion by small petrosal nerve; with sympathetic on middle meningeal by external petrosal nerve; with pneumogastric, glosso-pharyngeal, carotid plexus, auricularis magnus, auriculo-temporal and with the three divisions of fifth nerve.

Eighth. Auditory$^{\text{viii b}}$ (Fig. 80), or *portio mollis of the seventh nerve*, nerve of hearing; *superficial origin*, from medulla oblongata in groove between olivary and restiform bodies at lower border of pons; *deep origin*, superior vermiform process of cerebellum, gray substance of posterior pyramid and restiform body; *escapes*, by internal auditory meatus; *distribution*, to internal ear; its branches are,

Vestibular, to the vestibule, *Cochlear*, to the cochlea (see p. 320).

Ninth. Glosso-pharyngeal$^{\text{viii a}}$ (Fig. 80) or first divsion of eighth nerve, nerve of general sensation and taste; *superficial origin*, medulla oblongata just behind olivary body; *deep origin*, gray nucleus at lower part of floor of fourth ventricle; *escapes*, at central part of jugular foramen, after which it presents two gangliform enlargements, the *jugular and petrous ganglia*; *distribution*, to muscles of pharynx, mucous membrane of pharynx, fauces, tonsil, and tongue, and the middle ear; its branches are

Tympanic (Jacobson's),	*Pharyngeal,*	*Tonsillar,*
Carotid,	*Muscular,*	*Lingual.*

Describe the brachial plexus with its branches.

The anterior divisions of the fifth and sixth, with a branch of the seventh cervical nerves form the *outer cord,* the eighth cervical and first dorsal form the *inner cord,* a branch from the latter and one from the united fifth and sixth join the seventh nerve to form the *posterior cord*; this is the usual, but not *invariable* arrangement; its branches are,

N. above clavicle

Communicating, above clavicle to phrenic.

Muscular, to longus colli, scaleni, rhomboidei, and subclavius muscles.

Posterior or *long thoracic* (external respiratory nerve of Bell), *from* fifth, sixth, and seventh nerves, *to* serratus magnus muscle.

Suprascapular, from outer cord; *to* supra- and infra-spinatus muscles and shoulder-joint.

External anterior and internal anterior thoracic, the former *from* outer cord, the latter *from* inner cord; both *to* pectoral muscles.

Shoulder

The *three scapular,* the upper *from* communicating branch from outer to posterior cord, the other two from posterior cord; *to* subscapular, teres major, and latissimus dorsi muscles.

Circumflex, from posterior cord with musculo-spiral; *to* deltoid and teres minor muscles, the shoulder-joint, and skin of lower deltoid region.

3T To M. [illegible]

Musculo-cutaneous (external cutaneous), *from* outer cord and pierces the coraco-brachial muscle; *to* coraco-brachial, biceps, brachialis anticus muscles, elbow-joint, and skin of outer half of front of forearm.

Internal cutaneous, from inner cord with ulnar and inner head of median; *to* skin over biceps and that of inner half of forearm in front and behind.

Lesser internal cutaneous (nerve of Wrisberg), *from* inner cord alone, or a branch from this joined with intercosto-humeral (the lateral cutaneous branch of second intercostal nerve, piercing external intercostal muscle to supply skin of upper half of inside of arm), or, again, the intercosto-humeral may entirely replace it; *to* skin of inner side of arm.

Median, from outer and inner cords by two roots which embrace the axillary artery uniting in front or to its outer side, first lying external to brachial artery, then crossing to its inner side, passing between heads of pronator radii teres muscle to run between deep

outside, top, inside (Virg.)

and superficial flexor of fingers to within two inches of wrist, when it becomes superficial; its branches are,

Muscular, to all anterior superficial forearm muscles except flexor carpi ulnaris.

Anterior interosseous, to deep forearm muscles except inner half of flexor profundus digitorum.

Palmar cutaneous, piercing fascia above wrist; *to* skin of palm to radial side.

Muscular, to abductor, opponens, and outer head of flexor brevis pollicis.

Five digitals, supplying both sides of thumb, index, middle, and radial side of ring fingers.

Ulnar, from inner cord runs behind inner condyle, thence passing into forearm between heads of flexor carpi ulnaris to run some distance from ulnar artery at upper third, but close to it for rest of its extent; it partially supplies elbow and wrist-joints, both sides of little and ulnar side of ring finger and skin of forearm and hand; its branches are,

Articular.

Muscular, to flexor carpi ulnaris, inner half of flexor profundus digitorum, the little finger muscles, the interossei and lumbricals, palmaris brevis, adductor, and inner head of flexor brevis pollicis.

Cutaneous,
Dorsal cutaneous,
Superficial palmar.
Deep palmar (muscular).

Musculo spiral, from posterior cord and branch of inner cord, running in same-named groove with superior profunda vessels to divide in front of condyle into the *radial* and *posterior interosseus*; its branches are,

Radial,
Cutaneous,
Muscular,
Posterior interosseous.

Radial, to skin of ball and outer side of thumb, and that on back of index, middle, and part of ring fingers;

Posterior interosseous, supplying wrist-joint and all muscles on back of forearm except anconeus, supinator longus, and extensor carpi radialis longior.

Muscular, to triceps, anconeus, supinator longus, extensor carpi radialis longior, and brachialis anticus.

Cutaneous, to skin of lower, outer, and back part of arm, forearm, and hand.

Dorsal Nerves.

Describe them.

Twelve in number on each side, the first escapes between first and second dorsal vertebræ, the last between the last dorsal and first lumbar; they divide into an *anterior* and *posterior* division, the latter supplying spine, extensor muscles of back and dorsal integument; the former (anterior) are,

The *intercostal nerves*, each connected by one or two filaments with the adjacent sympathetic ganglia: the anterior division of the *first* nerve aids in the formation of the brachial plexus, its intercostal branch is small and gives off no lateral cutaneous branch: the lateral branch of the *second* nerve is the intercosto-humeral (see p. 206): the remaining nerves give off *lateral cutaneous* branches supplying skin of front of thorax and abdomen, while they give *muscular branches* to the intercostal and abdominal muscles.

Describe the lumbar nerves.

The posterior branches resemble in origin and arrangement those of the dorsal region, while their anterior branches form the lumbar and part of the sacral plexuses.

Describe the lumbar plexus and branches.

It is formed by loops of communication between the anterior divisions of the four upper lumbar nerves ~~*in the substance*~~ of the ~~psoas muscle~~, thus,

1 ~~The first nerve~~ gives off,

Ilio-hypogastric, to skin of gluteal and hypogastric regions.

Ilio-inguinal, to internal oblique muscle and skin of scrotum (labium in female), and upper inner part of thigh, and

2 ~~A~~ *Communicating loop, to* ~~second lumbar nerve~~, from which arises, in conjunction with a branch from the third nerve, the

External cutaneous, to skin of antero-external and posterior surfaces of thigh: from the second nerve and loop from first nerve comes, the

Genito-crural, to cremaster muscle and skin of front of upper part of thigh; and a

Communicating loop, to third lumbar nerve. By a branch of the

+ 4 ~~third~~ and fourth nerves with fibres from the second is formed the

Obturator, *to* obturator externus and adductor muscles, hip- and knee-joints, also sometimes to skin of thigh and leg.

The *accessory obturator*, either from obturator or formed by two filaments from third and fourth nerves, *to* pectineus muscle and hip-joint.

Communicating loop, between third and fourth nerves.

Communicating loop, between fourth and fifth nerves.

Anterior crural, *from* third and fourth, with communicating branch from second lumbar nerve. It descends beneath Poupart's ligament external to artery between the psoas and iliacus muscles after emerging from former, and divides into an anterior and posterior division: its branches are, within the pelvis, 2

Muscular to iliacus internus, to *femoral artery*; external to pelvis,

II Ant. { *Middle cutaneous*, Post. { *Articular* (hip and knee),
P. { *Internal cutaneous*, P. { *Muscular*.
Muscular *Long saphenous*,

It supplies all the anterior thigh-muscles except tensor ~~vaginæ~~ fasciae femoris, and skin of front and inner side of thigh, leg, and foot.

Describe the sacral plexus. cov. by pelvic fascia [illegible]

It is formed by the lumbo-sacral cord (the anterior division of fifth nerve with a branch of the fourth), and anterior divisions of three upper sacral nerves and part of that of fourth: it rests upon the anterior surface of the ~~pyriformis musc~~le, giving off the

Superior gluteal, *from* lumbo-sacral cord passing out through great sacro-sciatic foramen, *to* gluteus medius and minimus and tensor vaginæ Fasciae ~~femoris~~ ~~muscles~~.

Muscular branches, *to* pyriformis, obturator internus, gemelli, and quadratus femoris.

~~*Pudic*~~ passes out of pelvis by greater sacro-sciatic foramen to reënter it by lesser sacro-sciatic foramen: its branches are,

Inferior hæmorrhoidal, *Perineal*, *Dorsal of penis*.

Muscular, to transversus peronei, accelerator urinæ, erector penis and compressor urethræ muscles: the preceding branches supply analogous muscles and parts in female.

Small sciatic, escapes by greater sacro-sciatic foramen, *to* ~~skin of~~ scrotum, back of leg and thigh, and gluteus maximus muscle. INF. Glute

~~*Great sciatic*~~, is a continuation of lower part of sacral plexus, forming the largest nerve in the body. Escaping by the greater sacro-sciatic foramen below pyriformis muscle, it gives off these branches,

Articular (to hip).

Muscular, to biceps, semi-tendinous, semi-membranous and adductor magnus, and divides into the *internal* and *external popliteal nerves.*

Give the branches of the internal popliteal nerve.

Articular (to knee).

Muscular to gastrocnemius, plantaris, soleus, and popliteus.

External or short saphenous, a ~~cutaneous bra~~nch communicating with external popliteal, musculo-cutaneous, and small sciatic nerves. Ext. Long., Flex. [illegible] Foot. [illegible]

Describe the posterior tibial nerve.

It is a continuation of ~~former~~ Int. Pop., runs down middle of back of leg beneath the calf muscles to divide between the inner malleolus and the heel into the internal and external plantar: its branches are,

Muscular, to tibialis posticus, flexor longus digitorum and pollicis muscles.

Plantar cutaneous, to skin of heel and inner side of sole of foot.

Internal plantar, supplying skin of sole, tarsal, and metatarsal articulations, the two inner lumbricals, abductor pollicis, and flexor brevis digitorum, with four *digital* branches which supply both sides of first, second, and third toes and inner half of fourth.

External plantar, smaller than preceding, supplying flexor accessorius, abductor minimi digiti, flexor brevis minimi digiti, all the interossei, two outer lumbricales, adductor pollicis, transversus pedis muscles, and skin of little toe and adjoining side of fourth toe.

Describe the external popliteal nerve. (Peroneal.

It descends obliquely along outer side of popliteal space close to tendon of biceps, giving off,

Two *articular branches* to knee; *cutaneous branches* to skin of postero-external surface of leg, and a branch, the *communicans peronei*, joining external saphenous nerve: it divides into the

Anterior tibial, about one inch below head of fibula, giving *mus-*

cular branches to tibialis anticus, extensor longus digitorum, peroneous tertius, extensor proprius pollicis, and extensor brevis digitorum, *articular* to tarsal and metatarsal joints, and *cutaneous* to skin of adjacent sides of great and second toes;

Musculo-cutaneous, giving off *muscular branches*, to peroneus longus and brevis, and *cutaneous*, to skin of lower part of leg, and dorsum of foot and toes, except outer side of little and adjoining sides of great and second toes.

✓ **Describe the sacral and coccygeal nerves.**

They are five in number, their long roots forming the *cauda equina* in the spinal canal; each divides into an *anterior* (see page 209) and *posterior division*, the latter escaping by the posterior sacral foramina, except the fifth, which emerges between sacrum and coccyx, and supplies multifidus spinæ muscle and skin of posterior gluteal region; a branch goes from third nerve to bladder; the two lower nerves join with coccygeal, forming loops, which supply skin over coccygeal region and coccygeus, levator, and sphincter ani muscles.

The Sympathetic Nerve.

What is the sympathetic nerve or system?

It consists of a series of ganglia with intervening commissural bands, forming two cords on either side of spinal column connected above by *ganglion of Ribes* on the anterior cerebral communicating artery, and below by *ganglion impar*, in front of coccyx; they also communicate with the spinal system of nerves.

Locate and briefly describe the connections of the cranial and facial ganglia.

Ganglion of *Ribes*, on anterior communicating artery, *connected* with cavernous and carotid plexuses.

Carotid ganglion (Laumonier's) on under surface of vessel.

milunar - *Gasserian*, on fifth nerve (infra).

ital — *Ciliary* or *ophthalmic*, in orbit (infra).

yg. pal — *Spheno-palatine* (Meckel's), in spheno-maxillary fossa (infra). [illegible]

icular — *Otic* (Arnold's), on inner side of inferior maxillary nerve below foramen ovale (infra).

(*Ganglia* are occasionally found on middle meningeal, lingual, temporal, and pharyngeal arteries, receiving the name of their discoverer, as Bidder, Cloquet, etc.

Ganglion of Bochdalek is situated at point of junction of one of the nasal branches of Meckel's ganglion and anterior dental nerve. p. 597

Submaxillary lies above deep portion of submaxillary gland (infra).

[illegible]. Aq. of Fallop. Rt.= Large [illegible]

Describe the ciliary ganglion.

Situated in the orbit between the optic nerve and external rectus muscle, its roots are, *sensory*, from nasal branch of ophthalmic (fifth pair); *motor*, third nerve; *sympathetic*, cavernous plexus; its branches are,

Short ciliary to ciliary muscle and iris.

Describe the spheno-palatine ganglion.

Deeply placed in spheno-maxillary fossa, its roots are, *sensory*, from superior maxillary; *motor*, facial, through Vidian; *sympathetic*, from carotid plexus; its branches are,

Ascending,
Anterior palatine,
Posterior palatine,
Middle palatine,
Superior nasal,
Naso-palatine,
Vidian,
Pharyngeal, or
Pterygo-palatine.

Describe the otic ganglion.

It lies immediately beneath foramen ovale on the inner surface of inferior maxillary nerve; its roots are, *sensory*, from auriculo-temporal; *motor*, internal pterygoid branch of inferior maxillary, also facial and glosso-pharyngeal through continuation of small petrosal nerve; *sympathetic*, from middle meningeal plexus; its branches supply ten~~sor~~ tympani and te~~nsor palati~~ muscles.

Describe the submaxillary ganglion.

Situated above deep portion of submaxillary gland, its roots are, *sensory*, from gustatory branch of inferior maxillary; *motor*, facial, through chorda tympani; *sympathetic*, from facial plexus; its branches go to mucous membrane of mouth, to submaxillary gland and to its duct.

Describe the petrosal nerves.

The *great petrosal* (large superficial petrosal) is usually described as a branch of the spheno-palatine ganglion passing through the

Vidian canal, distributing twigs to mucous membrane of back part of nose, septum, and end of Eustachian tube, entering skull through foramen lacerum medium, where it ~~divides~~ into the ~~*large superficial*~~ *petrosal*, which enters hiatus Fallopii, receives a twig from Jacobson's nerve, and ~~terminates~~ in the ~~geniculate ganglion~~ of facial nerve, and the ~~*carotid* (*large deep petrosal*)~~ joining the carotid plexus.

~~*Small petrosal* (small superficial~~ petrosal) connects ~~geniculate~~ ganglion of seventh nerve with ~~otic~~ ganglion.

External petrosal (external superficial petrosal) passes between geniculate ganglion of seventh nerve to middle meningeal plexus.

More correctly, the great petrosal and carotid branches may be said to *form* the Vidian nerve, which runs forward through the same named canal to join the spheno-palatine ganglion, supplying it with motor and sympathetic fibres. According to this description, the twigs said to be given off by the Vidian nerve to the nasal mucous membrane must be considered branches of the *ganglion* running back in the same sheath.

Describe the cervical ganglia.

There are three on either side, viz., the

Superior cervical ganglion. It is placed opposite second and third cervical vertebræ behind carotid sheath, and gives off a

Superior branch, to internal carotid artery, forming by its division the cavernous plexuses and carotid plexus (with its subdivisions);

Descending branch, connecting superior with middle ganglion;

External branches to cranial and spinal nerves, and *anterior branches* forming plexuses on external carotid and its branches;

Pharyngeal, forming with branches from pneumogastric, glossopharyngeal, and external laryngeal nerves, the ~~*pharyngeal plexus*;~~

Laryngeal, uniting with superior laryngeal nerve and its branches;

~~*Superior cardiac,*~~ connected with other branches of sympathetic, and with some of pneumogastric, passes to back of aorta, the right joining the *deep*, and the left (usually) the *superficial* ~~cardiac plexus~~.

Middle cervical ganglion is placed opposite fifth cervical vertebra; by its

Superior and inferior branches it communicates respectively with superior and inferior cervical ganglia; the *external* filaments join fifth and sixth spinal nerves, the internal are the *thyroid* to inferior thyroid artery and gland, and the *middle* or ~~*great cardiac*~~ *nerve*, communicating with other sympathetic branches and recurrent laryngeal, to terminate in the deep cardiac plexus.

~~*Inferior cervical*~~ *ganglion* is placed between base of transverse process of seventh cervical vertebra and neck of first rib on inner side of superior intercostal artery; its *superior* and *inferior branches* connect it respectively with middle cervical, and first thoracic ganglia; the *external* branches join the spinal nerves, others form a plexus or vertebral artery; its chief branch is the

~~*Inferior cardiac nerve*,~~ communicating with recurrent laryngeal and middle cardiac nerves, terminating in the ~~deep~~ cardiac plexus.

What is the carotid plexus?

It is a plexus situated on the outer side of the internal carotid artery communicating with the Gasserian and spheno-palatine ganglia, the ~~sixth nerve~~, and tympanic branch of glosso-pharyngeal; it supplies the carotid artery and dura mater

p. 59

What is the cavernous plexus?

It is one situated below and internal to that part of the internal carotid artery running alongside of the sella turcica, in the cavernous sinus; it communicates with third, fourth, fifth, and sixth nerves, and ophthalmic ganglion, and supplying carotid, it, with the carotid plexus, forms plexuses embracing the cerebral and ophthalmic arteries.

What are the other ganglia of the sympathetic system?

Usually twelve thoracic, four lumbar, four or five sacral, one coccygeal (ganglion impar), and numerous ganglia connected with the various viscera, whence plexuses are formed named from their locality, or the organ.

What are the branches of the thoracic portion?

Communicating, with one another and the dorsal spinous nerves, filaments to aorta and its divisions and to posterior pulmonary plexus, and

The *Great Splanchnic nerve*, formed by internal branches from thoracic ganglia between sixth and tenth, with filaments from upper six; it perforates crus of diaphragm to terminate in semilunar ganglion, sending branches to renal plexus and suprarenal capsule.

The *Lesser Splanchnic*, springing from tenth and eleventh ganglia and cord between and communicating with great splanchnic, it pierces diaphragm with preceding, to join the cœliac plexus.

Smallest, or *Renal Splanchnic*, arises from last ganglion, occasionally communicates with preceding, pierces diaphragm, and terminates in renal and lower part of cœliac plexus.

Describe the solar plexus.

It supplies all the abdominal viscera, consisting of a network of nerves and ganglia lying in front of the aorta and crura of diaphragm, surrounding the cœliac axis and root of the superior mesenteric artery, extending below to the pancreas, laterally to the suprarenal capsules. The great and part of lesser splanchnic nerves of both sides and termination of the right pneumogastric form this plexus, in conjunction with the two semilunar ganglia, these latter being large, irregular gangliform masses, composed of smaller ganglia with interspaces between, placed by side of cœliac axis and superior mesenteric artery, close to suprarenal capsules, that on the right lying beneath the vena cava. From this are derived the following plexuses accompanying the same-named arteries to the various organs:

(L. → A
[illegible]
[illegible]

Phrenic, or *diaphragmatic*,
Cœliac,
Gastric,
Hepatic,
Splenic,
Aortic,
Suprarenal,
Renal,
Superior mesenteric,
Spermatic,
Inferior mesenteric.

Name some of the more important.

Tympanic (see p. 316).

Great, or *deep*, *cardiac* is placed i~~n front of bifurcation of trac~~hea, and above that of pulmonary artery, b~~ehind aortic arch~~, and is formed by all sympathetic cardiac nerves (except left superior), and cardiac branches of recurrent laryngeal and pneumogastric (except left superior cardiac and left inferior cardiac of pneumogastric).

G. of
Wrisb

Superficial cardiac lies ~~beneath aortic arch~~ in ~~front of right pul~~

monary artery, and is formed by left superior cardiac, left inferior cardiac of pneumogastric, and branches from deep cardiac plexus.

Coronary, the *posterior*, chiefly from deep, the *anterior* from superficial cardiac plexus. accompanying R. & L. Coron. art.

Aortic, on sides and front of vessel between roots of superior and inferior mesenteric arteries.

Hypogastric, lying in front of sacrum, between common iliac arteries, it supplies the pelvic viscera, and is formed by filaments from aortic plexus, and from lumbar and first two sacral ganglia, contains no ganglia, and divides into two lateral portions, forming the inferior hypogastric or ~~pelvic~~ plexuses.

Inferior hypogastric plexuses. These lie upon each ~~side of rectum and bladder~~ (rectum, vagina, and bladder, in females), and each is formed by a continuation of the hypogastric plexus and branches from second, third, and fourth sacral nerves, and a few filaments from sacral ganglia; the branches accompany those of internal iliac artery, and are distributed to all the pelvic viscera; their branches are

Inferior hemorrhoidal plexus.

Vesical plexus.	*Vaginal plexus.*
Prostatic plexus } *Small and large cavernous nerves* to penis.	*Uterine nerves.*

VISCERAL ANATOMY.

The Digestive Organs.

What is a viscus?

Any of the internal organs with their appendages, contained within the cranial, thoracic, or abdominal cavities.

Of what does the digestive apparatus consist?

Of the alimentary canal, a musculo-membranous tube, lined with mucous membrane, about thirty feet long, and extending from the mouth to the anus, and certain accessory organs.

Name the subdivisions of the alimentary canal.

Mouth, pharynx, œsophagus, stomach, small intestine (duodenum, jejunum, ilium), large intestine (cæcum, colon, rectum).

What are the accessory organs?

The teeth, salivary glands (parotid, submaxillary, sublingual), liver, pancreas, and spleen.

The Teeth.

What is their function?

To reduce the food to fragments, thus enabling the digestive fluids to act to best advantage.

Into what classes are they divided?

Into

Temporary, or *milk teeth*, ten in each jaw, viz., four incisors, two canines, and four molars;

Permanent teeth, sixteen in each jaw, viz., four incisors, two canines, four bicuspids or premolars, six molars.

Of what parts does every tooth consist?

Of a

Crown or *body* (Fig. 82), that part projecting above the gum.

Neck, the constricted portion between crown and fang.

[4]*Fang*, or *root*, that part occupying the alveolus, held there by the periodontium (periosteum), lining the socket; in addition, the teeth are steadied by the *gums*, composed of dense fibrous tissue covered with mucous membrane.

[3]*Pulp-cavity*, an interior cavity filled with the tooth-pulp, a vascular connective tissue with numerous nerves, both arteries and nerves reaching the pulp by a canal opening at apex of fang.

Describe the characteristics of each of the four varieties of teeth.

Incisor or *cutting teeth*. The crown is wedge-shaped, convex in front, bevelled and slightly concave behind; the fang is single, long, conical, and transversely flattened.

Canines. Crown is large, ~~conical~~, convex in front, ~~rises above level of other tee~~th; fang long, conical, compressed laterally; upper pair are called in common parlance "~~eye-teet~~h," the lower "s~~tomach-tee~~th."

Bicuspids. Crown has t~~wo-projecting cus~~ps, fang generally is ~~single,~~ laterally grooved with ~~bifid ape~~x; they are also called *premolars.*

Molar (grinders). Crown nearly ~~cubical~~, with four cusps in upper, and ~~five in lower~~ molars; ~~fangs~~, usually three for first two ~~upper~~, and ~~two f~~or first two ~~lower~~ molars; the third molar is called the "wisdom tooth" (*dens sapientiæ*), from its late appearance, and usually has but one fang with grooves indicating a tendency to formation of three fangs in upper, two in lower jaw.

The *second temporary molar* is the largest milk tooth, and is succeeded by the ~~second~~ permanent bicuspid.

FIG. 82.



Describe the structure of a tooth.

Each is formed chiefly of

Dentine, or *ivory*², composed of minute, wavy, branching tubes, called *dental tubuli,* embedded in a hard, homogeneous substance, the *intertubular tissue.* The tubules are about $\frac{1}{4500}$th of an inch in diameter, dividing dichotomously, giving the wavy appearance of the cut surface, and open into the pulp cavity. Chemically dentine consists of twenty-eight parts of animal, and seventy-two of earthy matter.

*Enamel*¹ forms a thin crust over crown, is the densest of all animal tissues, containing only 3.5 per cent. of animal matter, and is composed of minute parallel hexagonal rods, about $\frac{1}{5500}$th of an inch in diameter, pursuing a wavy course.

Cement, or *crusta petrosa*⁴, is a layer of true bone commencing at the neck and becoming thicker toward apex of fang.

Pulp, filling the pulp-cavity³, consisting of soft, very vascular connective tissue, with numerous nerves and cells, the latter being of two kinds, columnar, called *odontoblasts*—arranged in a layer lining pulp-cavity—and fusiform cells wedged in between these or permeating the pulp, both having fine processes, said to be prolonged into the dentinal tubules.

Whence do the teeth obtain their blood and nerve supply?

From the alveolar and infraorbital branches of internal maxillary, and from inferior dental artery; the nerves come from the anterior and posterior dental branches of the superior maxillary, and from the inferior maxillary division of the fifth nerve

When do the temporary teeth appear?

The time is variable, but, according to the latest authority, they crupt as follows, expressed in *months*:

5 3 4 2 1 1 2 4 3

	Molar.	Molar.	Canine.	Incisor.	Incisor	Incisor.	Incisor.	Canine.	Molar,	Molar.
Upper. . . .	20–30	12–15	18–24	8–10	8–10	8–10	8–10	18–24	12–15	20–30
Lower . . .	20–30	12–15	18–24	12–15	4–7	4–7	12–15	18–24	12–15	20–30

36 18 24 12 6

The order of their appearance is first, lower central incisors; then upper central incisors, closely followed by laterals; then lower laterals; next upper anterior molars followed by lower; then upper canines followed by lower; finally, lower back molars, followed by upper.

Describe the order of appearance of the permanent teeth.

The first to appear are first molars at end of sixth year, the lower teeth usually preceding the upper; thus, expressed in *years*, these teeth erupt as follows:

8, 7 1 5 4 6 3 2 2 3 1

	Wisdom.	Molar.	Molar.	Bicuspid.	Bicuspid.	Canine.	Incisor.	Incisor.	Incisor.	Incisor.	Canine.	Bicuspid	Bicuspid.	Molar.	Molar.	Wisdom
Upper	17–21	12–13	6½	10	9	11–12	8	7	7	8	11–12	9	10	6½	12–13	17–21
Lower	17–21	12–13	6½	10	9	11–12	8	7	7	8	11–12	9	10	6½	12–13	17–21

12, 6, 10, 9, 11 8, 7,

Describe the development of the teeth.

About the seventh fœtal week appears a depression in each jaw, the *primitive dental groove,* from the bottom of which is developed a ridge which, atrophying at intervals, leaves a row of projecting ~~papillæ~~, containing ~~*odontoblasts*~~, which become ~~capped with epithelial~~ masses developed from the margins and sides of the groove, the ~~*enamel organ*~~. Now the groove becomes converted into *follicles,* each containing a papilla, by contraction and projection of its margins and the ingrowth of membranous septæ.

About the thirteenth week the papillæ grow rapidly, the follicles deepen and become closed in by the coalescence of from two to five small membranous outgrowths springing from their margins, called *opercula.* The lips of the dental groove advance and fuse, completing the *saccular stage* about the end of the fifteenth week. The more superficial portion of the groove remains open, leaving the *secondary dental groove,* in which ten lunated depressions appear behind each of the sacs of the milk teeth, for the ten anterior permanent teeth; the secondary groove closes in like the primary one, the follicles becoming *cavities of reserve,* into which papillæ grow to be closed in as for the temporary teeth; the remaining teeth arise from successive expansions backward of the primitive dental groove. The dentine forms from without inward capped with enamel, and by the gradual growth of the fang the crown produces absorption by pressure upon the overlying bone and mucous membrane, when *eruption* occurs; the *cementum* is formed by the periodontal membrane last of all, increasing up to old age.

The Mouth.

What is the mouth?

It is an ovoid cavity in which food is masticated, bounded by lips in front, by cheeks and alveolar processes of both jaws with their contained teeth at sides, by hard and soft palate above, by tongue and floor of mouth below, and terminates at the anterior pillars of the fauces where it opens into pharynx by fauces, and is lined by mucous membrane covered by ~~scaly stratified epithelium~~ containing numerous ~~racemose glands~~, continuous with external skin; it presents for examination

The hard palate, formed by palatal processes of superior maxillary and palate bones covered by the intimately adherent periosteum and mucous membrane.

Soft palate, consisting of a ~~fold~~ of mucous membrane depending from posterior border of hard palate, e~~nclosing muscular tis~~sue, an ~~aponeur~~osis, ~~adenoid~~ tissue, mu~~cous~~ glands, etc.; the muscles on each side being the levator and tensor palati, palato-glossus, palato-pharyngeus, and azygos uvulæ—the latter with fellow forming the median projecting conical *uvula*.

~~Anterior~~ and ~~posterior~~ pillars of the fauces, the former containing within a fold of mucous membrane the palato-glossus muscle on each side, arching downward and forward from palate to base of tongue; the latter, the ~~palato-pharyngei~~ muscles, passing backward and downward to sides of pharynx.

Isthmus faucium, bounded by pillars of fauces, base of tongue, and free margin of soft palate.

Tonsils[7] (Fig. 83), situated between anterior and posterior pillars on each side, consisting of glandular tissue containing twelve to fifteen openings leading into crypts lined with ~~mucous mem~~brane, external to which is a layer of closed capsules analogous to those of Peyer's glands; the tonsil is only separated from the i~~nternal carotid~~ artery by superior constrictor muscle.

Openings of ducts of Steno, opposite second upper molar tooth on each side, delivering the secretion of parotid glands.

Openings of ducts of Wharton, one on either side of frænum of tongue, delivering secretion of submaxillary glands.

Openings of ducts of sublingual gland (*ducts of Rivinus*), from eight to twenty in number, they open on an elevated crest of mucous membrane on each side of frænum linguæ, one or more joining in a tube ope~~ning~~ into ~~Wharton~~'s duct called the *duct of Bartholine*.

What are the salivary glands?

Three compound racemose glands on each side. The *parotid*, the largest, lies below and in front of ear between zygoma above, mastoid behind, and ramus of jaw in front—its duct is *Stenson's*; the *submaxillary* lies below jaw in anterior part of submaxillary triangle—its duct is *Wharton's*; the *sublingual* lies beneath mucous membrane of floor of mouth against jaw, close to symphysis—its ducts are those of *Rivinus*.

Describe the tongue.

Its base is attached to hyoid bone by muscles, to ~~epiglottis~~ by the two ~~lateral~~ and one ~~median? glosso-epiglott~~idean fold of mucous membrane, and to soft palate by anterior pillars; its tip, sides, dorsum, and part of under surface are free; a median raphé

Fig 83.



and fibrous septum divide the organ into halves; its mucous membrane reflected over floor of mouth to gums forms in front an antero-posterior fold, the *frænum linguæ*; the tongue is formed by certain *intrinsic muscular fibres*, viz., those of lingualis, and by *extrinsic muscles*, the stylo-glossus, hyo-glossus, genio-hyo-glossus, and palato-glossus (see p. 121). The mucous membrane of tongue presents the

Circumvallate papillæ, numbering eight to ten, arranged on back part of dorsum in two lines converging behind like a letter V, presenting at their junction a deep ~~mucous follicle, the *foramen cæcum*~~[2].

Fungiform papillæ[3], scattered over dorsum, but chiefly found at sides and apex.

Filiform papillæ[4], with secondary papillæ projecting from apices, arranged in lines, cover the anterior two-thirds of dorsum of tongue.

Racemose glands[a], situated along sides beneath tip, but chiefly over posterior third of dorsum. = Pharyngeal

Lymphoid tissue, collected into masses or *follicles*, some resembling crypts of tonsil.

Describe the blood supply.

This is from,

Lingual { *Dorsalis linguæ*, *Sublingual*, *Ranine*,

Ascending pharyngeal.

Facial, by submental branch anastomosing with sublingual.

Describe the nerve supply.

The ~~gustatory~~ branch of ~~fifth~~ is distributed to ~~papillæ~~ Tactile at front and sides, which endows these parts with ~~general sensibility~~; and also by its connection with the ~~*chorda tympani*~~ becomes the nerve of ~~taste~~.

Lingual branch of the glosso-pharyngeal, supplying mucous membrane of base and sides and ~~circumvallate papillæ~~; it is the nerve of taste for these parts.

Hypoglossal, to muscles; the motor nerve.

Superior laryngeal, sends a few filaments to base from its internal branch.

What two anatomical points of surgical interest does the palate present in the neighborhood of the last molar tooth?

Just behind this tooth the *hamular process and internal pterygoid plate* can be felt, the point for division of tensor palati aponeurosis in the operation for cleft palate, and in front of this to inner side of last molar tooth is the *posterior palatine artery* as it emerges from the canal, sometimes requiring plugging after a cleft-palate operation.

The Pharynx.

What is the pharynx?

A conical ~~musculo-membranous~~ sac, about ~~four and a half inches~~ long, extending from under surface of base of skull to a point corresponding to the ~~cricoid cartilage~~ in front, and the ~~fifth cervical~~ vertebra behind. It is widest opposite cornua of hyoid bone, narrowest below, where it terminates in the ~~œsophagus~~; it is bounded above by basilar process of occiput, is connected *posteriorly* with the cervical vertebræ and longus colli and recti capitis antici muscles; *anteriorly* it is incomplete, opening into the mouth, being attached to internal pterygoid process, ~~pterygo-maxillary ligament~~, lower jaw, tongue, hyoid bone and larynx; *laterally*, it is attached to styloid processes and muscles arising from them, while the common and internal carotid arteries, the internal jugular veins, and eighth, ninth, and sympathetic nerves here lie in contact with it. ?

How many openings communicate with it?

Seven: the two posterior nares, two Eustachian tubes, mouth, larynx, and œsophagus.

Of how many coats does it consist?

Three: an internal ~~mucous~~, continuous with lining of mouth, etc., covered with ciliated epithelium down to the level of floor of nares, below by squamous epithelium, containing racemose glands most plentiful around orifices of Eustachian tubes, and much lymphoid tissue surrounding crypts like those of tonsil; a middle

~~fibrous coa~~t (*pharyngeal aponeurosis*), thick above, where the muscular fibres are wanting; and an ~~external muscular coat~~ composed of the superior, middle, and inferior constrictors, stylo-pharyngeus, and palato-pharyngeus muscles.

Describe the pharyngeal muscles.

***Inferior constrictor*[8]**: *origin*, side of cricoid and thyroid cartilages; *insertion*, posterior median raphé; *action*, constricts pharynx during swallowing; *nerves*, pharyngeal plexus, glosso-pharyngeal, external laryngeal, recurrent laryngeal.

***Middle constrictor*[9]**: *origin*, greater and lesser cornua of hyoid bone and stylo-hyoid ligament; *insertion*, posterior median raphé; *action*, same as superior constrictor; *nerves*, glosso-pharyngeal, pharyngeal plexus.

***Superior constrictor*[10]**: *origin*, lower third of margin of internal pterygoid plate and hamular process, contiguous surface of palate bone, reflected tendon of tensor palati muscle, pterygo-maxillary ligament, alveolar process above posterior end of mylo-hyoid ridge and side of tongue; *insertion*, posterior median raphé and pharyngeal spine of occipital bone; *action*, same as other constrictors; *nerves*, glosso-pharyngeal plexus.

FIG. 84.



***Stylo-pharyngeus*[6]**: *origin*, inner side of base of styloid process; *insertion*, blends with constrictor muscles, and is also inserted into the posterior border of thyroid cartilage; *action*, draws sides of pharynx upward and outward; *nerves*, glosso-pharyngeal, pharyngeal plexus, the former running on outer side, and crossing muscle to reach tongue.

What is the pharyngeal tonsil?

A considerable mass of lymphoid tissue extending across back of pharynx, between the orifices of the Eustachian tubes.

What arteries supply the pharynx?

Superior thyroid,	*Pterygo-palatine,*
Ascending pharyngeal,	*Descending palatine,* to *epiglottis,* etc.

What are its nerves?

Branches from pharyngeal plexus, formed by pharyngeal branches of ~~glosso-pharyngeal~~, pneumogastric, ~~superior~~ laryngeal, and sympathetic nerves.

The Œsophagus.

Describe it.

It is a muscular canal, about ~~nine inches~~ long, joining the pharynx and the stomach, commencing at lower border of cricoid cartilage, passing through the œsophageal opening of the diaphragm, to terminate at cardiac orifice of the stomach, opposite the ninth dorsal vertebra; in the *neck* it lies between trachea and spinal column and longus colli muscle, at lower part inclining to the left, having on either side the common carotid artery, with lateral lobes of thyroid gland—the ~~recurrent laryngeal~~ nerves ascend ~~between~~ it and trachea; in the *thorax,* while a little to the left at first, after passing behind aortic arch, it runs to right of the vessel, to pass in front and to the left again before piercing the diaphragm.

What coats has it?

It has three: an internal *mucous,* when empty thrown into longitudinal folds, containing numerous ~~racemose glands~~, its surface is studded with small papillæ, and it is covered by a thick layer of squamous epithelium; a middle *cellular,* loosely connecting the mucous and the succeeding coat; a *muscular,* composed of an external longitudinal and internal circular layer, the fibres *above* consisting chiefly of the striated, but *below* almost entirely of the unstriped involuntary variety.

Name its arteries and nerves.

Arteries, chiefly from thoracic aorta.

Veins, are drained by vena azygos ~~minor~~.

Nerves, from œsophageal plexus, formed by pneumogastrics with some sympathetic filaments.

The Stomach.

Describe it.

The stomach is irregularly conical, curved upon itself, presenting a round base turned to left side. Moderately distended, it measures about ~~twelve~~ inches in its longest, and ~~four inches~~ in its vertical diameter. It lies in left hypochondriac, epigastric, and part of right hypochondriac regions, immediately behind anterior wall of abdomen below the liver and diaphragm, and above the colon; it presents for examination the

Greater or *splenic end,* or *fundus,* situated to the left, being attached to the spleen by gastro-splenic omentum, and to diaphragm by gastro-phrenic ligament.

The *lesser,* or *pyloric end,* is in contact with abdominal wall and under surface of liver near end of cartilage of eighth rib.

The *œsophageal,* or *cardiac orifice,* is the highest point of the stomach, is funnel-shaped, and communicates with œsophagus.

The *pyloric orifice* communicates with duodenum, the opening being guarded by a ~~reduplication of the mucous lining~~ containing numerous circular muscular fibres, forming a ~~thick ring~~—this is called the *pyloric valve.*

The ~~*lesser curvature*~~ extends along the upper border of organ between the cardiac and pyloric orifices, is concave, and is connected with under surface of liver by the ~~lesser omentum~~.

The *greater curvature* is convex, extends along lower border between the two orifices, and gives attachment to great omentum.

How many coats has the stomach?

Four, as follows: a

Serous, derived from peritoneum covering all parts except along greater and lesser curvatures, where the omenta leave a ~~triangular space~~, along which pass the ~~vessels and nerves~~; a

Muscular, consisting of three sets of fibres: *longitudinal,* continuous with the longitudinal coat of œsophagus and small intestine, more distinct along the curvatures; the *circular fibres* form a continuous layer beneath the former set, most abundant at pyloric end, forming valve (supra); the *oblique fibres*, some passing obliquely from left to right, others *vice versâ*, around the cardiac orifice, to which part they are chiefly limited; a

Mucous, covered with columnar epithelium, and thrown into longitudinal folds or *rugæ* when stomach is empty, covered by small shallow polygonal-shaped alveoli from $\frac{1}{100}$th to $\frac{1}{350}$th of an inch in diameter, into which the gastric follicles open.

Describe the gastric glands.

They are of three kinds, the

Peptic follicles, lined throughout with columnar epithelium, consisting of a duct into which open several cæcal tubes, in which, beneath the epithelium, lie large spheroidal, coarsely granular [illegible] *peptic cells*; these glands are found in all parts of the stomach.

[illegible] *Mucous glands*, found in greatest numbers at pyloric end, and resemble the preceding, but are *without* the large peptic cells, and the cæcal tubes are longer.

Lenticular, or *simple solitary*, found especially in early life, consisting of masses of lymphoid tissue.

Give the blood and nerve supply.

The *arteries* are, *gastric*[13], *pyloric* and *right gastro-epiploic*[17] branches of the *hepatic artery*[14].

FIG. 85.



Left gastro-epiploic[21], } Branches of
Vasa brevia[23], } *Splenic artery*[20].

The *veins* empty into splenic and portal veins.

The *nerves* are terminal branches of both pneumogastrics and numerous branches from sympathetic.

The Small Intestine.

Describe it.

It is the narrowest part of the digestive tract, is a convoluted tube some ~~twenty feet~~ long, occupying the central and lower parts of the abdominal and pelvic cavities, and is ~~suspended~~ from the ~~spine by~~ a fold of peritoneum called the ~~*mesentery*~~; its three divisions are the

~~*Duodenum*~~[2] (Fig. 86), as long as the breadth of ~~twelve fingers~~, *i. e.*, eight to ten inches, passing upward and backward to the right to under surface of liver, near ~~gall-bladder~~—*ascending portion*—then descending in front of ~~right kidney~~—*descending portion*—thence running transversely across spine to end in the ~~jejunum~~ on ~~left~~ side of ~~second lumbar~~ vertebra—*transverse portion*—the junction being crossed by superior mesenteric artery; it is only partially covered by peritoneum, has no mesentery, and surrounds the head of the pancreas; into the descending portion, to the inner side, pass obliquely the common ~~bile and pancreatic ducts~~.

Fig. 86.



~~*Jejunum*~~[3], so-called because usually ~~empty after~~ ~~death~~, includes upper two-fifths of remainder of small intestine, and lies chiefly in the umbilical region.

~~Ileum~~, includes remainder (three-fifths) of small intestine, is named from its numerous ~~coils~~, occupies chiefly umbilical, hypogastric, right iliac, and occasionally pelvic regions, and opens into the inner side of the commencement of the large intestine in the right iliac fossa; the orifice is guarded by a two-leaved valve (see page 231).

What coats has the small intestine?

The same as stomach.

The *mucous membrane* is covered with columnar epithelium, and forms transverse folds encircling the tube for one-half to two-thirds its circumference, forming the

Valvulæ conniventes; they are absent in first two inches of duodenum and almost disappear in lower part of ileum; ~~villi~~ cover the surface, formed of a central ~~lacteal~~ vessel accompanied by unstriped ~~muscular~~ tissue, an encircling plexus of capillaries, lymphoid tissue and granular corpuscles, all enclosed by a basement membrane supporting columnar epithelium: their number is estimated at ~~four million~~; the

Cellular coat connects mucous with muscular coat, and contains the intestinal vessels; the

Muscular coat, externally is composed of longitudinal and internally of ~~circular~~ fibres, these ~~not forming complete~~ rings.

Describe the glands of the small intestine.

They are,

Brunner's glands, resembling pancreas in structure, and are limited to duodenum and commencement of jejunum.

Simple follicles or *glands of Lieberkühn*, are minute tubular depressions lined with columnar epithelium.

Solitary glands, most numerous in lower part of ileum, but found in all parts, and consist of masses of lymphoid tissue about half a line to a line in diameter.

Peyer's glands or *patches* are twenty to thirty ovoidal patches composed of numerous solitary glands (hence named *agminated glands*), situated opposite to mesenteric attachment, their long axis lengthwise, and are largest and most numerous in ileum, although occasionally seen in duodenum; the valvulæ conniventes cease at their margins.

[illegible] Lieberkühn (g [illegible]
[illegible] (Duodenal [illegible]
[illegible]

Name bloodvessels and nerves.

The *arteries* are from gastro-duodenal of hepatic and superior mesenteric arteries; the *veins* chiefly empty into portal vein; *nerves* are, from superior mesenteric plexus of sympathetic.

The Large Intestine.

Describe it[5],[7],[8],[9].

It extends from ileum to anus, is about ~~five feet long~~, is large and sacculated, has the same coats as small intestine, the mucus ~~destitute of~~ villi, but thrown into crescentic folds by the ~~longi~~tudinal ~~muscular~~ fibres which are disposed in ~~three bands~~, shorter than the other coats, so forming them into pouches.

Describe the subdivisions of the colon.

They are,

The *cæcum*[5] or *caput coli*, the dilated commencement of the large bowel situated in the right iliac fossa, and usually wholly covered by peritoneum.

The *appendix vermiformis*[6], a narrow, blind-ended, worm-like tube, from three to six inches long, springing from lower back part of cæcum, held, coiled upon itself, by a peritoneal fold.

The *ileo-cæcal valve* (or Bauhin's) is formed by ~~two horizontal semilunar folds~~ of mucous membrane at the termination of ileum in cæcum, opening toward large intestine and ~~guarding against reflux *from large*~~ *into* small bowel; the mucous folds are reinforced by circular muscular fibres, and the surfaces toward the ileum are covered with villi, while these are absent on the cæcal side.

The *ascending colon*[7] extends up from cæcum in front of right kidney, to under surface of liver to right of gall-bladder, where abruptly bending to the left, it forms the ~~*hepatic flexure*~~; it is almost—sometimes entirely—enveloped in peritoneum, in which latter event it has an *ascending meso-colon.*

The *transverse colon*[8] traverses the abdomen from right to left, just ~~below~~ liver, stomach, and ~~spleen~~, to left hypochondriac region, where it curves downward beneath lower end of spleen, forming the ~~*splenic flexure*~~; it has a wide transverse *meso-colon*, attaching it to the spine.

The *descending colon*[9] passes down in front of left kidney to left iliac fossa and is only ~~partially covered with periton~~eum, its ~~pos~~terior surface usually being ~~free.~~

The *sigmoid flexure*[10], the narrowest part, lies in left iliac fossa extending from descending colon at crest of ilium to left sacro-iliac symphysis; it is curved like an S, and is held in place by a loose peritoneal fold, the *sigmoid meso-colon.*

The ~~rectum~~[11] extends from left sacro-iliac symphysis tc anus, is from ~~six to eight inches~~ long, curves slightly to right and then adapts itself to sacral curve, and ~~near top of coccyx~~ inclines ~~back~~ward to anus; it is non-sacculated, but just above anus presents a considerable dilatation; it is o~~nly covered with~~ peritoneum at its anterior portion—except above—while its lower inch and a half is destitute of any such covering; the longitudinal muscular fibres form a continuous layer around the rectum.

What are the appendices epiploicæ?

They are little ~~peritoneal pouches filled with~~ fat placed along ~~colon~~ and upper part of ~~rectum~~.

What guards the anal orifice?

The sphincter ani and internal sphincter; a third has been described as some t~~hree~~ inches up the bowel (see page 331

What are the folds of Houston?

Three semilunar folds of mucous membrane, one at upper right side of rectum, one at left middle portion, and one projecting from front of rectum opposite bladder.

Name the glands of the large intestine.

Follicles of Lieberkühn and *solitary glands.*

Give the blood- and nerve-supply of the large intestine and rectum.

The *arteries* are branches of the *superior* and *inferior mesenteric,* the *middle hemorrhoidal* from the internal iliac, and the *inferior hemorrhoidal* from internal pudic.

The *nerves* come from sympathetic plexuses around mesenteric arteries, and, in case of rectum, the fourth anterior sacral nerve supplies the sphincter muscle.

) Diagram as tho looking at own liver from behind & below.

The Liver.

Describe it.

It occupies the right hypochondriac region extending across the epigastrium into the left hypochondriac. It is the largest gland in the body, weighing from t~~hree to four pou~~nds, its transverse diameter is from ten to twelve inches, its antero-posterior six to seven, its thickest part behind three inches. Covered by peritoneum, e~~xcept at~~ attachments ~~of~~ ligaments and ~~bottom~~ of ~~fissures~~, it possesses a *fibrous coat,* covering the whole organ continuous with the *capsule* of *Glisson* at transverse fissure; there are five fissures, five lobes, five ligaments, and five sets of vessels; the

Upper surface is convex, smooth, lying in contact with diaphragm above, and below with a small portion of abdominal wall; it is divided into two unequal lobes by a peritoneal fold, the suspensory ligament; the

Under surface is concave, is in relation with stomach, duodenum,

FIG. 87.



hepatic flexure of colon, right kidney, and suprarenal capsule, and is divided by a *longitudinal fissure* [8,9], into a right and left lobe; the

Anterior border[4] is thin, sharp, and deeply notched at site of round ligament; it corresponds usually with margins of ribs in males, projecting a little lower in women and children; the

Posterior border[3] is rounded and grooved, or even perforated, by inferior vena cava.

Describe the ligaments.

They are five in number, four being composed of folds of peritoneum, the fifth is the obliterated umbilical vein; the

Suspensory or broad ligament[18] (falciform) is a falciform antero-posterior peritoneal fold, with ~~apex backward~~, attached *to liver* from notch on anterior border to posterior border,—its anterior free edge enveloping the round ligament—and to *diaphragm* and sheath of right rectus muscle as low as umbilicus; the

Coronary ligaments consist of two layers reflected from diaphragm on upper and lower margins of posterior border of organ, ~~between~~ which firm areolar tissue bi~~nds the~~ liver to diaphragm; the

Lateral ligaments, right and left, ~~triangular~~, are formed of two layers of peritoneum extending from sides of diaphragm to adjacent margins of posterior border of liver; the

Round ligament[17], a fibrous cord, ascends from umbilicus in anterior free margin of broad ligament to longitudinal fissure, traceable back to vena cava, the back portion, the remains of ductus venosus, the anterior of umbilical vein.

Describe the fissures.

They are five in number, dividing the under surface of the organ into five lobes; the

Longitudinal fissure[8],[9], extends from notch on anterior border to posterior border, separating right and left lobes, and is joined by the *transverse fissure*[10] at posterior third; the anterior two-thirds of the longitudinal fissure is called the *umbilical fissure* from lodging fœtal umbilical vein, the posterior third the *fissure of the ductus venosus* lodging the obliterated fœtal vessel; the anterior part of the former is often partially bridged over by the *pons hepaticus*; the

Transverse fissure[10] is about two inches long, passing transversely across under surface of right lobe from junction of posterior and middle thirds of longitudinal fissure; here enter the portal vein, hepatic artery, and nerves, and the bile ducts and lymphatics emerge; the

Fissure of the gall-bladder[14] lies parallel and to right of longitudinal fissure reaching back nearly to transverse fissure; the

Fissure for vena cava is short and deep, sometimes converted into a canal, extending obliquely upward from a little behind right extremity of transverse fissure to posterior border of liver, there joining the fissure of ductus venosus; the hepatic veins here empty into vena cava.

Describe the lobes of the liver.

They are five in number; the

Right lobe[1], the largest, of a quadrilateral form, its under surface marked by three fissures: viz., transverse, for gall-bladder, and for inferior cava, and by an anterior depression for colon, a posterior one for right kidney; the longitudinal fissure forms the division between this and the

Left lobe[2], smaller and flattened, resting by its anterior concave surface on stomach; the

Lobus quadratus[5] is marked off from under surface of right lobe by transverse fissure behind, by that for the gall bladder on the right, and anterior part of longitudinal fissure on left; the

Lobus Spigelii[6] projects from back under surface of right lobe, bounded in front by transverse fissure, on right by that for cava, and on left by posterior third of longitudinal fissure; the

Lobus caudatus[7] is a small elevation running obliquely outward from base of lobus Spigelii to under surface of right lobe; it intervenes between right end of transverse fissure and commencement of that for vena cava.

Describe the vessels of the liver.

They number five, viz., hepatic artery, portal vein, hepatic vein, hepatic duct, and the lymphatics; the first three are enveloped in loose areolar tissue called *capsule of Glisson*, which accompanies the vessels through the *portal canals* in the interior of the organ; the

Portal vein[11], about four inches long, is formed by the junction of superior mesenteric and splenic veins, the latter receiving the blood returned by inferior mesenteric vein, while the gastric veins empty into the portal vein itself. Entering the transverse fissure it divides into branches for right and left lobe, which receive the *vaginal* and *capsular* veins, and subdividing form *interlobular veins*[b] (Fig. 88), which send branches to interior of each lobule forming a plexus converging to empty into the central *intra-lobular vein*, running at right angles to other vessels; the intra-lobular vein empties into a *sublobular vein*, a radicle of hepatic vein; the

Hepatic veins[16] (Fig. 87) commence in sublobular veins as just described, which unite to form three large and several smaller branches terminating in inferior vena cava in fissure for that vessel; the

FIG. 88.



Hepatic artery[12] springs from cœliac axis, enters transverse fissure, and divides into branches which ramify in the capsule of

Glisson—*vaginal,* supplying nutriment to vessels, ducts, etc., *capsular* to terminate in external fibrous coat—and ends in a plexus on outer surface of each lobule formed by ultimate branches of portal vein, the *interlobular plexus* above described; the

Hepatic duct above 13 is formed by the union of two trunks of nearly equal size which emerge from the transverse fissure, one from the right and other from left lobe; it passes downward to the right for one inch and a half to be joined by the *cystic duct* from the gall-bladder, to form the ductus communis choledochus (see p. 238); the

Lymphatics accompany bloodvessels, and consist of a deep and superficial set.

Whence do the nerves of the liver come?

From the hepatic plexus of the sympathetic, from the pneumo~~gastric nerves~~, especially the left, and right ~~phrenic~~.

Describe the liver structure.

It is composed of numerous ~~lobules~~ from one-twentieth to one-~~tenth~~ of an ~~inch in diameter~~, of irregular form in human liver, polygonal in pigs, their bases clustered round the sublobular vein, to which each is connected by its intralobular vein; the lobules are bound together by loose areolar tissue; each lobule is formed of *hepatic cells,* spheroidal or many-sided nucleolo-nucleated cells, measuring from one-one thousandth to one-two thousandths of an inch, containing granular yellow coloring matter and globules, which cells lie in the meshes of the dense capillary network occupying its interior, spaces between the cells forming *intercellular biliary passages*^{c}, the radicles of the bile ducts.

What is the capsule of Glisson?

The areolar tissue entering the transverse fissure, surrounding the hepatic vessels to their ultimate ramifications, and continuous with fibrous coat of organ.

How may the portal be distinguished from the hepatic veins in a liver section?

The portal veins are collapsed, being surrounded by the capsule of Glisson, while the hepatic are patulous, being closely connected with liver substance.

Describe the gall-bladder[14].

It is a pear-shaped, fibro-muscular receptacle for the bile; its fundus and under surface of body and neck covered by peritoneum, while its interior is lined with a cylindrical-celled, epithelial-coated mucous membrane, presenting a honeycombed appearance from the confluence of numerous minute rugæ; its length is from three to four inches, its capacity from eight to twelve fluidrachms, and it lies in a fissure on the under surface of liver, its *fundus*—most dilated portion—projecting slightly beyond the anterior border, touching abdominal wall just below ninth costal cartilage; the narrow *neck* curves upon itself like an italic *f*, to terminate in the *cystic duct*, which is about one inch long, and is lined by mucous membrane so disposed in a series of crescentic folds as to present the appearance of a *spiral valve*; it joins the hepatic duct to form the common bile-duct.

Describe the ductus communis choledochus[13] (Fig. 87).

It is about ~~three inches~~ long, three-sixteenths of an inch in diameter, and is formed by the junction of the hepatic and cystic ducts; it opens usually by a common orifice with the pancreatic duct, a little below the middle of inner side of descending portion of duodenum; its walls are composed of fibrous with a little muscular tissue, its mucous membrane is covered with columnar epithelium, and it ~~runs between~~ the layers of gastro-hepatic process of the peritoneum.

The Pancreas.

Describe the pancreas[9] (Fig. 85).

It is a compound racemose gland, from six to eight inches long, by one and one-half inches at its widest part, lying behind the stomach and in front of the first lumbar vertebra. It is composed of lobules connected by areolar tissue, each lobule consisting of an ultimate branch of the duct lined with columnar epithelium, terminating in cæcal pouches or acini, also lined with cylindrical epithelium, outside of which is a fine capillary network. The *Pancreatic duct* (*canal of Wirsung*) extends the whole length of gland and opens into middle of descending part of duodenum, to the inner side, with common bile-duct; its

Head or *right extremity* is embraced by the concavity of the duodenum, the common bile-duct lying behind, the pancreatico-duodenal artery in front; the

Tail or *left extremity* reaches spleen above left kidney and suprarenal body; the

Body is covered by the ascending layer of transverse meso-colon and posterior surface of stomach; posteriorly, the superior mesenteric artery and vein, the portal vein, the vena cava, and the aorta separate it from the first lumbar vertebra.

The *arteries*[18] (Fig. 85) are derived from the splenic, pancreatico duodenal branch of hepatic and superior mesenteric arteries.

The *veins* empty into splenic and superior mesenteric veins.

The *nerves* come from the splenic plexus of the sympathetic.

What is the lesser pancreas?

A lobular fold, passing transversely to the left behind the superior mesenteric vessels, which is sometimes detached from the gland when it may open into duodenum by a separate duct an inch or more above the pancreatic duct.

The Ductless Glands.

What are the ductless glands?

The spleen, suprarenal capsules, thyroid and thymus glands, bodies whose functions are uncertain and which have no ducts.

The spleen and suprarenal capsules lie in the abdominal cavity, but the others will be for convenience described after them.

The Spleen[10] (Fig. 85).

Describe it.

It is a soft, brittle, very vascular, oblong, flattened organ embracing the cardiac end of the stomach, to which it is attached by the gastro-splenic fold of peritoneum, which latter membrane completely invests the spleen, except where the gastro-splenic omentum and suspensory ligament are attached; the organ lies deeply in the left hypochondriac region; the

Outer surface is convex, smooth, lying in contact with ~~diaphrag~~m, which ~~separates~~ it from the ~~ninth, tenth, and~~ eleventh ribs; the

Inner surface is concave, lies in contact with stomach, and presents the

Hilum, a vertical fissure pierced at various points by bloodvessels, lymphatics, and nerves; the

Anterior border is often notched, and the

Posterior border is rounded and in relation with left kidney; the

~~*Suspensory ligament*~~, a peritoneal fold, connects it with the under surface of diaphragm; the

~~*Fibro-elastic coat*~~, beneath the serous (see above) which intimately adheres to it, forms the framework of the spleen, passing ~~inward at the hilu~~m with th~~e vessels as sheaths,~~ from which, and the inner surface of capsule, numerous bands or ~~*trabeculæ*~~ pass, uniting to form numerous small spaces o~~r *areolæ*~~ filled with the splenic pulp.

Describe the spleen-pulp.

It is of a dark reddish-brown color, and under the microscope is seen to consist of a network of branched connective-tissue cells, containing pigment granules (broken-down blood corpuscles), granular matter, nucleated cells, free nuclei, and red blood-cells in all stages of disintegration; the

Splenic artery[20] (Fig. 85) is large, tortuous, and divides at hilum into five or six branches which do not anastomose to any extent, each branch breaking up again and again into smaller ones, until they terminate in bundles of straight vessels which open into the interstices of the reticulum formed by the branched connective-tissue cells.

What are the Malpighian bodies?

Small, rounded, expanded masses of the lymphoid tissue of the external coat of the small ~~splenic ar~~teries, from one-one hundredth to one twenty-fifth of an inch in diameter, usually surrounding the vessel, but sometimes projecting from one side; they are composed of a delicate reticulum packed with lymph corpuscles and pervaded by capillaries; the

Splenic vein arises from the *lacunar* spaces in the pulp, and empties into the portal vein; the

Nerves are derived from the right pneumogastric nerve and the right and left semilunar ganglia.

The Thyroid Gland.

Describe it.

It consists of ~~two lateral lo~~bes situated at the sides of the upper portion of the trachea, connected about their lower thirds by a narrow transverse portion, the ~~*isthmus*~~, which is occasionally absent. A third lobe, the ~~*pyramid*~~, sometimes springs from the Rem. of left upper side of isthmus, or the left lobe, and extends to hyoid bone; it is at times detached; occasionally a slender muscular band—the *levator glandulæ thyroideæ*—extends from hyoid bone to isthmus or the pyramid. Structurally, the gland consists of a connective-tissue capsule, sending in septa subdividing the organ into lobes, each of which is formed of numerous closed vesicles filled with a viscid yellowish fluid embedded in connective tissue containing a dense vascular plexus surrounding the vesicles.

What arteries supply this gland?

The *superior* and *inferior thyroid* branches respectively of external carotid and thyroid axis, and at times the *middle thyroid*, or *thyroideæ ima*, springing either from the innominate artery or aortic arch, itself to pass up on the front of the trachea. These vessels are remarkable for their size and frequent anastomoses. (For description of those vessels, see pages 159 and 161.)

Describe the veins of the thyroid gland

They form a plexus on the surface of the gland, whence arise the

Superior thyroid vein, } Emptying into internal jugular vein.
Middle thyroid vein, }
Inferior thyroid vein opening into innominate vein.

Name the nerves.

Branches of the pneumogastric nerves and of middle and inferior cervical ganglia.

The Thymus Gland.

Describe it.

This is a temporary organ of ~~unknown function~~, reaching its full development at the end of the second year, after which it gradually atrophies, having nearly disappeared at puberty. When fully developed it extends from fourth costal cartilage to lower border of thyroid gland, consisting of two lateral lobes in close contact along the middle line, while sometimes an intermediate lobe exists; the sternum and origins of the ~~sterno-hyoid and sterno-~~thyroid muscles ~~cover it~~, as it lies in the neck behind these muscles upon front and sides of trachea; it rests, in the mediastinum, upon the pericardium, the ~~thoracic fascia~~ separating it from the aortic ~~arch and~~ great vessels. Its dimensions are: length, ~~two inches~~, breadth, one ~~and a~~ half inches, thickness, ~~three to~~ four lines; at birth it weighs about half an ounce.

Describe its structure.

It is formed of numerous ~~lobules~~ bound together by delicate areolar tissue, all being inclosed by a fibrous capsule; the lobules are composed of a ~~*cortical* lymphoid~~ tissue, and a *medullary* portion containing a few lymphoid cells, but toward the centre granular cells and ~~concentric corpuscles~~ exist. Each follicle is surrounded with a capillary plexus.

Describe the arterial and nerve supply.

The *arteries* are derived from internal mammary, and superior and inferior thyroid.

The *veins* empty into thyroid and left innominate veins.

The *lymphatics* arise in the substance of the gland, and are said to empty into internal jugular vein.

The *nerves* are very minute, coming from pneumogastric and sympathetic.

The Suprarenal Capsules.

What are they?

Two triangular or semilunar flattened bodies, situated behind peritoneum, immediately in front of upper part of each kidney.

They consist of a thin, closely adherent *fibrous capsule,* which sends numerous communicating partitions inward, a *~~cortical~~* and *~~medullary~~ portion,* the former containing groups of finely granular polyhedral cells, with a channel between them and the fibrous portions, believed to be a lymph sinus; the *medullary portion* is formed of stout connective-tissue bundles, between which lie coarsely granular columnar and branched cells, which have been thought to communicate with the nerve fibres of the very fine plexus found in the medulla.

What parts do these organs lie in contact with ?

The lower concave border rests upon upper end of kidney; their anterior surfaces touch respectively the under surface of liver, pancreas, and spleen; their posterior surfaces lie upon ~~crura of~~ diaphragm, opposite ~~tenth dorsal~~ vertebra, and ~~their inner~~ margin is related to the ~~great splanchnic~~ nerves and ~~semilunar ganglia~~, on the right side touching ~~vena cava~~, on left ~~aorta~~.

Name its vessels and nerves.

The *arteries* are the suprarenal and branches of inferior phrenic and renal arteries forming a dense capillary network in its interior; the

Veins return blood from the medullary venous plexus by suprarenal veins into inferior vena cava on right side, and on left into renal vein; the

Nerves are very numerous, chiefly distributed to ~~medulla,~~ and are derived from solar and renal plexuses and phrenic and pneumogastric nerves, having developed upon them numerous small ganglia, whence the opinion held by some that these bodies belong to the sympathetic nervous system.

The Abdominal Cavity.

What is the abdomen ?

An ovoidal cavity, limited above by the thorax, below by the brim of pelvis; in front and at sides, by lower ribs, abdominal muscles, and venter ilii; behind, by spine, psoas, and quadratus lumborum muscles; it is lined with peritoneum—a serous membrane—which is reflected to form partial or complete investments for the contained organs or *viscera*.

Name the abdominal viscera.

Stomach,
Liver,
Spleen,
Kidneys,
Bladder (when full),
Uterus (during pregnancy),
Receptaculum chyli,
Intestines,
Gall-bladder,
Pancreas,
Suprarenal capsules,
Abdominal aorta,
Inferior vena cava,
Thoracic duct.

Describe the openings in its walls.

They are the

Umbilical, for *fœtal* umbilical vessels, obliterated after birth, leaving a depression, the *umbilicus.*

Opening for inferior vena cava, in diaphragm.

Aortic opening, behind diaphragm between crura, for aorta, vena azygos major, thoracic duct, and sometimes left sympathetic nerve.

Œsophageal opening, in diaphragm for œsophagus and pneumogastric nerves.

Internal abdominal ring, on each side, for spermatic cord in male, round ligament in female.

Femoral, or *crural ring,* on each side, below Poupart's ligament, for femoral vessels.

Into what regions is the abdomen artificially divided?

Into nine, by two imaginary parallel lines encircling the body on a level with ninth costal cartilages and summits of iliac crests, and two other parallel lines drawn from the cartilage of the eighth rib on each side to the centre of Poupart's ligament.

What are these regions named, and what does each contain?

The right lateral are, from above downward (see cut), the

Right hypochondriac, which contains right lobe of liver, gall-bladder, hepatic flexure of colon, upper part of right kidney, and part of right suprarenal capsule.

Right lumbar, contains ascending colon, lower part of right kidney, and some coils of small intestine.

FIG. 89.



Right inguinal (*iliac*), contains cæcum and appendix vermiformis, ureter, spermatic vessels.

The median regions are, from above downward (see cut); the

Epigastric, contains middle and pyloric end of stomach, left lobe of liver, lobulus Spigelii, pancreas, duodenum, parts of kidneys and suprarenal capsules, aorta and branches, vena cava, semilunar ganglia, thoracic duct.

Umbilical, contains transverse colon, part of great omentum and mesentery, transverse portion of duodenum, and some coils of jejunum and ileum, part of both kidneys, and the receptaculum chyli.

Hypogastric, contains coils of small intestine, the bladder in children, or when distended in adults, and the uterus during latter six months of pregnancy, often cæcum, appendix vermiformis, and sigmoid flexure of colon.

Left lateral regions from above downward (see cut) are,

Left hypochondriac, contains splenic end of stomach, spleen, and tail of pancreas, splenic flexure of colon, upper half of left kidney, and part of left suprarenal capsule.

Left lumbar, contains descending colon, part of omentum, lower part of left kidney, and some coils of small intestine.

Left inguinal (iliac), contains sigmoid flexure of colon, ureter, spermatic vessels.

The Peritoneum. V. Dict. (lat, fin

Describe it.

A serous membrane forming a closed sac (in *male*), the layer covering the walls being called the *parietal*, that reflected over the viscera, the *visceral* layer; it is coated with a layer of flattened endothelium, its attached surface being connected with subjacent parts by the subperitoneal areolar tissue or fascia.

Does the peritoneum always form a closed sac?

No; in the female the Fallopian tubes open into its cavity.

How is the peritoneum divided for convenience of description?

Into the *greater sac*[15], or that covering upper anterior portion of liver[5], the stomach[6] behind and above, descending to ileum forming the anterior layer of great omentum[16], the under surface of mesocolon[20], the mesentery[21], and reflections[23] upon and between the rectum[11], uterus in female[22], and bladder[13], then lining the anterolateral abdominal walls to reach, from the under surface of diaphragm[17], the upper surface of liver, the starting-point of this description (see Fig. 90).

The *lesser sac,* or *cavity of the great omentum*[16], starting from diaphragm behind, this layer passes over back of liver[18], then covers back and under side of stomach[6], forms the inner layer of great omentum[16], passes over colon[7] completing meso-colon[20], and thence passes over duodenum[9] and pancreas[10] to line the posterior abdominal wall, reaching the point started from; the two cavities communicate through the foramen of Winslow.

FIG. 90.



Describe the foramen of Winslow[18] (Fig. 90).

It is an opening at the constricted portion of peritoneum where it curves around the hepatic vessels.

Does this foramen transmit anything?

No; it is merely the orifice of communication between the two sacs.

What are the omenta?

Three folds of peritoneum, viz. the

Cnn. St. w. Alt. visc.

Gastro-hepatic, or *lesser omentum*[18], extending between the transverse fissure of liver and lesser curvature of stomach, consisting of two layers, the anterior pertaining to the greater, the posterior to the lesser sac, which enfold at the right free border the *hepatic artery*, *common bile-duct*, *portal vein*, *lymphatics*, and *hepatic nerve plexus*.

The *great*, or *gastro-colic omentum*[19], is formed of four layers, two descending, one from the anterior, the other from posterior wall of stomach, and uniting below to pass as low as pelvis, when they ascend to transverse colon, separate and embrace this part of the bowel; the

Gastro-splenic omentum is the fold connecting spleen with stomach, contains the splenic vessels and vasa brevia, and is continuous below with the great omentum.

What are the mesenteries?

Double layers of peritoneum embracing various portions of intestinal tube (except duodenum), which they suspend from posterior abdominal walls; between the folds run the vessels of the part suspended. They are called, the

Mesentery[21], *meso-cæcum, meso-colon*[20], *meso-rectum.*

What other name is given to suspensory folds of peritoneum?

Ligaments, such as,

The *gastro-phrenic,* slinging stomach from diaphragm.

The *longitudinal, two lateral*[17], and *coronary* ligaments of liver.

The ~~*vesical,* five~~ so-called ~~false~~ ligaments.

Two *vesico-uterine*[23], two *recto-uterine*[22], two *broad ligaments* of uterus.

Suspensory ligament of spleen connecting this organ with diaphragm.

Mention the viscera wholly(1) and partially(2) covered with peritoneum and those totally deficient(3) in such investment.

(1) The *spleen, small intestine*[8], *cæcum, transverse colon*[7], *sigmoid flexure, ovaries, uterus, stomach*[6] and *liver*[5] practically, *duodenum* (first part), *rectum*[11] (upper third).

(2) *Duodenum*[9] (descending and transverse),
Ascending colon,
Descending colon,
Rectum (middle third[11]),
Vagina[12] (upper part),
Bladder[13] (posterior surface).

(3) *Rectum*[11] (lower third),
Bladder[13] (base, anterior surface),
Vagina (lower portion).
Pancreas[10],
Kidneys,
Suprarenal capsules,

✓

ORGANS OF VOICE AND RESPIRATION.

The Larynx.

What is the larynx[1] (Fig. 92)?

The voice-organ, formed of cartilages united by ligaments, the segments enjoying movement upon one another, which is effected by numerous muscles.

Mention the component cartilages.

They are nine in number, being the

Thyroid cartilage[1],	*Two arytenoid cartilages*,[7]
Cricoid cartilage[5],	*Two cuneiform cartilages*,
Epiglottis,	*Two cornicula laryngis*.

Describe the thyroid (shield-like) cartilage.

It consists of two ~~*alæ*~~, or lamellæ, united at an acute angle in front, forming a v~~ertical ridge~~e[1], whose more prominent upper portion is called the *pomum* ~~*Adami*[2], or Adam's~~ apple; the

Inner surfaces of the alæ are smooth, giving attachment in front, at the receding angle formed by their junction, to ~~epiglo~~ttis, true and false vo~~cal~~ cords, thyro~~-aryt~~enoid and thyr~~o-epig~~lottidean ~~muscles~~.

The *outer surface* affords attachment along an oblique ridge to sterno-thyroid and thyro-hyoid muscles, below and behind to inferior constrictor muscle.

The *upper border* presents a deep, ~~medium notch~~[1], and is slightly concave on either side.

FIG. 91.





The *lower border* is connected medianly with cricoid cartilage by crico-thyroid m~~embra~~ne, and on each side by crico-thyroid muscle.

The *posterior* borders are thick, rounded, and terminate above and below on each side in *cornua*, the two *superior*[3] having attached to their apices the thyro-hyoid ligament, while the *inferior*[4] ~~articu~~late internally by an oval facet with cricoid cartilage; to this border are also attached the ~~stylo-pharyn~~geus and pa~~lato-pharyn~~geus muscles.

Describe the cricoid ~~(ring-like~~) cartilage.

It is placed with its narrow portion[6] in front, and has on each side two ~~articular~~ facets, one on upper margin behind for arytenoid cartilages, one externally, near lower margin for inferior cornu of thyroid cartilage.

The *upper border* gives attachment in front and at sides to crico-thyroid membrane, at sides to lateral crico-arytenoid muscles.

The *lower border* is connected with ~~first tracheal ring~~ by a fibrous membrane.

The *posterior surface* presents a median vertical *ridge* for attachment of longitudinal ~~fibres of œsophagus~~, and on either side the posterior ~~crico-arytenoid muscles~~ are attached.

Describe the two arytenoid (pitcher-like) cartilages[7].

They are pyramidal in form, and are placed at the upper border of cricoid cartilage at back of larynx; the

Posterior surface of each affords attachment to arytenoideus muscle; the

Anterior surface of each has attached the thyro-arytenoid muscle and false vocal cord; the

Internal surfaces face each other; the

Base presents a concave, smooth, articular facet for cricoid cartilage; the

External angle has attached to it the posterior and lateral crico-arytenoid muscles; the

Anterior angle has attached to it ~~true vocal cord~~; the

Apex articulates with a corniculum laryngis.

Describe the cornicula laryngis.

They are two small conical cartilaginous nodules surmounting the apices of arytenoid cartilages affording attachment to aryteno-epiglottidean folds.

Describe the two cuneiform cartilages.

They are elongated cartilages contained in free borders of aryteno-epiglottidean folds.

Describe the epiglottis.

It is a thin, leaf-shaped lamella of *fibro-cartilage*, attached by apex to reëntering angle of thyroid cartilage just below median notch; it lies at base of tongue, in front of upper opening of larynx, and during deglutition shuts like a lid over laryngeal orifice. Its

Base is free, rounded, curving forward toward base of tongue; the

Apex is attached to receding angle of thyroid by thyro-epiglottic ligament; the

Anterior surface is covered with mucous membrane reflected on to sides and base of tongue, forming the three glosso-epiglottidean ligaments; the

Posterior surface shuts down over laryngeal opening during deglutition; the

Lateral margins give attachment to aryteno-epiglottidean folds.

How are the ligaments of the larynx divided?

Into *extrinsic*, or those connecting thyroid cartilage and epiglottis with hyoid bone; and *intrinsic*, those binding the various cartilages together.

Name the extrinsic ligaments.

Thyro-hyoid membrane, *Two lateral thyro-hyoid ligaments*, containing a small cartilaginous or bony nodule, the *cartilago triticea*; *Hyo-epiglottic.*

Name the intrinsic ligaments.

Crico-thyroid membrane,
Two crico-thyroid capsular ligaments,
Two crico-arytenoid ligaments,
Two crico-arytenoid capsular ligaments,
Two inferior thyro-arytenoid ligaments,
Hyo-epiglottic ligament,
Thyro-epiglottic ligament,
Three glosso-epiglottic folds,
Two superior thyro-arytenoid ligaments.

What are the vocal cords?

The *superior* or *false vocal cords*, on each side, are two folds of mucous membrane enclosing the ~~superior~~ thyro-arytenoid ~~ligaments~~ composed of elastic tissue, stretching between angle of thyroid cartilage below epiglottis and anterior surfaces of arytenoid cartilages; the lower margin forms the upper free crescentic margin of ventricle of larynx.

The *true*, or *inferior vocal cords*, are two strong, ~~yellow elastic~~ fibrous tissue bands, the ~~inferior~~ thyro-arytenoid ~~ligaments~~, covered by thin, tightly adherent mucous membrane, which pass from the receding angle of the thyroid to anterior angles of arytenoid cartilages, the upper border forming the lower margin of ventricle of larynx, the lower continuous with lateral portion of crico-thyroid membrane; each has the ~~thyro-arytenoid muscle~~ lying ~~parallel externally~~; these cords or membranes produce sound by their vibrations, the false cords do not, although indirectly influencing vocalization.

What is the glottis (rima glottidis)?

It is the narrow triangular interval between the true ~~vocal~~ cords; it measures, in males, somewhat less than an inch, by from one-third to one-half inch at its base *behind*, when dilated.

Describe the ventricle of the larynx.

It consists of an oval depression or *sinus* on each side, between the true and false vocal cords leading upward, external to superior cord, into a cæcal pouch, the *sacculus laryngis*.

Describe the sacculus laryngis.

It is a slightly curved conical membranous sac, situated between the inner surface of thyroid cartilage and false vocal cords on each side; internally lined with mucous membrane, perforated by orifices of ducts of sixty to ~~seventy follicular glands~~ lying in the submucous tissue; it is strengthened externally by a fibrous capsule continuous below with superior ~~thyro-arytenoid~~ ligament, and has its inner (laryngeal) surface covered by the inferior thyro-arytenoideus, and its external by thyro-~~epiglottideus muscles~~, which com~~press the sacculus~~, discharging the mucus upon, and for the lubrication of the vocal cords.

What kind of epithelium has the mucous membrane?

A~~bove the~~ false vocal cords squamous epithelium is found, except in front, which is covered by ciliated columnar cells as high as middle of epiglottis; ~~below the~~ false cords, only ciliated columnar epithelium is found.

Describe the muscles of the larynx and epiglottis.

Crico-thyroid[2] (2) (Fig. 92): *origin*, triangular from antero-lateral surface of cricoid cartilage; *insertion*, lower border of thyroid and anterior border of lower cornu; *action*, increases tension of vocal cords by tilting thyroid cartilage forward; *nerve*, superior laryngeal.

Crico-arytenoideus posticus (2): *origin*, lateral half of posterior surface of cricoid; *insertion*, outer angle of base of arytenoid cartilage; *action*, by rotating arytenoid cartilages outward it opens glottis and tightens cords; *nerve*, recurrent laryngeal.

Crico-arytenoideus lateralis (2): *origin*, upper border of side of cricoid; *insertion*, outer angle of base of arytenoid; *action*, by rotating arytenoids inward closes glottis; *nerve*, recurrent laryngeal.

Thyro-arytenoideus (2): *origin*, lower half of receding angle of thyroid and crico-thyroid membrane; *insertion*, base and front surface of arytenoid cartilage; *action*, advances arytenoid and cricoid cartilages thus relaxing vocal cords, also compresses sacculus laryngis; *nerve*, recurrent laryngeal.

Arytenoideus (1): *origin*, back surface and outer border of one arytenoid to be *inserted* at same part of other cartilage; *action*, approximates arytenoids closing back of glottis; *nerves*, superior and recurrent laryngeal.

Kerato-cricoideus: a small muscular bundle; *origin*, from near lower border of cricoid cartilage; *insertion*, inferior cornu of thyroid cartilage; *action*, possibly steadies one cartilage upon the other, only occasionally found, and usually on one side.

Triticeo-glossus: *origin*, cartilage of same name in lateral thyrohyoid ligament; *insertion*, tongue with hyo-glossus muscle; occurs on one or both sides; *action*, unknown.

Describe the muscles of the epiglottis.

Thyro-epiglottideus: *origin*, inner surface of thyroid cartilage; *insertion*, aryteno-epiglottidean fold, margin of epiglottis, outer surface of sacculus laryngis; *action*, depresses epiglottis; *nerve*, recurrent laryngeal.

Aryteno-epiglottideus superior: *origin*, apex of arytenoid cartilage; *insertion*, aryteno-epiglottidean folds; *action*, narrows upper laryngeal orifice; *nerve*, recurrent laryngeal.

Aryteno-epiglottideus inferior: *origin*, arytenoid cartilage above superior vocal cord; *insertion*, inner upper part of epiglottis; *action*, compresses sacculus laryngis; *nerve*, recurrent laryngeal.

Mention the arteries and veins of the larynx.

The *arteries*, are laryngeal branches of superior and inferior thyroid arteries, which inosculate freely; the

Veins, empty into superior, middle, and inferior thyroid veins.

What nerves supply the larynx?

The *superior*, and *inferior* or *recurrent laryngeal*, branches of pneumogastric, the former receiving branches from spinal accessory and superior cardiac sympathetic nerves; the

~~*Superior laryngeal*~~ is chiefly a nerve of ~~sensation~~, supplying laryngeal mucous membrane and ~~arytenoid~~ muscles by its *internal laryngeal* branch, after piercing the thyro-hyoid membrane; and the ~~*crico-thyroid muscle*~~ by its *external laryngeal branch*; the

Recurrent laryngeal is the ~~motor~~ nerve supplying all ~~the~~ laryngeal muscles except ~~*crico-thyroid*~~; the right nerve descends the neck to pass from before backward around the subclavian artery, the left winds in the same direction around the aortic arch, and both nerves ascend in groove between trachea and œsophagus to their distribution, giving off in their course cardiac, œsophageal, tracheal, and pharyngeal branches, and anastomosing with superior laryngeal nerves.

The Trachea and Bronchi.

What is the trachea[3] (Fig. 92)?

It is a membranous, cylindrical air-tube, flattened posteriorly, of a transverse diameter of three-quarters to one inch, extending from lower part of larynx (corresponding to sixth cervical vertebra), to opposite fourth or fifth dorsal vertebra, where it bifurcates to form the

Right and left bronchus; it is composed of ~~sixteen to twenty imper~~fect cartilaginous rings surrounding ~~two-thirds of the cylind~~er, enclosed in a double elastic fibrous membrane connecting the rings with one another, while the space behind has, in addition to the membrane—there forming a single layer—a layer of longitudinal and another of transverse unstriped muscular fibres, the transverse lying beneath the fibrous membrane called the *trachealis muscle*; the last ring, by a triangular hooked-process curving backward between the bronchi, forms two imperfect rings on either side for right and left bronchus; the lining mucous membrane contains much lymphoid tissue and numerous glands, and is covered by several layers of epithelium, the most superficial columnar ciliated.

Give the relations of the trachea in the neck.

It is covered in *front* from above downward by

Anastomosing branches between jugular veins,
Isthmus of thyroid gland
Inferior thyroid veins,
Arteria thyroidea ima (when present),
Sterno-hyoid muscle,
Stero-thyroid muscle,

Posteriorly, it lies upon
Œsophagus,
Laterally, lie
The common carotid arteries[13],
Lobes of thyroid gland,
Inferior thyroid arteries,
Recurrent laryngeal nerves.

Describe its relations in the thorax.

It is *covered* from before backward by

Manubrium of sternum,
Remains of thymus gland,
Left innominate vein,
Arch of aorta,
Innominate artery,
Left carotid artery[12],
Deep cardiac plexus.

Posteriorly, lies the
Œsophagus,
Laterally, on each side, the
Pneumogastric nerves,

What are the bronchi?

Two tubes, structurally like trachea, extending from its bifurcation into the lungs, dividing and subdividing to form the bronchial tubes, in whose walls only scattered cartilaginous plates exist until the diameter of one-fourth of a line is reached, when they become wholly membranous, while the muscular coat and elastic fibrous coat then form a continuous circular layer around the smallest tubes; the mucous membrane is covered with ciliated columnar epithelium; the

Right bronchus is wider, shorter (about ~~one~~ 1-2 inch long), and more horizontal than left; the

Left bronchus is smaller, more oblique, and longer (nearly ~~two~~ inches).

Describe the relations of each bronchus.

The *right*, beginning opposite fourth dorsal, enters lung opposite fifth dorsal vertebra, lying ~~behind~~ superior vena cava and right auricle of heart, having the right pulmonary artery at first below, then anterior to it, and the vena azygos major arching over it from behind.

The *left bronchus*, commencing at same point as right, passes in front of œsophagus, thoracic duct, and descending aorta beneath the aortic arch, the left pulmonary artery lying at first above, then in front of it to enter lung opposite sixth dorsal vertebra.

What vessels and nerves supply trachea and bronchi?

The *arteries* are tracheal branches of inferior thyroid arteries and bronchials from aorta; the

Veins empty into thyroid plexus and bronchial veins; the

Lymphatics empty into mediastinal glands; the

Nerves are branches from pneumogastric, recurrent laryngeal branch of same, and sympathetic nerve.

The Lungs.

Describe them.

They are the two organs of respiration, occupying the greater part of the thoracic cavity, separated by the heart and other contents of the mediastinum, covered by the pleuræ, of a specific gravity of 0.345 to 0.746, weighing together about ~~forty-two~~ ounces, the right being the heavier by two ounces. They are pinkish-white at birth, but irregularly marked by ~~slate-colored~~ patches as age advances from the deposit of carbonaceous matter, or perhaps altered ~~blood-pigment~~, in the lung tissue; they are divided into lobes, the right lung having three[4],[5],[6], the left only two[7],[8]; each lung has an

Apex, extending into root of neck beneath and above first rib, about one to ~~one~~ and a half inches, where it is in contact with first and second part of ~~subclavian artery~~; the

FIG. 92.



Base is broad and concave, resting upon diaphragm, its thin margins extending lower down externally and behind than in front; the

External surface is smooth, convex, marked by the interlobular fissures, and conforms to the shape of the thorax; the

Inner surface is concave, and the left lung presents a depression in front for heart; behind, each has a deep fissure, the *hilum pulmonis*, the point of attachment of root of lung; the

Root[21] is the point of entrance of bronchus, pulmonary and bronchial arteries, and of exit of the pulmonary and bronchial veins and lymphatics; in front of each lies the phrenic nerve and anterior pulmonary plexus, behind, the pneumogastric and posterior pulmonary plexus.

In what order are these various structures arranged?

From before backward, Naturally. - Mitral valve → V Rt. Ht. → A A. is P. to V is a X other

Pulmonary veins,
Pulmonary artery,
Bronchus, etc.
} may be recalled by letters, V, A, B.

From above downward; right lung,

Bronchus,
Pulmonary artery,
Pulmonary veins.
} to be recalled by letters, B, A, V.

(Lt. = tipped back & base toward Rt.)

Left lung,

Pulmonary artery,
Bronchus, etc.,
Pulmonary veins.
} to be recalled by letters, A, B, V.

Describe the structure of the lung.

They have externally a serous coat, a subserous areolar tissue penetrating between the lobules, and their bulk is composed of the pulmonary substance or parenchyma, which is formed of an aggregation of *lobules*, each composed of a terminal bronchial tube with its air cells, and ramifications of pulmonary and bronchial vessels, lymphatics, and nerves, thus forming a miniature of lung; they are pyramidal and large upon the surface, smaller and irregular in the interior; the *alveoli*, or *air-cells*, are small polyhedral alveolar recesses measuring about ~~one-two-hundredths~~ to one-seventieth of an inch, opening into the sides and ends of the intercellular passages —*i.e.*, the irregular terminations of the bronchioles—separated from one another by delicate membranous septa, between the layers of which lies the plexus formed by the pulmonary artery, thus exposing the ~~blood to air on two surfaces~~; they are lined with a layer of squamous epithelium.

What are the vessels and nerves of the lungs ?

They receive blood by bronchial arteries for their own nutrition, and venous blood by pulmonary artery for aëration, *i. e* , absorption of oxygen and emission of carbonic dioxide; the vessels are

Bronchial arteries, branches of aorta. *Pulmonary arteries*, from right side of heart. The *bronchial veins* empty on right side into the vena azygos, on the left, into ~~superior intercostal~~ vein ; the

Lymphatics empty into bronchial glands ; the

Nerves are derived from the anterior and posterior pulmonary plexuses, formed chiefly by branches from the ~~sympathetic~~ and ~~pneumogastric~~ nerves.

The Pleuræ.

What are the pleuræ ?

Two closed delicate serous sacs, consisting of a portion investing each lung and another lining, respectively, the right and left halves of thorax.

Are the pleuræ in contact internally ?

No, except at **one point in front**, leaving a space between them elsewhere.

What is the visceral layer (that covering lung) of each pleura called ?

The *pleura pulmonalis,* investing the lung as far as its root.

What is that called which lines the thorax ?

Pluera costalis. or *parietal layer.*

What is the ligamentum latum pulmonis ?

This *broad ligament of the lung* is **a triangular fold of the pleura** passing downward from lower border of root of lung to diaphragm, retaining lower part of lung in position.

Do the pleural cavities differ from each other ?

Yes, the right is shorter, wider, and extends higher in neck.

What is the mediastinum ?

The median space left between the two pleural sacs, extending from sternum to vertebral column, containing all the thoracic viscera but lungs.

Mention its subdivision, and the contents of each.

The *anterior mediastinum*, bounded in front by sternum, on each side by pleuræ, and behind by pericardium; its contents are

Origins of sterno-hyoid and sterno-thyroid muscles,
Origin of triangularis sterni muscle,
Left internal mammary vessels,
Remains of thymus gland,
Loose areolar tissue containing lymphatics.

The *middle mediastinum* is the broadest portion of interpleural space; its contents are the

Heart enclosed in the pericardium,	*Pulmonary artery,*
Ascending aorta,	*Pulmonary vein,*
Superior vena cava,	*Phrenic nerves.*
Bifurcation of trachea,	

The *posterior mediastinum*, bounded in front by pericardium and roots of lungs, behind by spinal column, and laterally by the pleuræ; its contents are the

Descending aorta,	*Pneumogastric nerves,*
The greater azygos vein,	*Splanchnic nerves,*
The lesser azygos vein,	*Œsophagus,*
Left superior intercostal vein,	*Thoracic duct,*
Lymphatic glands.	

The Urinary Organs.

Where are the kidneys situated ?

In the lumbar regions, one on each side of the vertebral column each kidney reaching from the eleventh rib nearly to the iliac crest, but the right a little lower than the left; they lie embedded in a mass of fat behind the peritoneum, which retains them in place with the aid of their bloodvessels.

What are their size and weight?

Each measures about four inches in length, two in breadth, and one inch in thickness; they weigh, in the male, from four and a half to six ounces; in the female, four to five and a half ounces.

Describe their relations.

The *Anterior surface of right kidney* is in relation with the right lobe of liver, descending portion of duodenum, and ascending colon.

The *Anterior surface of left kidney*, with great end of stomach, lower end of spleen, tail of pancreas, and descending colon.

The *Posterior surface of each* rests upon crus of diaphragm, the anterior lamellæ of ~~transversalis aponeurosis~~ separating it from quadratus lumborum and psoas magnus.

The *Superior extremity* is embraced by the suprarenal capsule.

What is the hilum of the kidney?

A notch or fissure one inch long at central portion of internal border, opening into a cavity, the *sinus*; through it pass the *lymphatics*; the *renal vein* is in front; the *renal artery*[A] next, and *ureter*[U], or excretory duct, behind and below. VAUL.

Describe the naked-eye appearances of a vertical section of a kidney.

The kidney proper consists of a

Cortical structure[1], composed of convoluted and straight uriniferous tubules, bloodvessels, nerves, lymphatics, connective tissue, and Malpighian bodies, and is prolonged down between the pyramids, forming the *columns of Bertin*; a fibrous *capsule* envelops the organ, passes into the sinus which it lines, blending with the sheaths of vessels, nerves, and the calices; the

Medullary substance[2] consists chiefly of straight tubuli uriniferi, "looped tubes of Henle", bloodvessels, etc., arranged in eight to eighteen

Pyramids of Malpighi[2], whose apices are embraced by the calices of the pelvis; the

Pelvis consists of from seven to eighteen *calices*, cup-like tubes, embracing the apices of one or more Malpighian pyramids, converging to form three *infundibula*, these again joining to form the *pelvis*[c], which gradually contracts into the *ureter*[U] or *excretory duct*

Describe the tubuli uriniferi[15] (Fig. 94).

They commence in a dilated cæcal extremity, the *Malpighian*[1] *capsule* and terminate by opening on the free surface of the papillæ[15]. They are called, in various portions of their course, the

FIG. 93.



1'' Labyrinth

1' med. Ray?

[illegible] Pyramids

(Def. [illegible]

Tubules red. brown)

Proximal convoluted tube[3], the much convoluted portion after leaving the Malpighian capsule and situated in the cortical portion.

The *spiral tube of Schachowa*[4], next succeeding, where the tube approaches the medullary portion in a spiral manner; now entering the medullary portion, the tubes suddenly become smaller, quite straight, dipping down into the pyramids to ascend, suddenly enlarge, and again become spiral, and reënter the cortical structure, constituting the descending

Loop of Henle[5], this ascends[7],[8] in an irregular angular manner through the cortex, and terminates in the

Distal convoluted tube[10],[11], which terminates in a narrow

Curved tube[12], continuous with a

Straight, or *collecting tube*[13],[14], terminating on summit of one of the papillæ.

Describe these straight tubes[13],[14].

Traced into pyramids from the papillæ, they run from apex to base, dividing dichotomously, receiving the curved tubes, to enter the cortex much increased in number, where they form conical masses with their bases toward the medullary portion.

What are these conical masses called?

The pyramids of Ferrein. = very St. Tub. + 1 & 2 Spiral

Describe the epithelium of the uriniferous tubules.

It varies in different portions, being flattened polyhedral, angular, and columnar.

Describe the Malpighian bodies (tufts, or vascular glomeruli).

Each is formed by a *renal afferent*[a] (Fig. 95) *arteriole*, which, after piercing the capsule—a pouch-like commencement of a tubulus uriniferus lined with flattened epithelium[k]—breaks up into a tuft of vessels forming a plexus[c], from which arises the *efferent*[e] *vessel* emerging from capsule near point of entrance of afferent arteriole.

Describe the renal circulation.

The

Renal artery, from aorta, divides just external to hilum, into four or five

Primary branches, which divide and subdivide to form

Arteriæ propriæ renales, two of which run along the sides of each Malpighian pyramid, giving off afferent branches to Malpighian bodies in columns, and, bending between bases of pyramids and cortex, send off the (1) interlobular, and (2) arteriolæ rectæ;

(1) *Interlobular arteries*, passing outward between the pyramids of Ferrein to capsule, terminating in stellate plexuses beneath it (*stars of Verheyen*), and also supplying afferent arterioles to Malpighian bodies of cortex, whence issue efferent vessels forming a dense *venous plexus* around adjacent uriniferous tubules; and the

FIG. 94.


Subcapsular layer without Malpighian corpuscles.
a
A
Cortex.
Inner stratum of cortex, without Malpighian corpuscles
c
B
C
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16

(2) *Arteriolæ rectæ*, or *descending branches*, passing from bases to apices of pyramids, there to terminate in the venous plexuses.

The blood is collected by the *stellate venous plexuses* beneath the capsule, forming *venæ interlobulares*, which pass between pyramids of Ferrein, are joined by branches emptying the plexuses around convoluted tubes of cortex , and join the venæ rectæ at bases of Malpighian bodies; these *venæ rectæ* being branches from *plexuses* at apices of medullary pyramids formed by arteriolæ rectæ. The junction of the *venæ rectæ* and *venæ interlobulares* form the *venæ propriæ renales*, accompanying the arteries of the same name, which receive venous twigs from Malpighian bodies of medulla and unite in the sinus to form the

Renal, or *emulgent vein*, which passes out of hilum to empty into inferior vena cava, the right vein being the shorter.

FIG 95.



Describe the nerves of the kidney.

They are small, fifteen in number, have ganglia developed upon them, and come from the solar plexus, lower and outer part of semilunar ganglia, and from lesser and smallest splanchnic nerves forming the *renal plexus*, communicating with spermatic plexus.

The Ureters.

What is the ureter?

It is the tubular, cylindrical, excretory duct of the kidney, of the size of a goose-quill, sixteen to eighteen inches long, extending from the pelvis of kidney to base of bladder, into which it opens by a constricted orifice, after having passed obliquely for nearly an inch between its muscular and mucous coats: it has a *fibrous coat* continuous with capsule of kidney and fibrous tissue of bladder, a *muscular* composed of longitudinal and circular fibres, a *mucous* covered with several layers of many shaped epithelial cells.

Describe its course and relations.

It passes obliquely downward along posterior abdominal wall beneath the peritoneum, over the iliac arteries, ~~behind ilium~~ on *right side*, and ~~sigmoid flexure~~ on *left*, enters the ~~posterior~~ false ligament of bladder in male, with ~~vas deferens~~ between it and bladder, and enters bladder obliquely (see supra) about one and one-half inches behind prostate, and two inches from its fellow, at ~~posterior angle of trigone~~; in female it passes along sides of ~~cervix uteri~~ and upper part ~~of vagina~~; the right ureter lies close to outer side of vena cava.

The Bladder.

What is the bladder[1]?

The musculo-membranous reservoir for the urine, situated behind the pubes, between it and rectum in male, between the pubes and uterus in female; when moderately distended its dimensions are, length ~~five inches~~, breadth three inches, capacity about one ~~pint~~; in the child it is an abdominal organ, and also in adults *when distended.* It presents for examination, the

Summit[10] (Fig. 98), connected with umbilicus by a fibro-muscular cord, the ~~*urachus*~~, and by two fibrous cords placed on either side, the obliterated fœtal hypogastric arteries; the

Body lies against posterior surface of pubes, triangular ligament, internal obturator muscles, and—when distended—abdominal walls, its posterior surface being covered by peritoneum, having some coils of ~~small intestine~~ interposed between it and ~~rectum, and uterus in~~ female; the obliterated hypogastric arteries cross its sides obliquely from below, upward and forward, all below them being uncovered with peritoneum; the ~~vas deferens~~ curves from before backward along each side to reach the base of bladder passing across the obliterated hypogastric vessels and along inner side of ureter; the

Fundus or *base*[11] (Fig. 98), is directed downward and backward, resting in the male upon the second portion of rectum, in the female upon lower part of cervix uteri, being adherent to upper part of vagina, but separated from cervix by a fold of peritoneum which is reflected so as to cover a small portion of base; in male the peritoneum passes from rectum to same portion of base; the

Neck, or *cervix*[12] (Fig. 98), is the constricted portion continuous with urethra, surrounded in male by prostate gland, and is directed downward and forward.

FIG. 96.



How is the bladder held in place?

By *true* and *false ligaments*; the true, being the

Anterior, or *~~pubo-prostatic~~*, formed by two folds of the recto-vesical fascia extending from either side of pubic symphysis to front of cervix over upper surface of prostate gland; the

Lateral, of same tissue, passing between lateral surfaces of prostate gland on each side, and sides of base of bladder; the

Urachus[22] (Fig. 98), a fibro-muscular cord covered with peritoneum, stretching between apex of bladder and umbilicus.

The ~~false~~ ligaments are,

Two Posterior, folds of peritoneum passing, in male from sides of rectum, in female from sides of uterus, to postero-lateral aspect of bladder, and contain the obliterated hypogastric arteries, the ureters, vessels, and nerves.

Two lateral, peritoneal folds passing from iliac fossa to sides of bladder.

Superior[23] (Fig. 98), a peritoneal fold passing over urachus and obliterated hypogastric arteries from apex of bladder to umbilicus.

What is the ~~urachus~~[22] (Fig. 98)?

The remains of a tubular canal connecting bladder with ~~allantois~~ *~~in foetus~~*.

Describe the structure of the bladder.

It has four coats, viz., a

Peritoneal[23],[24] (Fig. 98), covering posterior surface from entrance of ureters at base to its summit, passing on to sides whence it is reflected to abdominal and pelvic walls: a

Muscular, consisting of fibres spirally arranged forming figure-of eight loops, the more superficial being nearly *longitudinal*, the deeper nearly *circular*, corresponding to same-named layers of older authors; these form seven more or less distinctly marked layers, the loops directed toward, and embracing the urachus and urethra respectively, the aggregation of these loops around the neck forming the sphincter and the fibres passing in all directions are continued into prostatic urethra (Pettigrew): a

Cellular, which is a layer of areolar tissue intimately blended with mucous membranes binding it to the muscular coat; a

Mucous coat, covered by transitional epithelium, the superficial layer of polyhedral cells, below club-shaped and smaller spindle-celled: it contains a few mucous follicles and numerous racemose glands lined with columnar epithelium near the neck; it is thrown into folds or *rugæ* when bladder is empty.

FIG. 97.



What are the muscles of the ureters?

Two oblique bands rising behind the orifices of the ureters which converge, to be inserted by a fibrous process into the middle lobe of prostate gland.

What is the trigonum vesicæ or trigone vesical[2] (Fig. 97)?

A triangular smooth surface at base of bladder, with apex forward, of a paler color than remainder of mucous membrane, which is intimately adherent to subjacent tissues: the

Lateral boundaries are two slightly marked ridges on each side passing backward and outward from the apex formed by urethral opening to orifices of ureters[3], corresponding to the muscles of these ducts, the openings being placed about two inches from one another and one and a half inches behind urethral opening, a line connecting the two forming the *base of the trigone.*

What is the uvula vesicæ[4] (Fig. 97)?

A slight elevation of the mucous membrane projecting from lower anterior portion of floor of bladder into orifice of urethra.

Mention the arteries supplying the bladder.

Superior, middle and inferior vesical, and small twigs from *obturator and sciatic arteries* in male, with additional branches from *uterine and vaginal,* in female: the *veins* form intricate plexuses around neck, sides, and base, emptying into internal iliac vein; the *lymphatics* accompany the vessels.

Give the nerve-supply of the bladder.

Branches from hypogastric plexus of sympathetic supply upper part, and the third and fourth sacral nerves the base and neck.

The Male Urethra.

Describe this canal.

It extends from neck of bladder[4] to meatus urinarius, measuring from eight to nine inches, presenting a double curve when penis is flaccid, but a single one with convexity downward during erection.

Name and describe each of its divisions.

They are the prostatic[7,8], membranous, and spongy.

The *prostatic portion*[7,8] (Fig. 97) is that portion passing from base to apex of prostate gland, running nearer the upper surface, measuring one and a quarter inches long, is the widest and most dilatable section of urethra and is spindle-shaped; upon the median aspect of the floor is a narrow longitudinal ridge eight to nine lines long, the *verumontanum* or *caput gallinaginis*[5] (Fig. 97) containing muscular and erectile tissue: on either side is a slight fossa, its floor presenting the numerous openings of the prostatic ducts, called the *prostatic sinus.*

The *sinus pocularis*, or *uterus masculinus*[6] (Fig. 97), is a depression extending about a quarter of an inch upward and backward beneath middle lobe of prostate in the median line at fore part of verumontanum, upon or within whose margins are the slit-like orifices of the *seminal or ejaculatory ducts*[6] (Fig. 97).

The *membranous portion*[a] (Fig. 97) is that portion, three-quarters of an inch long, extending between apex of prostate and bulb of corpus spongiosum, which is included between the layers of the triangular ligament about one inch beneath the pubic arch; it is surrounded by the compressor urethræ muscle, and is the narrowest portion of urethra, except meatus.

The *spongy portion*[d,h] (Fig. 97) is the longest measuring about six inches, is so called from being contained in the corpus spongiosum, extends from membranous portion to meatus urinarius, and presents a posterior dilatation, that of the bulb[8] (Fig. 98), and one anterior, the *fossa navicularis*[16] (Fig. 98), the former having opening into it the ducts of Cowper's glands, the latter fossa being situated within the glans penis, having an opening directed forward in its roof, the *Lacunus magnus*, the orifice of a large mucous crypt; there are also here numerous other scattered openings of glands similarly directed.

FIG. 98.



What is the meatus urinarius?

The anterior orifice of the urethra, a vertical slit about three lines long bounded on each side by a small labium, and is the narrowest portion of the canal.

Describe the structure of the urethra.

It possesses three coats, viz., a mucous, forming a part of the genito-urinary membrane internally and continuous with the skin externally, having numerous mucous glands imbedded in its submucous tissue, which open on its free surface, and is covered with columnar epithelium except near meatus where it is squamous.

The *muscular coat* consists of longitudinal fibres externally, circular within, the latter lying beneath urethral mucous membrane for its whole length, the former leaving the circular fibres at the bulb to envelop the spongy body beneath its fibrous coat, again rejoining the circular fibres at the meatus urinarius; both muscular layers are in direct continuity with those of bladder. (For voluntary urethral muscles see p. 331.)

Erectile, this is a thin layer of erectile tissue surrounding the membranous and prostatic portions, anteriorly becoming continuous with that of corpus spongiosum.

Female Urethra.

Describe this canal.

It is about one and one-half inches long, extends from bladder to meatus urinarius above anterior vaginal wall, pierces the triangular ligament, and is embraced by compressor urethræ muscle as in male: its structure is similar to that of male urethra, it is lined by laminated squamous epithelium merging into spheroidal near bladder, is a quarter of an inch in diameter, but is capable of much greater dilatation; and its anterior opening (meatus) is situated near anterior margin of vagina, about one inch below clitoris, surrounded by a prominence of mucous membrane.

The Male Generative Organs.

Where is the prostate gland situated? (Fig. 97)?

It surrounds neck of bladder and commencement of urethra, lying in pelvic cavity posterior to deep perineal fascia, behind and below symphysis pubis upon the rectum: through it pass the urethra and seminal ducts.

Describe its form and size.

It resembles a ~~horse-chestnut~~ in form, measures ~~one-and-a-half~~ inches transversely, ~~one inch~~ antero-posteriorly, three-quarters of an inch in depth, and weighs about ~~six drachms~~: the base is directed backward toward bladder: it consists of two equal-sized lateral lobes and a middle lobe, (a small transverse band or rounded triangular eminence placed between lateral lobes immediately beneath neck of bladder behind commencement of urethra)

Describe its structure?

It is composed of numerous follicular pouched glands opening into elongated canals which join to form twelve to twenty excretory ducts imbedded in the interstices of a stroma formed of interlacing bundles of ~~unstriped muscle~~, the whole being inclosed in a ~~fibrous capsule~~.

What retains the gland in place?

The ~~pubo-prostatic~~ ligaments of bladder, posterior layer of deep ~~perineal~~ fascia, and anterior portion of ~~levator ani~~ muscle.

Describe the situation and structure of Cowper's glands?

They are two small lobulated glands of the size of [redacted], lying between the ~~two~~ layers of the ~~deep perineal~~ fascia in front, close behind bulb of urethra, ~~surrounded~~ by ~~compressor urethræ muscle~~ and opening, by ducts one inch long which pass obliquely forward beneath mucous membrane, on the floor of bulbous portion of urethra.

The Penis.

Describe the penis?

It has a root, body[5] (Fig. 98), and extremity or glans penis[6], and consists of three elongated cylindrical masses of erectile tissue, composed of a fibrous sheath which sends ~~inward numerous~~ interlacing bands (~~trabeculæ~~) forming numerous meshes in which lie the bloodvessels. The upper two cylindrical bodies lying side by side, like a ~~double-barrel~~ gun, are called the corpora cavernosa[5]; the third, much smaller, lying in median line beneath, like the ramrod of a gun, is the corpus spongiosum[7].

Describe the corpora cavernosa.

Situated as just described, they are intimately connected in their anterior three-fourths where they are in contact, presenting a median dorsal groove for vessels and nerves, an inferior median one for corpus spongiosum, while their posterior fourth diverges, forming the *crura*[4], which are attached to the rami of pubes and ischium anterior to the tuberosity; near their point of junction with one another they become slightly enlarged, forming, on either side, the ~~*bulb of*~~ *the corpus cavernosum.*

What other structure besides the crura secures the root of the penis to symphysis pubis?

The *suspensory ligament*, a fibrous membrane.

What is the septum p~~ectin~~iforme[5]? = bet 2 Corp. Cav.

The anterior portion of the v~~ertical fibrous~~ septum which is incomplete, the fibrous bands resembling in their arrangement the teeth of a comb; the septum and fibrous sheath contain numerous elastic and muscular fibres in addition to the white fibrous tissue.

Describe the corpus spongiosum.

It incloses the urethra[15], lying medianly below at junction of corpora cavernosa, commencing behind in front of deep perineal fascia between the crura of the corpora cavernosa as a rounded enlargement, the ~~*bulb*[8],~~ which is surrounded by the ~~accelerator urinae~~ muscle. Anteriorly it forms a conical enlargement, flattening from above downward, which caps the blunted end formed by the corpora cavernosa, the *glans penis*[6], the margin of whose base is called the *corona glandis*, the constriction behind the *cervix*.

What other parts of importance does the penis present?

The *meatus urinarius*, the external orifice of urethra opening at summit of glans penis.

The *prepuce*, a portion of skin of penis prolonged forward so ~~as~~ & then turned back int either completely or partially to cover glans penis, lined ~~internally~~ by a ~~tissue~~ re~~sembling muc~~ous membrane.

The *frænum præputii*, a fold of mucous membrane passing from behind meatus to bottom of a depressed raphé to become continuous with under margin of prepuce.

Glandulæ Tysonii odoriferæ, numerous small lenticular sebaceous glands upon corona and cervix of glans, secreting sebaceous matter with a peculiar odor.

What is meant by erectile tissue ?

An intricate venous plexus formed by the interspaces between the fibrous trabeculæ, the blood being delivered in the following ways: (1) by arteries terminating in ordinary capillaries—this is the arrangement in the corpus spongiosum and glans—(2) fine convoluted arterial twigs opening directly into venous network by funnel-shaped extremities. (Spleen)

What are the helicene arteries ?

Convoluted, tendril-like arterial branches opening directly into venous plexus (supra), as just explained, most abundant in back parts of spongy and cavernous bodies.

Name the arteries, veins, and lymphatics of the penis.

The arteries are all branches of internal pudic, viz.,

Artery of bulb, to corpus spongiosum.

Arteries of corpora cavernosa, to these bodies.

Dorsal arteries of penis, to corpora cavernosa, glans, prepuce, and skin; the

Veins return blood by dorsal vein, prostatic plexus, and pudental veins.

Lymphatics, the superficial end in inguinal glands, the deep join deep pelvic lymphatics; the

Nerves are branches of internal pudic and hypogastric plexus of sympathetic.

The Testes and their Coverings.

What are the testes or testicles (Fig. 99) ?

They are the procreating glands, those which secrete the seminal fluid, are of ovoid form compressed laterally, and are each obliquely suspended in the scrotum by their own spermatic cord.

They measure from one and a half to two inches long, one inch in breadth, and one and a quarter inches antero-posteriorly, weighing from six to eight drachms, the left being slightly larger.

What is the scrotum?

A c~~utaneous pou~~ch containing the testicles and part of spermatic cords, formed of ~~integument~~ externally and beneath this of a reddish contractile layer, the ~~dartos,~~ continuous with contiguous ~~superficial fascia~~ and sending inward a partition, the ~~septum scroti,~~ dividing the scrotum into two compartments.

Describe the coats of the testis.

really inside of Tunica Albuginea

They are, the

Tunica vaginalis testis, a serous coat originally derived from peritoneum, consisting of a portion investing the testis, the visceral layer, or *tunica vaginalis propria,* and a parietal layer, or *tunica vaginalis reflexa*; the

Tunica albuginea is formed of white fibrous tissue surrounding the gland and reflected into its interior at its postero-superior border to form an incomplete vertical partition, the *mediastinum testis,* or *corpus Highmorianum,* from which fibrous septa pass, separating the glandular lobules; the

Tunica vasculosa, or *pia mater testis,* consists of a plexus of blood-vessels bound together by areolar tissue, which invests the *inner surface* of the tunica albuginea and sends off processes between the lobules.

What is meant by the coverings of the testis?

The structures with which the testes become ~~invested~~ in their ~~passage, pre~~vious to birth, from the abdomen along the inguinal canals into the scrotum.

Name these coverings.

Skin, } scrotum, closely adhering to each other.
Dartos, } 2 Superficial fascia.
Intercolumnar, or *external spermatic fascia,* derived from margins of e~~xternal abdo~~minal ring.
Cremaster muscle, derived from lower border of ~~internal obli~~que muscle (some authors deny this).

Infundibuliform, or *fascia propria*, a downward continuation of infundibuliform process of transversalis fascia.

Tunica vaginalis, derived from peritoneum (supra).

Describe the structure of the testis.

It consists of some hundreds of *seminiferous* tubules, one-one-hundred and fiftieth to one-two-hundredths of an inch in diameter, convoluted so as to form three hundred conical *lobules*, with apices toward mediastinum; the tubes then unite to form twenty or thirty larger ducts, one-one-fiftieth of an inch in diameter, called *vasa recta*[c], which, passing upward and into the mediastinum, form a close network of anastomosing tubules, the *rete testis*; these, at the upper end of the mediastinum, terminate into twelve to twenty *vasa efferentia*[e], perforate the tunica albuginea, and become enlarged and convoluted, forming *coni vasculosi*[f], which aggregated compose the globus major of epididymis; the tubules consist of a membrana propria lined with a layer of polyhedral cells, with two or more inner layers of spheroidal cells, which divide into epithelial cells ultimately to become converted into spermatozoids; the vasa recta and tubes of retè testis have thin walls lined by one layer of squamous cells; the vasa efferentia and epididymis have thicker walls, containing muscular tissue and are lined with columnar epithelium.

What is the epididymis?

A convoluted tube, some twenty feet long, lying along posterior border of testis, commencing on testicle side by convergence of tubes of *coni vasculosi*, and ending in the single vas deferens beyond.

Describe its various parts.

The *globus major*[e] is formed by the coni vasculosi, or *efferent ducts*, which open at intervals into the single duct forming epididymis; the

Body[g] is the central portion formed by the convolutions of the tube bound together by delicate areolar tissue; the

Globus minor[g] is the lower enlarged portion, composed as body is, and connected to testis by fibrous bands and areolar tissue; the

Vas aberrensi is a narrow tube extending up into cord for two or three inches, ending in a blind extremity, occasionally connected with lower part of tube of epididymis, or commencement of vas deferens; the

Hydatids of Morgagni are small bodies attached to epididymis or upper portion of testicle by pedicles; one of these is probably the re~~mains of the *duct of Müller*~~.

Describe the vas deferens[h].

It is the excretory duct of testis, and a co~~ntinuation of epididy~~mis, commencing at globus minor[g] to ascend along p~~osterior sid~~e of ~~testis and~~ epididymis, back of spermatic cord, through the inguinal canal to internal abdominal ring, whence, passing into pelvis, it cr~~osses exter~~nal iliac, curves aro~~und epigastric~~ artery to reach side, then by a curve downward and backward, the base of bladder ~~internal~~ to the u~~reter~~ (Fig. 96); here it lies between the bladder and rectum, coursing along inner border of seminal vesicle[5] (Fig. 96) on each side, becoming enlarged and sacculated, but narrowing at base of prostate where, joined by duct of vesiculæ seminales[6] (Fig. 96), it forms the ejaculatory duct[7] (Fig. 96). It measures about ~~two feet~~ long, is about one and a quarter lines in diameter, its lumen measuring but one-third line, and has thick, rigid walls, presenting a hard, cord-like sensation when rubbed between the fingers.

FIG. 99.



What are the arteries of the testis and its coverings?

The gland itself is supplied by the *spermatic artery*[m] (Fig. 99), the coverings receive blood from *superficial external pudic* and *deep external pudic* from femoral, *superficial perineal* from internal pudic, *cremasteric* from epigastric, and *artery of vas deferens* from superior vesical.

The *veins* leave back of testicle forming the *pampiniform plexus*, which ends in a single trunk emptying on right side into ascending vena cava, on left into left renal vein.

Veins of the same name as arteries supplying coverings, return blood to femoral, external and internal iliac veins.

What nerves go to each testicle and its coverings?

Branches from spermatic plexus of sympathetic to testicle itself, to coverings and cord the *ilio-inguinal, ilio-hypogastric, superficial perineal, inferior pudendal,* and *genital branch of genito-crural nerve.*

What are the component parts of the spermatic cord, and how are they disposed?

Vas deferens,
Spermatic artery,
Cremasteric artery,
Artery of vas deferens,
Spermatic veins,
Spermatic nerve-plexus,
Branch of ilio-inguinal nerve,
Branch of genito-crural nerve,
Vas aberrens (when present),
Lymphatics.

These form a cord bound together by connective tissue about ~~four inches~~ long, extending from globus minor to internal ~~abdom~~inal ring.

Describe the vesiculæ seminales.

They are two lobulated membranous receptacles for ~~the semen~~, which they ~~dilute~~ with their own secretion. They are pyramidal in form, are about two and a half inches long by five lines broad, by two to three lines thick, and lie in contact with base of bladder, diverging from each other from base of prostate to near entrance of ureters; they join by their anterior pointed extremities with vas deferens, forming on each side the *ejaculatory ducts*[7] (Fig. 96), terminating in prostatic urethra by slit-like orifices on each side of the *sinus*[5] (Fig. 97) at front of verumontanum.

Describe the descent of the testes.

During early fœtal life the testes lie at back part of abdomen, behind the ~~peritoneum~~, just below and in front of kidneys; attached to lower end of epididymis, and attaining its full development from fifth to sixth fœtal month is the ~~*gubernaculum*~~ *testes*, which = Rud: contains ~~muscular~~ tissue; this divides below into three portions, passing to Poupart's ligament, to os pubis and rectus muscle, and to ~~dartos at~~ bottom of scrotum; the gubernaculum is supposed to contract and so cause descent of testicle, but this is a moot point. Between the fifth and sixth month each testis reaches the iliac fossa, by seventh it enters internal abdominal ring, by eighth month it has reached the scrotum, carrying before it the peritoneal sac, the upper part of which usually becomes obliterated just before birth, the lower portion then forming the *~~tunica vaginalis~~ testis* completely cut off from abdominal cavity; the other structures in front of the testis are likewise carried onward, forming its covering, as already described (see page 275).

Female Organs of Generation.

What is the vulva?

The term includes the following organs:

- *Labia minora*[13],
- *Mons veneris*[4],
- *Labia majora*[5],
- *Vaginal orifice*[15],
- *Clitoris*[6],
- *Meatus urinarius*[14],

These parts are also called the *pudendum* and *external organs of generation.*

Describe the mons veneris[4].

It is a rounded eminence surmounting vulva in front of pubes, formed by a collection of fatty tissue, and at puberty becomes covered with hair.

Describe the labia majora[5] and minora[13].

The *Labia majora*[5] are two prominent longitudinal ~~cutaneous folds~~ passing downward from mons veneris to anterior part of perineum enclosing the common *urino-sexual opening.* Each labium is formed ~~externally of hair-covered~~ skin, ~~internally,~~ of mucous ~~membrane~~ enclosing between these *layers* ~~fatty areolar~~ tissue and a structure resembling the ~~dartos~~; the junction of the labia in front constitutes the *anterior commissure*, that behind, the *posterior commissure*; the

Labia minora, or *Nymphæ*[13], are two thin folds of mucous membrane, containing numerous ~~sebaceous~~ glands, inside the labia majora, running from clitoris[6],—above which they form a hood-like prepuce,—obliquely downward for about one and a half inches on each side of vaginal orifice, to be lost in the labia majora.

Describe the remaining structures forming the vulva.

The *clitoris*[6] is a small penis with root, body, and glans, composed of two corpora cavernosa formed of erectile tissue attached to rami of pubes and ischium by two crura[7]; it has a suspensory ligament and two erectores clitoridis muscles; the

Vestibule is the triangular smooth surface below clitoris, bounded on each side by labia minora; the

Meatus urinarius[14] is placed at back part of vestibule, about one inch below clitoris, near vaginal margin~~, surrou~~nded by a ~~prominence of mucous mem~~brane; the

Vaginal orifice[15] is of an elliptical form, usually more or less closed in the virgin by hymen, and is surrounded by **sphincter vaginæ muscle**, the anologue of male accelerator urinæ muscle; the

FIG. 100.



Hymen is a thin fold of mucous membrane, variously shaped, but usually semilunar, with concavity upward, stretched across lower part of vaginal orifice; it may occasionally form a complete occluding membrane, the condition being then known as *imperforate hymen*; it may be practically absent in the virgin, and again, may persist after copulation, but at that time is usually ruptured, giving rise to small, rounded elevations surrounding vaginal outlet, called *carunculæ myrtiformes*; the

Glands of Bartholine[16] are small, oblong, reddish-yellow bodies, lying on each side of commencement of vagina, each opening by a long single ~~duct~~ e~~xternal to hymen~~ on inner side of each labium ~~majorum~~; the = Vulvo Vag. Gl. . lubricates [illegible]

Bulbi vestibuli[12] are two oblong masses—a ve~~nous plexus~~ enclosed in a fibrous envelope—extending from clitoris along each side of vestibule, a little behind labia majora; the

Pars intermedia[11] is another small venous plexus immediately in front of preceding, with which it is continuous as well as with glans clitoridis; the

Fourchette is a small transverse mucous fold, just within posterior commissure; the

Fossa navicularis is the space between fourchette and posterior commissure.

The Vagina.

Describe it.

It is a cylindrical membranous canal flattened from before backward, lying between rectum and bladder, extending from vulva to uterus, curved forward and downward conforming to axis of pelvis and of outlet. Narrow at its orifice it is larger above, its anterior wall measuring about f~~our in~~ches, its posterior ~~six,~~ being attached higher up on os uteri *behind* than in *front.*

Describe its structure.

It is formed of an ~~external~~ layer of ~~erectile t~~issue, a muscular coat, and a lining mucous membrane with a *median anterior* and *posterior raphé* or *columns of the vagina*; it contains mucous glands; the epithelium is of the ~~squamo~~us variety.

Give the relations of the vagina.

Its *anterior surface* is in relation with base of bladder and urethra;

The *posterior surface* is connected with lower three-fourths of rectum, the upper fourth being separated from bowel by recto-uterine fold of peritoneum forming *Douglas's cul-de-sac or pouch*;

Laterally, above it gives attachment to broad ligaments of uterus, below to levatores ani muscles and recto-vesical fascia.

The Uterus and its Appendages.

Describe the position and parts of uterus.

It is the organ of gestation, situated in pelvic cavity between rectum and bladder, opening below into vagina, which is attached around its cervix. It is ~~pear-shaped~~, flattened from before backward, is about ~~three inches~~ long, ~~two~~ inches in breadth at upper part, and ~~one~~ inch thick, weighing from ~~one~~ to one and a half ounces; it presents for examination the

Fundus[2], which is convex, covered with peritoneum, and placed below level of brim of pelvis; the

Body[1] gradually narrows from fundus to neck, with ~~anterior flattened~~ surface covered with peritoneum for its upper three-fourths, and a posterior peritoneum-covered convex surface, while each lateral margin is concave, affording ~~attachment~~ to ~~Fallopian~~[9] tube above, to ~~round ligament~~ in front and below this, and to ~~ovarian ligament~~ below and behind both; the

Cervix[3] is the lower, rounded, constricted portion of uterus projecting into upper portion of vagina, which is attached further up behind than in front; the

Cavity of the Body is small, ~~triangular~~, flattened from before backward, its upper extended lateral angles forming funnel-shaped cavities at the bottom of each of which opens, by a minute orifice, the Fallopian tube; at the inferior angle is the small opening into the cavity of the cervix called *os internum* or *internal os uteri*; the

FIG. 101.



Cavity of the cervix is ~~spindle-shaped~~, ~~flattened~~ antero-posteriorly, and opens into cavity of uterus above by *internal os uteri*, and below into vagina by the *external os uteri*; a median longitudinal ~~crest~~ of mucous membrane on back and front walls from which proceed obliquely other smaller ones, forms the ~~*arbor vitæ*~~, hardly noticeable after the first labor; the

Os externum or external os uteri (also os tinctæ)[4] is a transversely ovoidal opening from cervix into vagina which presents an *anterior and posterior lip*.

Describe ligaments of uterus.

Six are peritoneal folds, viz.,

Anterior or vesico-uterine, two crescentic folds passing between uterine neck and back of bladder; the

Posterior or recto-uterine, passing between sides of uterus and rectum; the peritoneal pouch thus formed between rectum and uterus being called *Douglas's cul-de-sac* or *recto-vaginal pouch*; the

Two lateral or broad ligaments[7] (Fig. 101), stretched between sides of uterus and ~~lateral walls of pelvis~~, dividing this cavity into two portions, and containing between their folds the ~~Fallopian[8] tubes~~ and the ~~round ligaments~~[8]. = Retra - peritneal

Describe the round ligaments of the uterus[8].

They are two rounded cords, ~~four or five inches~~ long, commencing at superior angles of uterus to pass forward and outward through internal abdominal ring into inguinal canal to become lost, after dividing into three fasciculi, in the fatty tissue of mons veneris; they consist of ~~dense fibrous tissue~~ and ~~unstriped muscle~~, enclosed, in the fœtal state, by a process of peritoneum extending a short distance into inguinal canal, the so-called *canal of Nuck,* usually obliterated in adult, but sometimes pervious throughout life.

Describe the structure of uterus.

The womb has three coats, viz., a

Serous, derived from peritoneum covering fundus of organ and its posterior surface, but only the superior three-fourths of its anterior surface; a

Muscular, forming bulk of uterus, composed of bundles of unstriped muscular tissue interspersed with areolar tissue, blood-vessels, lymphatics, and nerves. The fibres are disposed in three layers, viz., transverse, in front and behind fundus; middle layers, passing longitudinally, obliquely, and transversely; and circular, arranged in two hollow cones whose apices surround the orifices of the Fallopian tubes, whose bases fuse in middle of uterine body; at cervix these fibres pass transversely; a

Mucous, thin, smooth, and closely adherent to subjacent parts, covered with columnar ~~ciliated~~ epithelium, and containing numerous tu~~bular follicl~~es, most numerous in the cervix; when their ducts become obliterated their secretion is retained, forming small vesicular elevations within os and cervix uteri called ~~Ovula of Nabothi.~~ = Retent. Cyst.

Name the uterine vessels and nerves.

The

Arteries are the *uterine*, from internal iliac, and *ovarian*, from aorta, which pursue a very torturous course and freely anastomose.

The *veins* accompany arteries and terminate in uterine plexuses; during pregnancy they are called *uterine sinuses*, consisting of the lining membrane of the vein adhering to the walls of canals channelled through uterine substance; the

Lymphatics, terminate in pelvic and lumbar glands; the

Nerves, are branches of inferior hypogastric, and spermatic plexuses, and of third and fourth sacral nerves.

What are the Fallopian tubes?

They are the *oviducts*, two tubes about ~~four~~ inches long, extending between layers of broad ligament on each side from superior angle of uterus to sides of pelvis; the canal of each tube is very minute at its commencement the *ostium internum* at the superior angle of the uterine cavity, continuing so for inner half, when it gradually widens into the trumpet-shaped *ostium abdominale*[10], called from the fringe-like ~~*fimbriæ*~~[10] which surround the opening—~~one of which is attached to~~ ovary—the *fimbriated extremity*[10]. Their walls are formed of a serous or peritoneal coat, a muscular, formed of longitudinal and circular fibres, and a mucous coat covered with ~~ciliated~~ columnar epithelium, continuous on one side with uterine lining, on other with peritoneum.

Describe the ovaries[11]. Entirely in perit.

They are two flattened, ovoid bodies suspended by their anterior margins from back of broad ligaments[7], behind and below Fallopian tubes, upon either side of uterus; they are attached by their inner extremities to uterus by the ovarian ligaments and by their outer ends to one of fimbriæ of Fallopian tube; their dimensions are, *length*, ~~one~~ and one-half inches; *width*, three-quarters of an inch; *thickness*, one-third of an inch.

[illegible] as signet in ring. [illegible]

[illegible] not Covr. = Col. ep. [illegible]

" " = Germinal ep.

ova [illegible] = Graafian follicles.

Describe the structure of the ovaries.

They consist of numerous *Graafian vesicles* imbedded in a fibrous stroma, covered externally by ~~modified peritoneum~~, having ~~columnar instead of squamous~~ cells.

The *stroma* consists of numerous spindle-cells with connective tissue and abundant bloodvessels; the ~~condensed peripheral~~ layer surrounding the organ was formerly described as the *tunica albuginea.*

What are the Graafian vesicles?

Ovisacs, minute vesicles from one-one-hundredth of an inch in diameter, to even ~~one-one-twentieth of an inch~~ after puberty; microscopically, they are seen to consist of an *external fibro-vascular coat* connected with stroma by a vascular network, an internal coat, or *ovicapsule* lined by a layer of nucleated cells, the *membrana granulosa,* which are heaped up around ovum at that part of *Graafian vesicle* nearest the ovarian surface, forming the *discus proligerus*; it contains also a transparent, albuminous fluid, suspending the ovum in the immature vesicles.

Describe the human ovum.

It is a spherical mass of protoplasm, one-one-hundred and twenty-fifth of an inch in diameter, and consists of the

Vitelline membrane, or *zona pellucida,* a transparent sac surrounding the

Vitellus, or *yelk,* a fluid containing granules, among which is found the

Germinal vesicle, containing a smaller body, the

Germinal spot, one-thirty-six hundredths of an inch in diameter.

What is the corpus luteum?

An irregular yellow spot in ovary at site of a ~~ruptured Graafian~~ vesicle, which differs in appearance according as it is a *true* or *false corpus luteum.*

Describe the differences between these two kinds of corpora lutea.

The ~~*true corpus luteum* is that of pregnancy~~ which increases up to a certain point, reaching sometimes the size of ~~a mulberry~~, and usually does not entirely disappear until ~~full term~~.

The *false corpus luteum* is that following ordinary menstruation, is much ~~smaller~~, and d~~isappears in about three mo~~nths.

Name the ovarian arteries, veins, and nerves.

The *arteries* are the ovarian, from aorta, anastomosing with uterine artery; these vessels also supply Fallopian tubes; the *Veins* follow arteries and form a plexus near ovary, called the *Pampiniform plexus*; the *nerves* are derived from inferior hypogastric or pelvic plexus, from ovarian plexus, and the Fallopian tubes receive branches from one of the uterine nerves.

What is the parovarium ?

It is also called the *organ of Rosenmüller*, is the remains of a fœtal structure, and in the adult consists of a few closed epithelial-lined, convoluted tubes, one commonly ending in a bulbous hydatid-like swelling; at its uterine end the parovarium is connected with the remains of the Wolffian duct—the *duct of Gaertner*.

The Mammary Glands.

Describe the mammæ.

They exist in both sexes, being in male only rudimentary, but in female are two large hemispherical eminences situated toward lateral aspect of pectoral region on each side, reaching from ~~axilla to sternu~~m, and from third to sixth ~~or seventh~~ ribs; just below the centre projects a small, dark-colored conical prominence, the ~~*nipple*~~, surrounded by a lighter colored ~~*areola*~~, in which are numerous prominent ~~sebaceous~~ glands, which enlarge during pregnancy; the color of both nipple and areola darken during pregnancy, and the latter also extends its area.

Describe the structure of the mammary glands. Rac

They are composed of gland-tissue, of fibrous tissue connecting the lobes, and fatty tissue in intervals between lobules; the lobes consist of lobules formed of a number of rounded vesicles grouped about a small lactiferous tube into which their ducts open, which, by their union, finally form ~~fifteen or twenty excretory~~ ducts, or *tubuli lactiferi*, which converge toward the areolæ, dilating beneath it into the ~~*ampullæ at base of nippl*~~ they contract into straight tubes perforating the summit of the nipple; the lobules are surrounded by a ~~dense capillary~~ network during lactation, as is also the ~~nipple~~, which becomes ~~erected when irritated~~, partly from ~~fulness of blood~~, partly from contraction of its muscular tissue.

Name the vessels and nerves of the mammæ.

The *arteries* are derived from thoracic branches of axillary, intercostal, and internal mammary arteries; the

Veins form an anastomotic circle around the base of the nipple, called the *circulus venosus*, from which large vessels radiate to terminate in axillary and internal mammary veins; the

Lymphatics chiefly run along lower border of pectoralis major to axillary glands, a few pass through intercostal spaces to anterior mediastinal glands; the *nerves* come from anterior and lateral thoracic cutaneous.

THE ORGANS OF SENSE.

The Skin and its Appendages.

Of what parts does the skin consist?

Epidermis, or *cuticle*, formed of	*superficial epithelial layers* and the *rete mucosum*, or *deep epithelial layers*, which contain the pigment.
Derma, *cutis vera*, or *true skin*, formed of	*papillary layer*, upon which lies rete with an interposed basement membrane; *corium*.

Enumerate the accessory structures contained in the skin.

The *tactile corpuscles*, ~~in papillæ~~ of sensitive parts.

Ducts of sweat-glands, *Hair-follicles*, into which open sebaceous glands,	passing through all the layers of the skin.

Where do sweat- and sebaceous-glands with the hair follicles lie?

Chiefly in the subcutaneous fatty tissue, but sometimes in the deepest layers of the corium.

Describe the epidermis.

It is composed solely of epithelial cells, the deepest layer being columnar, more rounded in the middle portions, and flat, scaly and horny on free surface; the deepest, softest layer is accurately moulded upon the papillary layer of the derma, and contains the skin pigment; it is called the *rete mucosum.*

Describe the derma.

Is a tough, flexible, and highly elastic tissue, protecting subjacent parts, and is the chief organ of the sense of touch; excretion is effected by its various glands, and absorption also takes place from its surface. The derma consists of the

Papillary layer, situated upon its free surface, presenting innumerable, minute, vascular conical eminences, averaging in length one-one-hundredth of an inch by one-two-hundred and fiftieth of an inch in diameter at their bases, scattered irregularly in slightly sensitive parts, but arranged in parallel curved lines, forming ridges, in highly sensitive regions; each papilla contains one or more capillary loops, and one or more nerve fibres, some terminating in oval-shaped bodies, the

Tactile corpuscles, where touch is most highly developed.

The *corium* is composed of interlacing bands of white fibrous tissue, with some yellow elastic fibres, unstriped muscular fibres wherever hair exists, lymphatics, bloodvessels, and nerve-plexuses.

What are the nails?

Flattened, horny structures of modified epithelium, covering the dorsal aspects of derma of distal phalanges of fingers and toes. They are convex externally, and have a *root,* imbedded in a groove of skin, whence the growth in length comes; a *matrix,* that portion of derma beneath, by which the nail grows in thickness; the *lunula,* the white crescentic portion next to root, its color owing to diminution in size, number, and vascularity of the papillæ, which are disposed in longitudinal rows elsewhere in the matrix.

What are the hairs?

They are also modified epidermis, found everywhere in skin, except palms of hands and soles of feet, but varying in size. They each have a

Root, bulbous in form, springing from a vascular papilla at the bottom of an involution of the epidermis and corium, called a hair-follicle, which sometimes extends into subcutaneous cellular tissue; into the hair-follicle open one or more sebaceous glands; the

Shaft consists of a *central* or *medullary portion*, a *fibrous portion*, and a *cortical* of thin flattened scales; the medulla is commonly absent in fine hairs; the

Point is formed only of the fibrous and cortical portions.

Describe the sebaceous glands.

Most abundant in scalp, face, arm-pits, around nasal and oral apertures, and in external auditory canal; the largest are the Meibomian glands of eyelids; they are small sacculated glands, lodged in the corium or subcutaneous tissue of nearly every part, except palmar and plantar surfaces; the ducts usually open into hair follicles, but sometimes on general surface of skin.

Describe the sweat glands.

They are each formed of a single convoluted tube, situated either in deepest portion of corium, or more usually in subcutaneous cellular tissue, and opening on free surface by a spiral duct; the tubes are formed of an external fibro-cellular coat, continuous with superficial layer of corium, and a lining of epithelial cells continuous with epidermis.

What is their estimated number and area?

Nearly two and a half millions, forming an evaporating area of about eight square inches.

Organs of Taste.

Where does the sense of taste reside?

In the mucous membrane of dorsum of tongue, upper portion of pharynx, soft palate, and fauces.

Are there any special taste-organs?

Yes, the *taste-goblets*, flask-like bodies with their bases resting on corium, and a neck opening between epithelial cells. They consist of spindle-shaped, flattened cells externally, with central spindle-cells, whose inner fine terminations are continuous with a terminal nerve fibril; these bodies are found in the circumvallate and some of the fungiform papillæ, and at the side of base of tongue.

What are the special nerves of taste?

For its anterior two-thirds, *chorda tympani of facial,* recognizing acid, saline, and styptic qualities (according to Flint).

For posterior third, *lingual branch of the glosso-pharyngeal,* for perception of alkaline, metallic, sweet, and bitter tastes (Flint).

Organs of Smell.

Where does the sense of smell reside?

In the mucous membrane lining upper portion of nasal fossæ, where the *olfactory nerve filaments* are distributed.

Describe the nasal mucous membrane.

It is also called the *Schneiderian,* or *pituitary* membrane, and is continuous with that of pharynx, Eustachian tube, tympanum,

FIG. 102.



and mastoid cells, and with that of the accessory nasal cavities, the frontal, ethmoidal, and sphenoidal sinuses and antrum, also through lachrymo-nasal duct, with conjunctiva. Its epithelium is ciliated columnar below, and simple columnar above; it contains numerous mucous glands, and in the deeper portions of epithelial layer certain nucleated bodies are found, the *olfactory cells of Schultze* with two processes, one running between epithelial cells to free surface, the other inward, to connect with a terminal nerve fibre.

What vessels supply the exterior of the nose?

Lateralis nasi, from facial to alæ of nose; the

Artery of septum, from superior coronary to septum; the

Nasal branches of ophthalmic and infraorbital to dorsum and sides of nose; the *veins* terminate in the facial and ophthalmic.

What are the vessels of the nasal fossæ?

Anterior and posterior ethmoidal from ophthalmic to ethmoidal cells, frontal sinuses, and roof; the

Spheno-palatine and twig from *small meningeal*, from internal maxillary to membrane over spongy bones, meatuses, and septum; the

Alveolar, from internal maxillary to antrum; the

Veins usually accompany the arteries, terminating in facial, ophthalmic, and other trunks.

What are the nerves of the nose?

Branches of *facial*, *infraorbital*, *infratrochlear*, and *nasal branch* of *ophthalmic*.

Describe the nerve-supply to the nasal fossæ.

The *olfactory*[1] is distributed to mucous membrane over upper third of septum, and superior and middle spongy bones; the

Nasal branch of ophthalmic[2] supplies upper anterior part of septum and outer wall; the

Anterior dental branch of superior maxillary supplies inferior turbinated bone and inferior meatus; the

Spheno-palatine[6], by *naso-palatine* branch[7], supplies middle of septum, by *anterior palatine*, the middle and inferior spongy bones.

The *Vidian*[10] supplies the upper back part of septum and superior spongy bones, parts to which are also distributed the upper anterior nasal branches of spheno-palatine ganglion[7].

(For bones forming nasal fossæ, see p. 58.)

The Eye.

Describe the eyeball.

It is a spherical organ, with the segment of a smaller transparent sphere, the cornea, forming its anterior portion, lying in the orbit in a cushion of fat, and held in position by the ocular muscles, the optic nerve, conjunctiva, and eyelids. The following points should be studied, the

Diameters, antero-posterior in adult about ninety-five hundredths of an inch, transverse ninety-two hundredths of an inch, vertical ninety hundredths of an inch.

Anterior and posterior poles, the centres respectively of the cornea and fundus oculi.

Axes of eyeballs, or ocular axes which pass through the poles of each eye and are nearly parallel, *not corresponding to axes of orbits*, which diverge.

Visual axis passes from the yellow spot through the centre of curvature of the refracting media, so that *these axes converge.*

Nodal point, the centre of curvature of refracting media.

Equatorial plane, that passing through centre of eyeball at right angles to the axis, dividing the globe into two segments, an anterior and a posterior hemisphere.

Equator, the line where the just-mentioned plane cuts the surface of eyeball.

Meridional planes, planes coinciding with ocular axis.

Meridians, the lines where meridional planes cut surface of eyeball.

What is the capsule of Tenon?

The *tunica vaginalis oculi*, and consists of a thin membranous sac isolating the eyeball and allowing free movement, arising from optic foramen behind to become lost upon sclerotic in front, having a visceral layer investing posterior portion of globe, and a parietal layer lining the fatty cushion in which eye rests, both layers being lined by flattened endothelial cells: the lymphatic vessels of the sclerotic open into it, the cavity being a lymph space also communicating with the lymph-spaces of the optic nerve sheath; the ocular muscles pierce this capsule: that portion of the capsule behind the point of passage of the tendons is sometimes called the

Capsule of Bonnet, while to that in front the name of

Capsule of Tenon has been restricted; this is an unnecessary refinement.

What are the tunics or coats of the eye?

The *sclerotic*[2] *and cornea*[2], the most external, protective tunic.

Choroid, iris, and ciliary processes, the middle or vascular tunic.

Retina, the innermost, nervous tunic.

What are the humors of the eye?

The refracting media, three in number, viz., the

Aqueous[3], filling anterior chamber.

Crystalline[4], or crystalline lens.

Vitreous[5], filling posterior four-fifths of the globe.

Sclerotic and Cornea.

Describe the sclerotic.

It is a firm, unyielding fibrous membrane maintaining the form of the posterior five-sixths of the globe, thicker behind than in front, with smooth white external surface except where tendons are attached, covered in front by conjunctival membrane, and having an inner brown-stained surface grooved for ciliary nerves. Posteriorly to the nasal side it is pierced by the optic nerve with whose sheath it is continuous, the sclerotic at this point forming a thin

Lamina cribrosa through whose openings the nerve filaments pass, the most central orifice, called the *porus opticus*, transmitting the arteria centralis retinæ. Surrounding the cribriform lamella are numerous small openings for the ciliary vessels and nerves; the venæ vorticosæ emerge midway between margin of cornea and entrance of optic nerve, and the anterior ciliary vessels around the corneal border. The sclerotic is formed of white fibrous tissue, mingled with fine elastic fibres and fusiform nucleated cells. The following points should be noted; the

Thickness, about one-twenty-fifth of an inch posteriorly, one-sixtieth of an inch anteriorly;

Lamina fusca, the fine cellular tissue connecting sclerotic with choroid; the

Subconjunctival tissue, loose areolar tissue binding conjunctiva to sclerotic; the

Arteries are few, come from *ciliary*, and form a coarse network; the

Anterior vascular zone, surrounding cornea, formed by subconjunctival branches of anterior ciliary vessels; the

Nerves, whose existence is very doubtful.

Describe the cornea.

It is the transparent, projecting, anterior sixth of the external tunic. It is not quite circular, being a little broader in the transverse than the vertical diameter, and is set in sclerotic somewhat like a watch-crystal in its case: its thickness, one twenty-second of an inch to one thirty-second of an inch, is nearly same throughout except toward outer margin where it is thinner: it is *non-vascular*, the capillary vessels forming loops at its circumference: the *nerves* are numerous, estimated at from twenty-four to forty-five, and are derived from the ciliary nerves.

FIG. 103.



Section of eyeball.

Describe the structure of the cornea.

It consists from without inward, of the following layers:

Layer of conjunctival epithelium, the deepest columnar, next polyhedral, most superficial three or four layers of scaly cells; the

Anterior elastic lamina or membrane of Bowman, one two-thousandth to one-twelve-hundredths of an inch thick, forming a thin, firm, elastic layer consisting of extremely closely interwoven fibrils similar to those of corneal tissue proper, but with no corneal corpuscles; the

True corneal tissue is fibrous, tough, unyielding, continuous with sclerotic and composed of sixty, superimposed, flattened lamellæ made up of fibrillar connective-tissue continuous with fibres of sclerotic: the layers are held together by a cement substance in which are *corneal spaces*, stellate in form, communicating with adjacent spaces by processes, each space containing a similarly shaped but smaller *corneal corpuscle* which does not entirely fill it; the

Posterior elastic lamina (membrane of Descemet or Demours) is a structureless, elastic, brittle, extremely thin membrane, not rendered opaque by water, alcohol, or acids: it chief peculiarity is the tendency to roll upon itself with the attached surface innermost, when separated from cornea proper: at its circumference it breaks up into bundles of fibres joining the *ligamentum pectinatum iridis*; the

Posterior epithelial layer is a single layer of polygonal transparent nucleated cells.

Describe the choroid.

It is a thin, dark-brown, vascular membrane investing the posterior five-sixths of globe terminating in front at ciliary ligament, there bending inward to form the ciliary processes: it is connected externally with sclerotic by the *lamina fusca*, and consists of two layers, the

External layer is formed of the larger branches of the ciliary arteries and the venæ vorticosæf, interspersed with stellate pigment cells by the union of whose processes a delicate network is formed; externally the choroid is covered by a layer of elastic fibres, the *membrana suprachoroidea*, coated externally with endothelial cells; the

Internal layer or *membrana Ruyschiana* consists of a fine capillary plexus formed by the short ciliary vessels, which is separated from pigmentary layers of retina by the

Vitreous membrane, a thin, structureless layer.

The arteries of the choroid are the short, posterior ciliary, and recurrent branches from the long posterior and anterior ciliary; the

Veins, form four to five *venæ vorticosæ*[f] (Fig. 104), which pierce the sclerotic midway between corneal margin and optic nerve; the

Nerves are the long and short ciliary nerves.

What are the ciliary processes?

Sixty to eighty folds formed by plaiting and folding inward of choroid at its anterior margin, which are received between corresponding folds of the suspensory ligament of the lens; they consist of large and small processes, irregularly alternating, the former about one-tenth of an inch long.

What is the ciliary body?

It is that portion of the middle tunic between the choroid and iris formed of the ciliary muscle and ciliary processes.

Describe the ciliary muscle[b].

A grayish circular band of unstriped muscular fibre about one-eighth of an inch broad, thickest in front, thinnest behind, consisting of radiating and circular fibres, the former, the more numerous, arising from junction of cornea and sclerotic to pass backward to choroid opposite the ciliary processes; the latter are internal, and pursue a circular course around the insertion of the iris; they are called the *"ring-muscle" of Müller* and were formerly described as the *ciliary ligament.*

Describe the circulus arteriosus major and minor.

The former is an anastomotic ring formed by long ciliary arteries in the ciliary muscle; the latter, a second anastomotic circle formed by transverse offsets from the converging branches passing forward from the circulus major.

Describe the iris[c] (Fig. 104).

It is a thin, circular, perforated contractile curtain, suspended behind the cornea in the aqueous humor in front of the crystalline lens, forming the anterior portion of the middle ocular tunic, and is formed of radiating and circular muscular fibres, and of a fibrous stroma. Anteriorly it is covered by a layer of polyhedral cells resting on a fine basement membrane, being continuous with the membrane of Descemet. It presents for examination the

Pupil, the nearly central opening in iris placed a little to nasal side of centre; the

Ligamentum pectinatum iridis, a reticular tissue, connecting iris with cornea, derived from membrane of Descemet which sends fibrous prolongations to iris and sclero-corneal junction; the

Fig. 104.



Spaces of Fontana, the intervals between the reticulated fibres of the ligamentum pectinatum at outer angle of anterior chamber, which communicate with a larger space in the sclerotic close to sclero-corneal junction; the

Canal of Schlemm or Sinus circularis iridis, lined with endothelium, a venous sinus, containing two or three plexiform veins receiving blood from the sclerotic and ciliary plexuses and communicating with anterior ciliary veins; the

Uvea, a layer of purplish-hued pigment-cells on posterior surface of iris continuous with pigment layer of ciliary processes; the

Sphincter pupillæ, the narrow band of circular muscular fibres surrounding pupil on its posterior surface, about one-thirtieth of an inch wide, supplied by third nerve through ophthalmic ganglion; the

Dilator pupillæ, consisting of the radiating muscular fibres converging from circumference of iris toward pupillary margin, where they blend with the circular fibres; it is supplied by sympathetic fibres from ophthalmic ganglion; the

Membrana pupillaris, a delicate, transparent, vascular membrane which occluded the pupil in the fœtus, usually disappearing about eighth fœtal month, but occasionally persisting; it is nourished by many small vessels continued from margin of iris to those on front part of lens capsule; the

Arteries, are branches of long and anterior ciliary forming the circulis iridis major and minor (see p. 296); the

Veins, empty into those of ciliary processes and anterior ciliary veins; the

Nerves, are branches of third, fifth, and sympathetic, through long and short ciliary nerves, the third going to sphincter pupillæ (circular fibres), the sympathetic to dilator pupillæ (radiating fibres), and the fifth supplying common sensation.

What is the retina?

The innermost ocular tunic, forming a delicate, grayish, transparent nervous membrane, thicker behind than in front, terminating nearly as far forward as the ciliary body by a ragged margin, the *ora serrata*, but its fibrous stroma covered by the pigment layer passes forward to iris as the *pars ciliaris retinæ*. It presents the

Macula lutea or yellow spot of *Sömmering*, a round elevated spot exactly in centre of retina posteriorly, which corresponds to axis of eye; here vision is most perfect, the retina being thin, crowded with nerve elements, but destitute of rods and the nerve-fibre layer; the

Fovea centralis is a minute central depression at summit of yellow spot; the

Optic papilla or disk is the point of entrance of optic nerve, centrally pierced by arteria centralis retinæ, lying about one-tenth of an inch to inner side of yellow spot; it is the only portion of retina where the sense of vision is wanting, and is therefore called the *blind spot*; the

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

5. *Inner nuclear layer*, containing (1) oval nuclei with two processes, one passing inward to inner molecular layer, believed to become continuous with processes of ganglion cells, and another passing into outer molecular layer, these bifurcating and (according to some) communicating with rod and cone fibres; (2) nucleated cells without branches; (3) cells connected with fibres of Müller.

6. *Outer molecular layer*, is thinner, resembles the inner molecular layer, but contains branched stellate cells—probably ganglion cells.

7. *Outer nuclear layer*, consists of (1) *rod-granules*, transversely striated cells, with an external fine process connected with a single rod of Jacobs's membrane, and an internal prolongation which enlarges, then breaks up into minute fibrils entering the outer molecular layer; and (2) *cone-granules*, closely connected with cones of Jacobs's membrane, with a thick process passing inward, becoming bulbous and terminating by numerous fibrillæ, which enter outer molecular layer.

8. *External limiting membrane*, a membrane formed of retinal connective tissue, perforated by numerous openings.

9. ~~*Jacobs's membrane or rods and cones*,~~ composed of *rods* arranged perpendicularly to surface, each composed of an outer and inner portion joined by cement substance; and *cones*, with apices directed toward choroid, formed of two portions and like rods their outer segments are transversely striated; this is probably the perceptive layer of retina.

10. *Pigmentary layer*, formerly considered part of choroid, consisting of a single layer of hexagonal, pigmented epithelial cells.

The connective tissue is modified to form a fenestrated, sponge-like structure the *membrane of Müller*.

The Humors.

What is the aqueous humor?

A small amount of clear alkaline fluid composed of water 96.7, albumen 0.1, sodium chloride and extractive 3.2, filling anterior and posterior chambers of the eye.

What are the anterior and posterior chambers of the eye?

The anterior chamber is a space filled with aqueous humor bounded in front by cornea, behind by front of iris, and communicating through pupil with the

Posterior chamber, also filled with aqueous humor, which is now thought to be only a narrow chink between peripheral portion of iris, the suspensory ligament, and ciliary processes.

Describe the vitreous humor.

This forms about four-fifths of entire globe, is transparent, of the consistence of thin jelly, and is composed of an albuminous saline fluid enclosed, except where hollowed out for crystalline lens, in a delicate *hyoloid membrane*, beneath which are small, granular, nucleated cells; it possesses neither vessels nor nerves, and is hollowed out in front where the crystalline lens reposes. Running from entrance of optic nerve to posterior surface of crystalline lens is the *canal of Stilling or hyaloid canal*, filled with fluid and lined with a process of the hyaloid membrane.

Describe the crystalline lens with its ligaments, etc.

It is a transparent double convex body, more convex posteriorly than in front, enclosed in a *capsule*, and is lodged in a depression of the vitreous humor, where it is retained by its suspensory ligament. It lies immediately behind the pupil surrounded and slightly overlapped by the ciliary processes, measuring about one-third of an inch transversely by one-fourth of an inch antero-posteriorly, and is composed of water 60, soluble albuminous matter 35, insoluble albuminous matter 2.5, fat and cholesterine 2 per cent.; its

Capsule, is transparent, very elastic, and brittle, measuring one-two thousandth of inch in front, one-six thousandth of an inch behind, and is attached anteriorly to lens by a single layer of polygonal nucleated cells, which absorb fluid post-mortem, and breaking down from the *liquor Morgagni*; the

Suspensory ligament or zonula of Zinn, is a thin transparent membrane, containing longitudinal elastic fibres, connecting anterior margin of vitreous with anterior surface of lens near its circumference, presenting upon its outer surface a number of folds between which fit the similar projections of the ciliary processes; the

Canal of Petit is about one-tenth of an inch wide, passing around circumference of lens, bounded in front by suspensory ligament, behind by vitreous humor, its base being formed by capsule of lens; when inflated it becomes sacculated.

What is the structure of the crystalline lens?

It is formed of numerous laminæ, composed of hexagonal fibres one-five thousandth of an inch broad, with serrated margins, those of contiguous fibres accurately fitting one another; the lens is also divisible after boiling or immersion in alcohol into three triangular segments with their bases outward. The central harder portion is called the *nucleus*, the peripheral portions the *cortex*; it possesses neither vessels nor nerves.

Describe the muscles of the eyeball.

Rectus superior[4]: *origin*, upper margin of optic foramen and sheath of the optic nerve; *insertion*, sclerotic coat three to four lines from cornea; *action*, turns eyeball upward; *nerve*, third cranial.

Rectus inferior[5]: *origin*, ligament of Zinn; *insertion*, sclerotic three to four lines from cornea; *action*, turns eye downward; *nerve*, third cranial.

Rectus internus: *origin*, ligament of Zinn; *insertion*, sclerotic three to four lines from cornea; *action*, turns eye inward; *nerve*, third cranial.

Rectus externus[6]: *origin*, by two heads, the upper[8] from the outer margin of optic foramen, the lower[7] from ligament of Zinn, and a pointed bony process at the lower margin of the sphenoidal fissure; *insertion*, sclerotic, as other recti; *action*, turns eye outward; *nerve*, sixth cranial (abducens): between the two heads pass the ophthalmic vein, the third, nasal branch of the fifth, and the sixth cranial nerves.

Obliquus superior[2]: *origin*, one line above the inner margin of the optic foramen, terminating in a rounded tendon which plays through a fibro-cartilaginous ring or pulley[3] beneath the internal angular process of the frontal bone, whence it passes beneath the superior rectus; *insertion*, sclerotic between superior and external recti, midway between the cornea and entrance of the optic nerve; *action,* rotates eye on its antero-posterior axis; *nerve,* fourth cranial (patheticus).

Obliquus inferior[10]: *origin,* orbital plate of superior maxillary; *insertion,* sclerotic between superior and external recti; *action,* rotates eyeball on antero-posterior axis; *nerve,* third cranial.

(For muscles of lids see pp. 116, 117.)

FIG. 106.



Recapitulate the nerve supply of the ocular muscles.

Third cranial, or motor oculi, supplies superior, internal and inferior recti, and inferior oblique muscles.

Fourth cranial or patheticus, supplies superior oblique muscle.

Sixth cranial or abducens, supplies external rectus muscle.

Describe the vascular supply of the eyeball and its appendages.

(All branches of the vessels not supplying these parts *will be omitted,* having been already described on page 161.)

Ophthalmic, arising from cavernous portion of internal carotid, entering orbit by optic foramen, giving off the

Lachrymal to lachrymal gland, conjunctiva, and upper eyelid, inosculating with palpebral arteries.

Supra-orbital, supplies levator palpebræ and superior rectus muscles.

Superior and inferior palpebral, supply eyelids.

Nasal, supplies lachrymal sac.

Short ciliary, twelve to fifteen in number, penetrate sclerotic around optic nerve entrance to supply choroid and ciliary processes.

Long ciliary, two in number, penetrate sclerotic, pass forward between it and choroid to supply iris, forming *circulus iridis major* near ciliary margin, and *circulus iridis minor,* near margin of pupil.

Anterior ciliary, spring from muscular branches, perforate sclerotic, and join circulus iridis major.

Arteria centralis retinæ, obliquely traverses optic nerve to be distributed to retina.

Muscular branches, one superior, one inferior, to ocular muscles.

Anterior cerebral, from internal carotid, gives twigs to optic nerve.

Infra-orbital, from internal maxillary, supplies inferior rectus and inferior oblique muscles and lachrymal gland.

What are the chief lymph-spaces of the eyeball?

The *perichoroidal space,* between choroid and sclerotic.

Cavity of capsule of Tenon, between eyeball and capsule.

Vaginal space, between optic nerve and sheath.

What veins return the blood from the eye?

The *superior and inferior ophthalmic* emptying into the cavernous sinus; they also freely anastomose with the facial veins.

Name the nerves of the eye.

Optic or second cranial, nerve of special sense: motor nerves, the *third, fourth, sixth,* and *sympathetic filaments.*

The *Ophthalmic division of fifth,* supplies general sensation by

Lachrymal, to same-named gland, conjunctiva, and skin of upper eyelid; also the following branches:

Frontal { *Supra-trochlear* / *Infra-orbital* ; and Nasal { *Ganglionic.* / *Long ciliary.* / *Infra-trochlear.*

The *sympathetic* arises from cavernous and carotid plexus receiving communicating filaments indirectly from spinal nerves, and sending branches to third, fourth, fifth, and sixth nerves; the dilator fibres (radiating) of iris, ciliary ganglion, muscles of orbit and lids, with walls of arteries, receive their sympathetic nerve supply from this source; the

Short ciliary, numbering about ten, arise from ciliary (ophthalmic) ganglion, and pierce the sclerotic posteriorly, to ramify in sheath of optic nerve, choroid, ciliary muscle, iris, and cornea; the

Ascending branches of the spheno-palatine (Meckel's) ganglion, reaching orbit by spheno-maxillary fissure, usually considered to supply only periosteum, are believed by some to go to optic and sixth nerves and ciliary ganglion.

Give a brief description of optic tracts.

The deep origin of the nerve fibres of these tracts are the optic thalami, corpora geniculata, and nates of corpora quadrigemina, which, forming two flattened bands, wind around crus cerebri, to which they are slightly attached, as well as to lamina cinerea and tuber cinereum, and joining in front of latter they form the *optic commissure* (see page 200 for illustration and more elaborate description).

What course do the nerve fibres pursue in the optic commissure or chiasm?

The most numerous, the *decussating* fibres, connect the right hemisphere with the left retina, and the left hemisphere with the right retina, occupying the centre of the commissure.

Other fibres (*longitudinal, cerebro-retinal*) connect each hemisphere with the retina of its own side.

Others (*inter-cerebral*) pass posteriorly, connecting the two optic tracts.

Others (*inter-retinal*) pass anteriorly, connecting the two retinæ.

What are the origin and course of the optic nerves?

They arise on either side from the optic commissure, each passes into the orbit through optic foramen with central artery of retina, which pierces it, and enter eyeball through lamina cribrosa one-tenth inch to nasal side of globe, to aid in forming retina; these nerves are surrounded with a sheath derived from the dura mater between which and nerve is a lymph space communicating with subarachnoid space.

APPENDAGES OF EYE.

(Tutamina Oculi.)

Name these.

Eyebrows, or *supercilia*,
Eyelids, or *palpebræ*,
Conjunctiva[a], a mucous membrane,
Lachrymal gland[1] and *ducts*[2],
Lachrymal sac,
Lachrymo-nasal duct.

What are the eyebrows?

The arched eminences of skin surmounting upper margins of orbits on each side, from which grow several rows of short, obliquely placed hairs; by the attached muscles the eyebrows can partially shut off light from eyes.

What are the eyelids?

Two movable curtains protecting by their closure the eyes from injury; the upper lid is the larger and more movable, having its own elevator muscle; the

Palpebral fissure is the elliptical space between the margins of lids when opened, the internal and external angles being called, respectively, the *internal* and *external canthus*, the former being prolonged inward, leaving a triangular space between the lids, the *lacus lachrymalis*, at the commencement of which is a small elevation on each lid, the *lachrymal papilla*[3], whose summit is pierced by a minute opening, the *punctum lachrymale*[3], the commencement of the lachrymal canal.

What structures form the eyelids?

Externally the skin, beneath which is much loose areolar tissue, separating it from the fibres of the *orbicularis palpebrarum muscle*. The lids retain their shape by means of the so-called *tarsal carti-*

lages—in reality dense connective tissue *without* cartilage cells—that for the upper lid being semilunar and the larger, that for lower lid elliptical; these structures are connected at their orbital margins with the circumference of the orbit by a fibrous membrane, the *tarsal ligaments*, while in addition a rounded fibrous cord, the *tendo-oculi*, passes between the inner angle of each and the nasal process of superior maxillary bone.

Imbedded in the cartilages are numerous sebaceous glands, the *Meibomian glands*, discharging their secretion upon the free edge of lids, preventing their adhesion; they number thirty or forty in the upper lid, fewer in the lower lid.

Attached to the free margins of lids is a double or triple row of stiff hairs, curved in each lid so that their convexities meet, preventing interlacing, these are the *cilia*, or *eyelashes*; the inner surfaces of lids are lined by a mucous membrane, the *conjunctiva*.

Their *vascular supply* is from the *palpebral* branches of ophthalmic artery anastomosing with contiguous branches of facial.

The *nerves* are the third, seventh, and sympathetic to muscles, the ophthalmic branch of fifth to skin and conjunctiva.

Describe the conjunctiva.

It is a mucous membrane lining the inner surfaces of the eyelids and reflected thence upon anterior segment of sclerotic, its epithelial layer passing over cornea; the

Palpebral conjunctiva is thick, opaque, vascular, and covered with papillæ, containing numerous glands; at the inner angle of the eye it forms a semilunar fold, the *plica semilunaris*[4] (Fig. 108). The folds formed by the passage of the conjunctiva from lids to eyeball are called the *superior* and *inferior palpebral folds*, the former being the deeper; they contain numerous convoluted mucous glands; the

Ocular conjunctiva is thin, transparent, possesses few vessels in health, and is loosely attached to globe by the subconjunctival areolar tissue.

What is the caruncula lachrymalis[5] (Fig. 108)?

It is a small, conical, reddish body, at inner canthus, occupying the space called the lacus lachrymalis, and is formed of a group of follicles, resembling the Meibomian glands, covered with conjunctiva; from the caruncle project a few slender hairs.

Describe the lachrymal apparatus.

It consists of the lachrymal gland with its ducts, the canaliculi, the lachrymal sac, and the nasal duct. The

Lachrymal gland[1] (Fig. 107) is an oval, compound racemose gland of the size and shape of an almond, lodged in a depression at upper, outer portion of orbit, its concave under surface resting upon the globe of eye, the conjunctiva, superior and external recti muscles intervening; it is held in contact with orbital periosteum by a few fibrous bands. It has six or seven

Fig. 107.



Fig. 108.



Ducts[2] (Fig. 107), opening by minute orifices on upper outer half of palpebral conjunctiva near superior retro-tarsal fold. The secretion, or the

Tears, are taken up by the puncta, thence passing through canaliculi into lachrymal sac and nasal duct, thus reaching inferior meatus of nose.

Describe the canaliculi.

They are two minute canals[1] about one-half a line in diameter, commencing at the puncta, the superior passing upward, then bending acutely to pass inward to lachrymal sac, the inferior passing downward, then upward and inward; they are lined with mucous membrane continuous with conjunctiva, and are one-third of an inch long.

What is the lachrymal sac[2] (Fig. 108)?

It is the flattened, ovoidal, sacciform dilatation of upper part of nasal duct, lodged in the groove formed by lachrymal bone and nasal process of superior maxillary; its walls are formed of fibrous elastic tissue covered in by tensor tarsi muscle and a fibrous expansion from tendo-oculi attached to ridge of lachrymal bone, while internally it is lined with mucous membrane.

Describe the nasal duct[3].

It is a membranous canal about three-fourths of an inch long extending from the termination of lachrymal sac through osseous nasal duct to inferior meatus of nose, passing in a direction downward, backward, and slightly outward, its lumen being narrowest about the midpoint; externally, it is composed of fibro-areolar tissue; internally, of mucous membrane continuous with that of nose and lachrymal sac, the epithelial coating of both being of ciliated epithelium, that of the canaliculi of scaly cells.

What is the valve of Hasner?

An imperfect valve of mucous membrane guarding the somewhat expanded terminal opening of the duct into the inferior meatus of nose.

The Ear.

Where is the auditory apparatus lodged?

In the petrous and mastoid portions of the temporal bone.

Enumerate the divisions.

- The *External ear*, comprising the
  - *Auricle*, or *pinna*,
  - *External auditory canal.*
- *Middle ear*, or *tympanum*, comprising
  - *Membrana tympani,*
  - *Cavity of tympanum,*
  - *Mastoid cells,*
  - *Eustachian tube,*
- *Internal ear*, or *labyrinth*, comprising
  - *Vestibule,*
  - *Semicircular canals,*
  - *Cochlea,*
  - *Auditory nerve.*

The External Ear.

What is the auricle?

Its foundation is an expanded layer of fibro-cartilage, so disposed in ridges as to concentrate and direct the waves of sound into the external auditory canal, to which it is attached; the cartilage is deficient at certain points where it is connected by fibrous tissue; it is covered with perichondrium, outside of which is thin, firmly adherent skin, containing sweat and sebaceous glands, and provided with short downy hairs. The various ridges and depressions are as follows: the

Concha, the deep cavity leading into meatus; the

Tragus, the pointed prominence in front of concha projecting back over meatus, bearing on its under surface tufts of hairs; the

Antitragus, a small tubercle opposite tragus, separated by a deep notch, the *incisura intertragica*; the

Helix, the external prominent margin of auricle; the

Antihelix, a parallel prominence anterior to former, bifurcating above to form the

Fossa of the antihelix; the

Fossa of helix, the narrow, curved depression between the helix and antihelix.

What is the lobe or lobule of the ear?

The inferior, soft pendulous portion, formed of fat and connective tissue enclosed by integument.

Describe the muscles of the external ear.

The attolens aurem, attrahens aurem, and retrahens aurem, have been described on page 116.

The muscles of the auricle, but slightly developed in man, are the

Helicis major: a narrow, vertical band on anterior border of helix;

Helicis minor: an oblique band at root of helix;

Tragicus. a short, vertical band on outer surface of tragus;

Antitragicus: stretching from outer part of antitragus to lower part of helix;

Transversus auriculæ: radiating from posterior surface of convexity of concha to prominence caused by groove of helix;

Obliquus auris: a few fibres passing from upper back part of concha to convexity immediately above.

Give the vascular and nervous supply to the auricle.

The arteries are *Posterior auricular*, from external carotid.
Anterior auricular, branch of temporal.
Auricular, branch of occipital.

The *veins* accompany the corresponding arteries.

The nerves are, *Auricularis magnus*, from cervical plexus.
Posterior auricular, from facial.
Auricular branch (Arnold's) of pneumogastric.
Auriculo-temporal, from inferior maxillary division of fifth nerve.
Occipitalis major and *minor*.

What is the external auditory canal?

It is an osseo-cartilaginous, oval, cylindrical canal, with ~~greatest diameter vertical at external orifice~~, but reversed at tympanic end; it is narrowest at middle. About one and a quarter inches long, it is directed obliquely forward and inward, and slightly curved with convexity upward, and is lined by thin adherent skin, containing hair follicles, sebaceous and ceruminous glands in its cartilaginous portion. A narrow groove at the tympanic extremity, for insertion of membrana tympani, is sometimes called the *sulcus tympanicus*, interrupted above by the *notch of Rivinus*.

What are the arteries and nerves?

The *arteries* are branches of posterior auricular, internal maxillary and temporal.

The *nerves* come chiefly from auriculo-temporal branch of superior maxillary division of fifth nerve.

The Middle Ear or Tympanum.

Describe the membrana tympani.

It forms the outer wall of the tympanum.
lucent membrane placed obliquely at

tory meatus, with its anterior, shallow funnel-shaped surface facing downward, forward, and outward; at its upper anterior border is a white, pointed *tubercle,* formed by the short process of the malleus, while a yellowish-white stripe passing from this downward and backward toward the centre, indicates the handle of the malleus. During life, when illuminated, the membrana tympani presents a triangular *light spot* or "*cone of light,*" having its apex at end of malleus handle, whence it spreads downward and forward, and a darker central portion, the *umbo*, or shadow. The upper anterior part bridging a small notch in the bony ring to which the membrane is attached (the notch of Rivinus), is laxer, consisting of loose connective tissue, vessels, and nerves, covered with skin and mucous membrane, and has received the name of the *membrana flaccida*, or *Shrapnell's membrane.*

Of what tissues is the membrana tympani composed?

Of the skin, which is derived from the lining of the meatus; of a fibrous layer, some of whose fibres radiate from near the centre to the circumference, others form a dense ring around the attached margin; and of mucous membrane, derived from that lining the tympanum. The handle of the malleus passes between the inner and middle layers—according to von Tröltsch, it is received *between* the circular and radiating fibres of the middle coat; an *anterior and a posterior pouch* have also been described upon the posterior surface by von Tröltsch, opening downward, formed by a "duplicature of the membrana tympani extending from its border forward to the manubrium."

Describe the arterial supply of the membrana tympani.

The *deep auricular* branch of internal maxillary supplies the external layers forming a plexus, which communicates at the periphery with one in the mucous membrane formed from

Tympanic branches of internal maxillary and internal carotid arteries, and by the

Vidian, from internal maxillary, and

Stylo-mastoid, from posterior auricular.

Mention the nerves supplying the membrana tympani.

To the external layer run filaments from the *superficial temporal* branch of the fifth, while the mucous layer is supplied by the *tympanic* plexus.

The Tympanum.

Describe it.

It is an irregular cavity, measuring about one-half of an inch antero-posteriorly, one-third of an inch vertically, and one-fifth of an inch transversely, situated in the petrous bone, compressed from without inward, being placed above the jugular fossa, having the carotid canal in front, the mastoid cells behind, the external meatus externally, and the labyrinth internally; it communicates with the pharynx by the Eustachian tube, and presents for examination the following points; the

Roof, a very thin plate of bone, indicated on cranial surface by a depression on posterior surface of petrous portion of temporal bone; the

Floor, narrow, corresponding to jugular fossa, and presenting near inner wall a small *foramen for Jacobson's nerve*; the

Outer wall, the membrana tympani and bony ring to which it is attached, presenting three small orifices, the

Iter chordæ posterius, opening in angle of junction between posterior and external walls, just behind membrana tympani on a level with its centre, for entrance of chorda tympani nerve; the

Glaserian fissure, just above and in front of bony ring giving origin to membrana tympani, for passage of laxator tympani muscle and some tympanic vessels, and the lodgement of the long process of malleus; the

Iter chordæ anterius or *canal of Huguier*, opening just above preceding, for *escape* of chorda tympani nerve; the

Inner wall is vertical, looking directly outward, and presents the

Fenestra ovalis or oval window, a kidney-shaped opening leading into vestibule, closed by base of stapes with its ligament; the

Fenestra rotunda or round window, below, at bottom of a funnel-shaped depression, opening into scala tympani of cochlea, closed by the *membrana tympani secundaria*; the

Promontory, a rounded hollow eminence—the first turn of the cochlea—situated between oval and round windows, and presenting three grooves lodging branches of tympanic plexus; the

Eminence of the aquæductus Fallopii, passes above oval window, along inner tympanic wall, to curve behind that opening nearly vertically downward along posterior wall; the

Pyramid, a hollow conical projection, behind the oval window and in front of vertical portion of aquæductus Fallopii, containing the stapedius muscle whose tendon escapes from its summit; a minute canal communicates with the aquæductus Fallopii transmitting the nerve to stapedius muscle; the

Posterior wall, presenting the

Openings of the mastoid cells, one large and several smaller openings. The anterior wall, is wider above than below, corresponding with carotid canal, the thin bony partition being perforated by tympanic branch of carotid; it presents the

Orifice of the canal for the tensor tympani muscle, which is situated above, close to aquæductus Fallopii, on the summit of a small conical eminence, *the anterior pyramid*; the

Opening of Eustachian tube, is immediately below preceding, separated by a thin horizontal bony plate, the *processus cochleariformis.*

What are the ossicles of the tympanum?

Three bones forming a movable chain, passing between membrana tympani and oval window, called the

FIG. 109.



Malleus or hammer, consisting of an oval *head* articulating with incus, a neck, a *manubrium* or handle, affording attachment near its root to tensor tympani muscle, a *short process* coming in contact with membrana tympani, and a *processus gracilis* or long process, lodged in the Glaserian fissure, and giving attachment to so-called laxator tympani muscle; the

Incus or anvil, having a body articulating with malleus, a *long process* terminating in a rounded end, the *os orbiculare*, which articulates with head of stapes, and a *short process* attached to margin of opening into mastoid cells; the

Stapes or stirrup, consists of a *head* articulating with os orbiculare, a *neck* receiving insertion of stapedius muscle, two branches or *crura* joining the oval *base*, which latter is connected with margins of oval window by ligamentous fibres.

Describe the ligaments of the ossicles.

The *suspensory ligament of the malleus*, passing between tympanic roof and head of malleus: the

Posterior ligament of malleus, or laxator tympani minor muscle of Sömmering, passing from upper back part of external meatus to handle of malleus and processus brevis; the

Posterior ligament of the incus, passing between posterior tympanic wall near margin of opening into mastoid cells and end of short process of incus; the

Suspensory ligament of the incus, descending from tympanic roof to incus, near its articulation with malleus; the

Annular ligament of stapes, connecting circumference of its base to margins of oval window; the

Capsular ligaments, around articulations between incus and malleus, and incus and stapes, these joints having their surfaces coated with cartilage, and being provided with a synovial membrane.

Other ligaments have been described under special names, as the anterior and external ligaments of the malleus, and the *obturator ligament* of the stapes, this latter being a membrane filling up opening between crura of stapes.

Describe the muscles of the tympanum.

***Tensor tympani**: origin*, under surface of petrous bone, cartilaginous Eustachian tube, its own osseous canal; *insertion*, handle of malleus near root; *action*, draws membrana tympani inward—*i. e.*, increases *tension*; *nerve*, from otic ganglion.

***Laxator tympani major**: origin*, spinous process sphenoid bone, and cartilaginous Eustachian tube; *insertion*, neck of malleus above processss gracilis; *action*, relaxes membrana tympani; *nerve*, tympanic branch of facial. (Now called *ant. ligament of malleus.*)

***Laxator tympani minor**: origin*, upper back part external meatus; *insertion*, handle of malleus and processus brevis; *action*, relaxes membrana tympani. (Now called *posterior ligament of malleus.*)

Stapedius: origin, from interior of pyramid; *insertion*, neck of stapes; *action*, compresses contents of vestibule; *nerve*, filament of facial.

Describe arterial supply of the tympanum.

The *tympanic branch* of the internal maxillary enters by Glaserian fissure and is distributed to membrana tympani (tympanic branch of carotid also supplies this), anastomosing with *Vidian* and

Stylo-mastoid, from posterior auricular to back part of tympanum and mastoid cells.

Petrosal, from middle meningeal, entering through hiatus Fallopii.

Tympanic, from internal carotid.

Vidian, from internal maxillary.

Branch, from ascending pharyngeal passing up Eustachian tube.

What nerves supply the tympanum?

The *Tympanic branch of the glosso-pharyngeal* (Jacobson's nerve), which pierces floor of tympanum to supply fenestræ and mucous membrane, and aids in forming *tympanic plexus*; the

Tympanic branch of facial, supplying stapedius muscle; the

Branch from otic ganglion, supplying tensor tympani muscle.

(The *chorda tympani from facial*, passing into tympanum by iter chordæ posterius, emerging by iter chordæ anterius, arching across cavity between handle of malleus and long process of incus; it is covered by mucous membrane and gives off no branches.)

Describe the formation of the tympanic plexus.

The tympanic branch of glosso-pharyngeal, or Jacobson's nerve, divides into three branches, lying in grooves on the promontory, one passes forward and downward to join *carotid plexus*, another passes upward to join *greater superficial petrosal nerve* in hiatus Fallopii, the third passes upward and forward through petrous bone close to ganglionic enlargement of *facial*, from which it receives a filament, and becomes the *lesser superficial petrosal nerve*; the tympanic mucous membrane is supplied by branches from this plexus.

Describe the Eustachian tube.

It is about one and one-half inches long, passing downward, forward, and inward, the passage by which the air in the middle ear freely communicates with that in the pharynx, thus permitting free vibration, as the hole in the side of a drum does. It consists of about one-third bone, and two-thirds fibro-cartilage and fibrous tissue; the tympanic orifice measures about one and one-half inch, the oval pharyngeal opening one-third inch.

The *osseous portion*, is one-half inch long, commencing at lower part of anterior tympanic wall, gradually narrowing to terminate at angle of junction of petrous and squamous portions of temporal bone; the

Cartilaginous portion is about one inch long, formed by a triangular plate of elastic fibro-cartilage curved upon itself into a partial tube, but inferiorly the margins are not in contact, the defect being filled by fibrous tissue; the

Mucous membrane is continuous with that of pharynx, and is covered by ciliated epithelium.

Where is the pharyngeal orifice situated?

At upper lateral portion of pharynx behind posterior part of inferior meatus, just above the level of nasal floor.

Has this tube any special muscles?

Yes, the following have been described:

Spheno-salpingo-staphylinis: *origin,* sphenoid bone and cartilaginous tube; *insertion,* convex external border of cartilage; *nerve,* internal pterygoid; *action,* abductor or dilator of tube.

Salpingo-pharyngeus: a thin layer beneath mucous membrane, probably really a fascia.

(The ***Levator palati:*** *origin,* temporal bone and cartilaginous tube; *nerve,* pneumogastric; *insertion,* osseous tube, cartilage, and mucous membrane; *action,* dilates tube transversely.)

Give the arterial supply of the Eustachian tube?

The *ascending pharyngeal,* from external carotid.

Branches of middle meningeal, from internal maxillary.

Branch, from *stylo-mastoid artery.*

The *nerves* are, in addition to those supplying muscles of tube (supra), from the *fifth* and *seventh pairs* and *glosso-pharyngeal.*

The Internal Ear or Labyrinth.

Name its divisions.

The *osseous labyrinth,* consisting of
- *Vestibule*[3] (Fig. 110),
- *Semicircular canals*[5,6,7],
- *Cochlea*[8].

Membranous labyrinth, consisting of
- *Utricle,*
- *Membranous semicircular canals,*
- *Saccule,*
- *Membranous cochlea.*

Internal auditory canal, at bottom of which the auditory nerve enters.

Auditory nerve (eighth cranial, or portio mollis of seventh), with the

Organ of Corti, what the retina is to the eye, viz., the terminal and receptive auditory apparatus.

Describe the internal ear.

It is formed by a series of cavities excavated in the petrous bone, communicating externally with middle ear by the round and oval windows, internally, with internal auditory meatus: within the osseous labyrinth, surrounded by the

Perilymph, is the *membranous labyrinth*—filled with endolymph—upon which the auditory nerve filaments are distributed.

Describe the various subdivisions of the osseous labyrinth.

The *vestibule* is the common central cavity with which all parts of the internal ear communicate, placed behind cochlea in front of semicircular canals, at inner side of tympanum ; it is ovoid, measuring about one-fifth of an inch from before backward, as well as from above downward, and presents the

Fenestra ovalis, or *oval window,* on its outer wall, closed in the fresh state by base of stapes and its ligament; the

Fovea hemispherica[2], a small, circular depression on its inner wall, perforated antero-inferiorly by several minute foramina for branches of auditory nerve, the perforated plate, called the *macula cribrosa*; the

FIG. 110.



Pyramidal eminence, a vertical ridge behind macula cribrosa; the

Aquæductus vestibuli, at back of inner wall, transmitting a small vein, and, according to some, containing a tubular prolongation of lining membrane of vestibule, ending in a blind end in cranial cavity between layers of dura mater; the

Fovea semi-elliptica[1], a transversely-oval depression on roof, separated by pyramidal eminence from fovea hemispherica; the

Orifices of semicircular canals[3], these are five in number, opening behind; the

Apertura scalæ vestibuli cochleæ, a large oval opening, is situated anteriorly.

Describe the semicircular canals[5], [6], [7].

They are three curved bony tubes, each describing the greater part of a circle, of unequal length, of a diameter of about one-twentieth of an inch, each lying at a right angle to the other two, and presenting at one end a dilatation, or *ampulla*, more than double the diameter of the tube; the

Superior semicircular canal[5] is vertical, describes about two-thirds of a circle, passes at right angles to posterior surface of petrous bone, and forms a projection upon its anterior surface; its outer end is its *ampulla*, opening separately into vestibule; its inner undilated extremity joins with that of the

Posterior semicircular canal[6], opening by a common orifice at back of vestibule; this canal is also vertical, is nearly parallel to posterior surface of petrous bone, is the longest, the dilated extremity opening at lower back part of vestibule; the

External, or *horizontal semicircular canal*[7], is the shortest, is directed outward and backward, is ampullated, and opens into vestibule just above oval window, its other extremity by a separate orifice at upper back part of vestibule.

The *cochlea*[8] is conical, somewhat resembles a snail-shell, forms the anterior part of the labyrinth, is placed almost horizontally in front of the vestibule, its apex directed toward the upper front part of tympanic wall, and its base corresponding to anterior perforated depression at bottom of internal auditory meatus; it presents for examination the

Modiolus, a central, conical axis, perforated by numerous canals for filaments of cochlear branch of auditory nerve, its apex terminating in a delicate expanded lamella, like a funnel divided longitudinally, called the *infundibulum*; extending from base to apex is a canal, the *canalis centralis modioli*, for a small nerve and artery; making two and one-half spiral turns around the modiolus is the

FIG. 111.



Spiral canal[8], about one-tenth of an inch in diameter, and one and a half inches long, narrowing from base to apex, there forming the *cupola*, where the scala tympani and scala vestibuli, formed by the lamina spiralis, communicate by the *helicotrema.*

The *cochlea* presents three openings, viz., the round window, communicating with tympanum, the oval one opening into vestibule, and that of the aquæductus cochleæ[9]; projecting from the modiolus is the

Lamina spiralis ossea[l], a process formed of two bony lamellæ, between which are numerous canals for nerve fibres, defective in last half turn of cochlea leaving an aperture, the helicotrema, and terminating by the hook-like hamular process[h]; winding around the modiolus at the point of attachment of the lamina spiralis is the *canalis spiralis modioli*, lodging the enlargement of the cochlear nerve containing ganglion cells, called the *ganglion spirale*, whence come the nerve branches to the organ of Corti.

What is the scala vestibuli[s. v.] (Fig. 111)?

It is that portion of the canal of the cochlea above the lamina spiralis and membrane of Reissner, communicating with vestibule below and scala tympani above, by the helicotrema; it is filled with perilymph.

Describe the scala tympani[s, t].

It is that portion of the spiral canal below the lamina spiralis[b, s] and membranous cochlea, terminating below inferiorly at the round window—which is closed by membrane, the *membrana tympani secundaria*—and communicating above with scala vestibuli at helicotrema; it is filled with perilymph.

What is the aquæductus cochleæ[d]?

A small canal transmitting a vein from cochlea to jugular vein, commencing at lower wall of scala tympani, and ending to inner side of carotid canal close to posterior surface of petrous bone.

Describe and name the subdivisions of the membranous labyrinth.

It consists of a series of closed membranous sacs containing *endolymph*; its various parts are called

The *vestibule*, consisting of the $\left\{\begin{array}{l}\textit{Utricle,}\\ \textit{Saccule,}\end{array}\right\}$ two membranous sacs, with the former of which communicate three

Membranous semicircular canals.

The utricle and saccule are separate, but are indirectly connected by a minute canal passing from saccule to the similar one from utricle, passing along aquæductus vestibuli to end by a blind pouch near posterior surface of petrous bone; the saccule communicates with scala media (or membranous cochlea) by the *canalis reuniens*. The endolymph in all parts is thus seen to communicate freely.

Describe the utricle.

It is an oblong, laterally compressed sac, filled with endolymph, placed in upper back part of vestibule in fovea semielliptica, communicating behind with semicircular canals by five openings, and has distributed, chiefly at one part of its walls, numerous branches of the auditory nerve, at which point is a round mass of minute crystals of calcium carbonate, bound together by delicate fibrous tissue forming the *otoliths*, *otoconia*, or *ear-stones*; the thickening of

the wall both of utricle and saccule, where the nerves penetrate, is sometimes called the *macula acoustica.*

FIG. 112.



What are the membranous semicircular canals?

They are three canals of the same shape but smaller than the containing osseous canals, to which they are fastened by numerous fibrous bands, conveying the auditory nerve filaments to the ampullæ; like the osseous canals they have five orifices which open into the utricle.

Describe the saccule.

FIG. 113.



It is a globular sac, smaller than utricle, to which it is attached at one point, and lies in fovea hemispherica; it is surrounded with perilymph, and indirectly communicates by a short canal with utricle by means of the blind canal passing along aquæductus vestibuli to terminate near posterior surface of petrous bone; and with membranous cochlea or *scala media by canalis reuniens*, a small passage connecting the two cavities.

What is the membranous cochlea^{cc} (Fig. 113).

More usually called *scala media* or *ductus cochlearis* (Fig. 113) it begins by a blind extremity at lower anterior portion of vestibule, enters cochlea, where it forms in cross section a triangular canal, its base being outer bony walls of cochlea, its lower wall the *membrana basilaris*^{b}, its upper the *membrane of Reissner*^{r}; it is filled with endolymph and contains the *organ of Corti*^{co}, covered by the delicate *membrana tectoria*^{t} parallel with basilar membrane^{b}. The periosteum on upper surface of osseous spiral lamina forms an elevation, the

Limbus laminæ spiralis^{lls}, which presents an upper and lower margin called respectively the *labium vestibulare* and *labium tympanicum*, the groove formed between these being termed the *sulcus spiralis*^{s}.

What is the membrane of Reissner^{r}?

A delicate membrane stretched between outer wall of cochlea and periosteum of vestibular surface of osseous spiral lamina near commencement of limbus laminæ spiralis^{lls}, forming the upper wall of membranous cochlea.

What is the membrana basilaris^{b}?

A thin membrane passing from labium tympanicum to bony wall of cochlea, where it expands into the triangular *ligamentum spirale*^{lsp}. It forms the lower wall of membranous cochlea, supporting on its upper surface the organ of Corti^{co}.

Describe the membrana tectoria or membrane of Corti^{t}.

A delicate membrane passing from between labium vestibulare and attachment of the membrane of Reissner^{r} to outer wall of cochlea, running nearly parallel with basilar membrane^{b}; some contend that it is free, resting upon the organ of Corti^{co}, and in its fresh pulpy condition serves as a damper of vibrations.

Describe the organ of Corti^{co}.

It consists of a complex body formed by a series of some three thousand arches roofing over a space called the *zona arcuata*, formed of rods and epithelial hair cells, lying upon the membrana basilaris^{b} and covered by the membrana tectoria^{t}; the

Rods of Corti are disposed in two rows, the *inner*^{ir} and *outer*^{or} *rods*, each rod with its swollen base resting upon the basilar membrane^{bb}, and its expanded upper end inclined toward and in contact with the opposite one, forming an arched canal filled with endolymph; according to Waldeyer there are about six thousand rods in the inner row, and four thousand five hundred in the outer; a nucleated mass of protoplasm occupies the angle between the base of each rod and the basilar membrane; the

FIG. 114.



Hair cells^{a} (Fig. 115) are epithelial cells with stiff hair-like ciliaβ disposed in three or four rows upon the superior surfaces of the outer rods^{m} to the number of about eighteen thousand, and in a single row upon inner rods^{cd} numbering about thirty-three hundred; the

Lamina reticularis (Fig. 115) is formed of several rows of small fiddle-shaped cuticular structures called *phalanges*, connected together and to heads of outer rods, forming rings^{mno}, through which project the hairs of the outer hair cells.

FIG. 115.



Where does the auditory nerve arise and to what parts of the ear is it distributed?

Its *superficial origin* is from groove between olivary and restiform bodies at lower border of pons; its *deep origin* is threefold (1) from superior vermiformis process of cerebellum, and from (2) inner and (3) outer auditory nuclei formed chiefly by gray substance of posterior pyramid and restiform body, from which latter it receives fibres as it winds around it. Entering with facial nerve the internal auditory canal, at its bottom it divides into the *cochlear* and *vestibular branches*, both of which contain numerous ganglion cells; the

Vestibular division divides into three branches, the superior splitting into many filaments which pass through upper back part of bottom of meatus, enter the vestibule through macula cribrosa to be distributed to utricle and ampulla of external and superior semicircular canals; the middle branch passes through the openings of the fovea hemispherica to supply the saccule; the inferior passes to ampulla of posterior semicircular canal; the

Cochlear nerve divides into numerous filaments at base of modiolus which enter its canals, pass between the two plates forming the lamina spiralis, forming a plexus which contains ganglion cells, and sends branches, it is thought, to communicate with the hair cells; the

Intumescentia ganglioniformis Scarpæ is the gangliform swelling on the vestibular branch in the internal auditory canal; the

Ganglion spirale is the gangliform enlargement on the cochlear nerve as it occupies the *canalis spiralis modioli.*

Give the arterial supply of the labyrinth?

The *internal auditory*, a branch of basilar artery, enters with nerve the internal auditory canal, there to divide into a *vestibular* and *cochlear branch*; the

Stylo-mastoid, from posterior auricular; occasionally

Branches from occipital.

What is the internal auditory canal?

A canal about one-third of an inch long, passing outward from posterior surface of petrous portion of temporal bone to end in a cribriform partition perforated with numerous foramina, some of them arranged in a spiral-shaped depression, the

Tractus spiralis foraminulentus, others in three groups in as many depressions; at the upper part is the commencement of the aquæductus Fallopii for the seventh or portio dura nerve.

Anatomy of Inguinal Hernia.

What is the inguinal canal?

The canal passing downward and inward for one and a half inches which lodges the spermatic cord in the male, the round ligament in the female, pursuing a course parallel to Poupart's ligament through or between the abdominal muscles, commencing at internal abdominal ring and terminating at the external abdominal ring.[8]

Describe the internal abdominal ring.

It is an ovoidal opening in the transversalis fascia[13] ~~midway~~ between anterior superior iliac ~~spine a~~nd spine of pubes, and about half an inch above Poupart's ligament[6]. It is bounded above and externally by arched fibres of transversalis muscle[10], below and internally by epigastric vessels[13]; from its circumference passes the *infundibuliform fascia* on to the spermatic cord or round ligament.

What is the external abdominal ring[8]?

An obliquely triangular opening in the aponeurosis[2] of external oblique muscle[1], just above and to outer side of crest of pubes; from base to apex it averages one inch by half an inch transversely. It is bounded below by the crest of pubes, above by the curved aponeurotic *intercolumnar fibres*[7], and on each side by the free borders of the aponeurosis called the *internal*[5] *and external columns*[6] of the ring; from the margins of the ring passes on to the cord or round ligament, the *intercolumnar fascia*.

Give the boundaries of the inguinal canal.

Anteriorly, the skin, superficial fascia, aponeurosis of external oblique, and outer third of internal oblique muscles.(upper pt.

Posteriorly, the triangular ligament, conjoined tendon[11] of internal oblique and transversalis muscles, transversalis fascia[13], subperitoneal fat, and peritoneum.

Superiorly, the arched fibres of internal oblique[10] and transversalis muscles.

Inferiorly, the union of the transversalis fascia[13] and Poupart's ligament[6].

What is Poupart's ligament[6]?

The infolded, thickened margin of aponeurosis of ~~external oblique~~ muscle, extending from anterior iliac space[9] to spine of pubes, from which it is reflected a short distance along the pectineal line, forming *Gimbernat's ligament*, while a triangular band of tendinous fibres attached by its ~~apex~~ to the reflected portion of Poupart's ligament along the pectineal line, passing inward beneath spermatic cord behind the inner pillar in front of the conjoined tendon, and interlacing with its fellow at the linea alba, is called the *triangular ligament*: the lower part of Poupart's ligament forms the external pillar of the external abdominal ring[6].

FIG. 116.



Describe the relations of the epigastric artery and the internal abdominal ring.

It passes between the transversalis fascia and the peritoneum along the lower inner margin of internal ring beneath the spermatic cord.

What is the cremasteric fascia?

A series of muscular loops connected by areolar tissue investing the spermatic cord, supposed to have been derived from the internal oblique muscle during the descent of the testicle, and, of course, absent in the female.

What is Hesselbach's triangle?

A triangular space at lower part of inner surface of abdominal wall, bounded externally by the epigastric artery, internally by margin of rectus muscle, and below by Poupart's ligament: the conjoined tendon stretches across the inner two-thirds of this space, the remainder being filled in by the transversalis fascia.

What is an oblique inguinal hernia?

A protrusion of bowel, omentum, or both, following the course of the spermatic cord in the inguinal canal through both internal and external rings (*when complete*), the neck of the sac being to *outer* side of epigastric artery.

What are the coverings of an oblique inguinal hernia?

Skin; *superficial fascia*; *intercolumnar fascia*; *cremaster muscle*[12]; *infundibuliform fascia*; *subserous cellular tissue*; *and peritoneum* forming *sac*.

Where is the seat of stricture most apt to be?

(1) At internal ring, (2) inguinal canal by fibres of internal oblique or transversalis muscles, (3) at external ring, provided it is not at thickened neck of sac, the most usual site in *old herniæ*.

What is a direct inguinal hernia?

One where the protrusion passes through some part of Hesselbach's triangle, passing directly through the external abdominal ring, the neck of the sac being *internal* to the epigastric vessels.

What coverings has a direct hernia? = same w. dif.

The same as an oblique, except that the *transversalis fascia* take the place of the *infundibuliform fascia*, and the *conjoined tendon* is substituted for the cremasteric fascia, but this latter covering is probably a *theoretical*, not an actual one.

Femoral Hernia.

Describe the femoral or crural canal.

It lies beneath Poupart's ligament[6], to inner side of femoral vein[19], extending from *femoral ring* above to *saphenous opening*[17] below: it is about half an inch long, closed above by the *septum crurale*, formed of condensed areolar tissue, and below by the *cribriform fascia*, a portion of the deep layer of the superficial fascia covering the saphenous opening: the

Femoral ~~or crural~~ ring is an oval space between femoral vein and Gimbernat's ligament, half an inch in diameter, larger in female than in male, situated below and external to internal abdominal ring: it is closed by the septum crurale and a lymphatic gland; the *Saphenous opening*[17] is an ovoidal opening one and a half inches long by half an inch wide, below inner portion of Poupart's ligament[6], formed by the pubic portion[17] of fascia lata passing behind the saphenous vein leaving a margin concave outward, while continuous with it is a strong f~~alciform process~~, the ili~~ac portion~~ of the fascia lata, the *falciform process of Burns*[16], passing in *~~front~~* of the vein, its narrow pubic portion blending with the attachment of Poupart's and Gimbernat's ligaments; this is called *Hey's* ligament[16]: the opening is covered in by the *cribriform fascia.*

Bound the femoral canal.

Anteriorly, lie the transversalis fascia[13], Poupart's ligament[6], and falciform process of fascia lata[16]; Pectineal

Posteriorly, iliac fascia, covering pubic portion of fascia lata;

Externally, the fibrous septum separating it from femoral vein;

Internally, the junction of the processes of the transversalis and iliac fasciæ forming femoral sheath lying against outer edge of Gimbernat's ligament.

Describe the position of parts around the ring.

The *Spermatic cord* in male, the *round ligament* in female, lie just above anterior margin; the

Femoral vein[19] lies upon its outer side; the

Epigastric artery crosses upper outer angle of ring; the

Obturator artery once in three and a half cases arises in common with epigastric, when it closely skirts around the upper and inner margins of the ring.

Describe the septum crurale.

It is a layer of condensed cellular tissue upon which lies a small lymphatic gland, closing the femoral ring. = Sub peritoneal

What is the crural sheath?

It is a continuation downward of the ~~fascia transversalis~~ in front, and of the iliac fascia behind the femoral vessels forming their sheath, divided by septa into separate compartments for femoral artery externally, femoral vein next, and leaving a third internally, the femoral canal.

Describe the deep crural arch.

When present, for it is not infrequently absent, it is a band of fibres derived from transversalis fascia, crossing the front of, and adherent to the crural sheath, passing from the centre of Poupart's ligament to the pectineal line behind the conjoined tendon.

Where is the seat of stricture most apt to be?

At (1) junction of Hey's and Gimbernat's ligaments, (2) margin of saphenous opening, (3) neck of hernial sac.

What are the coverings of a femoral hernia?

They are *Skin, Superficial fascia, Cribriform fascia, Crural sheath, Septum crurale, Subserous areolar tissue, Peritoneum* forming *sac.*

Perineum and Ischio-rectal Regions.

What is the ischio-rectal region?

That portion of the outlet of the pelvis immediately behind the perineum containing the rectum, upon either side of which is a deep fossa filled with fat, the

Ischio-rectal fossa[5], of a triangular shape, between the end of rectum[x] and tuberosity of the ischium[c] on each side; its base corresponds to the skin, and apex to point of division of obturator fascia and origin of anal fascia; its base measures about one inch, its depth two inches, being deepest behind, and is bounded *internally*, by sphincter ani[x], levator ani[5], and coccygeus[6] muscles; *externally*, by tuberosity of ischium[c] and obturator fascia; in *front*, by line of junction of superficial and deep perineal fascia; and *behind*, by margin of gluteus maximus muscle[11] and great sacro-sciatic ligament; the

Internal pudic artery and nerve run about one and a half inches above margin of ischiatic tuberosity, becoming more superficial as they pass forward along inner margin of pubic arch; the

Inferior hemorrhoidal vessels, occasionally large enough to give trouble after lithotomy, traverse the centre of each fossa.

What is the perineum?

It is the anterior portion of the pelvic outlet, in front of ischio-rectal fossæ, is of a triangular form, bounded, deeply, by rami of pubes and ischia, forming, superiorly, the pubic arch, and behind, by an imaginary line drawn between ischiatic tuberosities; the lateral boundaries, in adult, vary from three to three and one-half inches.

Describe the muscles of the perineum.

External sphincter ani[x]: *origin*, tip of coccyx[b] and superficial fascia in front of bone; *insertion*, tendinous centre of perineum[2]: *action*, closes anus; *nerve*, fourth sacral.

Internal sphincter ani: consists of an aggregation of the involuntary circular fibres of intestine, forming a muscular ring two lines thick and one inch broad.

FIG. 117.



add. longus.
gracilis
add. magnus.
Biceps a Semiten

Sphincter tertius (Hyrtl): *origin*, sacrum, encircling rectum four inches above anus (not described by many anatomists).

Bulbo-Cav.

Accelerator urinæ[1]: consists of two symmetrical halves with median tendinous raphé; *origin*, central tendon of perineum[2] and median raphé in front; *insertion,* fibres spread over front of triangular ligament, encircle the bulb and corpus spongiosum, spread over sides of corpora cavernosa, to which they are partly attached and partly terminate in a tendinous expansion covering dorsal vessels of penis; *action,* empties urethra after bladder ceases to contract, perhaps aids in erection of penis; *nerve,* perineal branch of pudic. (Ischio-cav.)

Erector penis[3]: *origin,* inner surface of each tuberosity of ischium, surface of crus penis and adjacent portion of pubic ramus (*Erector clitoridis* in female *arises* from ischial tuberosity; *inserted* on side of crus clitoridis); *insertion,* by an aponeurosis into sides and under surface of each crus penis; *action,* maintains erection; *nerve,* perineal.

Transversus perinei[4]: *origin,* inner front side of tuberosity of ischium; *insertion,* tendinous centre of perineum (in female, side of sphincter vaginæ); *action,* steadies perineal centre; *nerve,* perineal.

Levator ani[5]: *origin,* body and ramus of pubes posteriorly, inner surface of spine of ischium, from angle of division between obturator and recto-vesical layers of pelvic fascia; *insertion,* sides of apex of coccyx[b], opposite muscle by median fibrous raphé, extending from coccyx to anal margin, side of rectum (and vagina in female) blending with sphincter muscles, and side of prostate gland, uniting beneath it with opposite muscle mingling with external sphincter and transversus peronei muscles at perineal centre[2]; *action,* supports rectum, vagina, and pelvic viscera, with its fellow helps form the floor of the pelvis; *nerves,* inferior hemorrhoidal and fourth sacral.

Compressor urethræ: *origin,* aponeurotic, from three-fourths of an inch of upper part of pubic ramus on each side; *insertion,* each muscle divides surrounding urethra from prostate gland to bulb of urethra, uniting with the opposite muscle above and below this tube by a tendinous raphé; *action,* shut-off muscle; *nerves,* perineal.

Coccygeus[6]: *origin,* by apex from spine of ischium and lesser sacro-sciatic ligament; *insertion,* by base into margin of coccyx and side of lowest segment of sacrum; *action,* raises coccyx; completes pelvic floor behind; *nerve,* anterior division fifth sacral.

Sphincter vaginæ: (in female) surrounds vaginal orifice, is homologous with accelerator urinæ in male—*origin*, central tendon of perineum; *insertion*, corpora cavernosa of clitoris.

Erector clitoridis, in female replaces the erector penis muscle.

Describe the perineal fasciæ.

The *superficial layer of the superficial fascia*, is thick, loose, loaded with fat and continuous with the fasciæ of ischio-rectal and crural regions; the

Deep layer of the superficial fascia is a thin, moderately strong, aponeurotic layer, continuous in front with dartos layer of scrotum, is firmly attached to the margins of the rami of ischium and pubes external to crura of penis as far back as ischial tuberosity, and passes down behind the transversus perinei muscles to blend with lower margin of the deep perineal fascia; the

Anterior layer of the deep perineal fascia is attached above and laterally to pubic arch, subpubic ligament and rami of ischium and pubes beneath crura of penis, forming a dense membranous lamina, extending and attached to tendinous perineal centre, becoming continuous with deep layer of the superficial fascia behind transverse perineal muscles: it embraces and is continued forward on anterior extremity of membranous urethra, which passes through it one inch below symphysis pubis; the

Posterior layer of the deep perineal fascia is attached like preceding, but more posteriorly, embraces back portion of membranous urethra and is continued backward over outer surface of prostate gland, the two layers forming the *triangular ligament*: the portion of urethra embraced between these layers is the *membranous urethra*; the

Obturator fascia is a continuation of transversalis and iliac fasciæ below a white band at level of a line passing from lower part of symphysis pubis to spine of ischium: it covers the internal obturator muscle and is continuous with posterior layer of triangular ligament; the

Recto-vesical fascia, or *visceral layer of pelvic fascia*, descends into pelvis upon upper surface of levator ani muscle, investing prostate, bladder, and rectum.

Describe Buck's fascia.

It is a moderately dense fascia investing the penis as far as glans, being a prolongation forward of deep layer of superficial fascia, blending with dartos layer of scrotum: it directs the urine forward into scrotum, penis, and upon abdomen when the urethra is ruptured.

What layers of fascia form the triangular ligament?

The superficial and deep layers of the *deep perineal fascia* according to the description of most authorities.

What structures lie between the layers of the deep perineal fascia?

Membranous urethra,
Compressor urethræ muscle,
Subpubic ligament,
Dorsal vein of penis,
Cowper's glands and ducts.
Pudic vessels,
Pudic nerves,
Arteries and nerves of bulb,
Venous plexus.

What structures are exposed by the removal of the deep layer of the superficial fascia?

The erector penis muscles covering the crura of penis.
The accelerator urinæ muscle covering corpus spongiosum and bulb.
Transversus perinei muscles.
Transversus perinei arteries.
Superficial perineal vessels.

What parts are divided in lateral lithotomy?

Skin.
Superficial fascia.
Inferior hemorrhoidal vessels and nerves.
Posterior fibres of accelerator urinæ muscle.
Transversus perinei muscle and vessels.
Superficial perineal vessels and nerves, (probably) *deep perineal fascia.*
Anterior fibres of levator ani.
Part of compressor urethræ.
Membranous and prostatic portions of urethra.
Part of prostate gland (neck of bladder, according to Gross).

What structures must be avoided in this operation?

In front, the bulb and its artery.

Externally, pudic artery.

Toward median line and posteriorly, the rectum.

At neck of bladder, the entire division of lateral lobe and, therefore, the *capsule* of prostate gland and venous plexus.

Briefly describe the special points of interest connected with the female perineum.

The *perineal body* is the point of junction of all the muscular and aponeurotic structures of perineum, forming a pyramidal mass of tissue extending for some distance up between rectum and vagina, upon the integrity of which depends the proper support afforded to the pelvic viscera by the *floor of* the *pelvis*

The length of the female perineum is only about one inch, extending from posterior commissure to verge of anus: the *accelerator urinæ muscle* is replaced by the *sphincter vaginæ* encircling the vaginal outlet: the triangular ligament, formed essentially as in male, and perforated by the urethra, is much smaller.

INTRODUCTION

TO

PLATES ON OSTEOLOGY

The following illustrations, taken from GRAY, are added with the idea of making this work a complete guide to the Student in his practical work. In the majority of our schools a careful study of osteology precedes dissection, the Student being given each bone for examination, and being required to master its anatomical details before receiving another one. The assistance offered by a complete set of plates needs no comment. Supplemented, as these illustrations are, by an admirably clear and condensed text, the Student has, in this work, a most efficient aid in that portion of his anatomical studies which is usually considered most dull and uninteresting.

THE PUBLISHER.

INDEX OF OSTEOLOGICAL PLATES.

Fig. 118 Cervical Vertebra....... 338
Fig. 119 First Cervical Vetetra or Atlas............ 338
Fig. 120 Second Cervical Vertebra or Axis.............. 338
Fig. 121 Seventh Cervical Vertebra......................... 339
Fig. 122 A Dorsal Vertebra....... 339
Fig. 123 Lumbar Vertebra....... 339
Fig. 124 Peculiar Dorsal Vertebra........................... 340
Fig. 125 Sacrum, anterior surface 341
Fig. 126 Sacrum, posterior surface.......................... 341
Fig. 127 Occipital Bone, outer surface..................... 342
Fig. 128 Occipital Bone, inner surface..................... 342
Fig. 129 Left Parietal Bone, external surface......... 343
Fig. 130 Left Parietal Bone, internal surface......... 343
Fig. 131 Frontal Bone, outer surface..................... 344
Fig. 132 Frontal Bone, inner surface..................... 344
Fig. 133 Left Temporal Bone, outer surface.......... 345
Fig. 134 Left Temporal Bone, inner surface.......... 345
Fig. 135 Petrous Portion......... 346
Fig. 136 Sphenoid Bones, superior surface.............. 347
Fig. 137 Sphenoid Bones, anterior surface......... 347
Fig. 138 Ethmoid Bone............ 348
Fig. 139 Perpendicular Plate of Ethmoid (enlarged) 348
Fig. 140 Right Nasal Bone, outer surface.......... 349
Fig. 141 Left Nasal Bone, inner surface..................... 349
Fig. 142 Left Superior Maxillary Bone, outer surface..................... 349
Fig. 143 Left Superior Maxillary Bone, inner surface 350
Fig. 144 Left Malar Bone, outer surface..................... 350
Fig. 145 Left Malar Bone, inner surface 350
Fig. 146 Palate and Alveolar Arch 351
Fig. 147 Left Palate Bone, internal view 351
Fig. 148 Left Palate Bone, posterior view 352
Fig. 149 Left Lachrymal Bone, external surface... 352
Fig. 150 Right Inferior Turburated Bone, internal surface.............. 352
Fig. 151 Right Inferior Turburated Bone, outer surface..................... 352
Fig. 152 Vomer.......................... 352
Fig. 153 Inferior Maxillary Bone, outer surface, side view....... 353
Fig. 154 Inferior Maxillary Bone, inner surface, side view....... 353
Fig. 155 Base of the Skull, inner surface............. 354
Fig. 156 Base of Skull, external surface..................... 355
Fig. 157 Anterior Region of the Skull................. 356
Fig. 158 Hyoid Bone, anterior surface..................... 356
Fig. 159 Roof, Floor and Outer Wall of Nasal Fossa 357
Fig. 160 Left Clavicle, anterior surface 357
Fig. 161 Left Clavicle, inferior surface.... 357
Fig. 162 Sternum and Costal Cartilages 358
Fig. 162* Posterior Surface of Sternum.................. 358
Fig. 163 Peculiar Ribs.............. 359
Fig. 164 Left Scapula, anterior surface or venter... 360
Fig. 165 Left Scapula, posterior surface or dorsum 361
Fig. 166 Left Humerus, anterior view................. 362
Fig. 167 Left Humerus, posterior view.................. 362
Fig. 168 Bones of the Forearm, anterior surface...... 363
Fig. 169 Bones of the Forearm, posterior surface... 364
Fig. 170 Bones of the Hand, dorsal surface......... 365
Fig. 171 Bones of the Hand, palmer surface....... 366
Fig. 172 Right Os Innominatum, external surface.......................... 367
Fig. 173 Right Os Innominatum, internal surface 368
Fig. 174 Right Patella, anterior and posterior surface 368
Fig. 175 Right Femur, anterior surface.............. 369
Fig. 176 Right Femur, posterior surface 369
Eig. 177 Bones of the Leg, anterior surface..... 370
Fig. 178 Bones of the Leg, posterior surface..... 370
Fig. 179 Bones of the Foot, dorsal surface... 371
Fig. 180 Bones of the Foot, plantar surface....... 372

FIG. 124.



PECULIAR DORSAL VERTEBRÆ.

FIG. 121.



SEVENTH CERVICAL VERTEBRA OR VERTEBRA PROMINENS.

FIG. 122.


Superior Artic. Process
Demi facet for head of Rib
Facet for Tubercle of Rib
Trans. Proc.
Pedicle
Body
Spin. Proc.
Infer. Artic. Proc.
Demi facet for head of Rib

A DORSAL VERTEBRA.

FIG. 123.



LUMBAR VERTEBRA.

FIG. 124.



PECULIAR DORSAL VERTEBRÆ.

FIG. 125.



SACRUM, ANTERIOR SURFACE.

FIG. 126.



SACRUM, POSTERIOR SURFACE.

Fig. 127.



Occipital Bone, outer surface.

Fig. 128.



Occipital Bone, inner surface.

FIG. 129.


Articulates with opposite Parietal B.
Parietal Foramen
Parietal eminence
Articulates with Frontal B.
Temporal Ridge
TEMPORAL MUSCLE
Articulates with Occipital B.
Sphenoid
Squamous Portn of Temp. B.
Mastoid Portn

LEFT PARIETAL BONE, EXTERNAL SURFACE.

FIG. 130.



LEFT PARIETAL BONE, INTERNAL SURFACE.

FIG. 131.



FRONTAL BONE, OUTER SURFACE.

FIG. 132.


Articulates with Parietal Bone
Pacchionian Depressions
for Superior Longitudinal Sinus
Meningeal Grooves
for Falx Cerebri
with Lesser Wing
with Greater Wing of Sphenoid
with Malar
Posterior and Anterior Ethmoidal for.
with Ethmoid
Orbital Plate
Lachrymal Fossa
with Lachrymal
with Sup. Maxill.
with Nasal
with Perpendicular plate of Ethmoid
Frontal Sinus
Expanded base of Nasal Spine, forming part of Roof of Nose

FRONTAL BONE, INNER SURFACE.

FIG. 133.



LEFT TEMPORAL BONE, OUTER SURFACE.

FIG. 134.


Articulates with Parietal Bone
Squamous Portion
Inner Surface
Petrous Portion
with Occipital Bone
Posterior Surface
Sphenoid
and Great Wing
Sup. Petrosal
Aqueductus Vestibuli
Depression for Dura-mater
Meatus Auditorius internus
Eminence for Superior Semicircular Canal.
Hiatus Fallopii
Opening for Smaller Petrosal Nerve
Depression for Casserian ganglion
Bristle passed through Carotid Canal

LEFT TEMPORAL BONE, INNER SURFACE.

FIG. 135.



PETROUS PORTION, INFERIOR SURFACE.

FIG. 136.



SPHENOID BONE, SUPERIOR SURFACE.

FIG. 137.



SPHENOID BONE, ANTERIOR SURFACE.

FIG. 138.



ETHMOID BONE, OUTER SURFACE OF RIGHT LATERAL MASS, (ENLARGED).

FIG. 139.


with Ethmoidal Spine, and Rostrum of Sphenoid
Articulates with Frontal with Nasal
Perpendicular Plate
Artic. with Vomer
with Triangular Cartilage of Septum

PERPENDICULAR PLATE OF ETHMOID (ENLARGED), SHOWN BY REMOVING THE RIGHT LATERAL MASS.

FIG. 140.



RIGHT NASAL BONE,
OUTER SURFACE.

FIG. 141.



LEFT NASAL BONE,
INNER SURFACE.

FIG. 142.



LEFT SUPERIOR MAXILLARY BONE, OUTER SURFACE.

FIG. 143.



LEFT SUPERIOR MAXILLARY BONE, INNER SURFACE.

FIG. 144.



LEFT MALAR BONE, OUTER SURFACE.

FIG. 145.



LEFT MALAR BONE, INNER SURFACE.

FIG. 146.



THE PALATE AND ADVEOLAR ARCH.

FIG. 147.



LEFT PALATE BONE, INTERNAL VIEW.

FIG. 148.



LEFT PALATE BONE, POSTERIOR VIEW.

FIG. 149.



LEFT LACHRYMAL BONE, EXTERNAL SURFACE (ENLARGED).

FIG. 150.



RIGHT INFERIOR TURBINATED BONE, INTERNAL SURFACE.

FIG. 151.



RIGHT INFERIOR TURBINATED BONE, EXTERNAL SURFACE.

FIG. 152.


Artic. with Sphenoid.
with Ethmoid.
with Triangular Cartilage.
Naso-palatine groove.
Alæ.
with Sup.-Maxill. bones and Palate.

VOMER.

FIG. 153.



INFERIOR MAXILLARY BONE, OUTER SURFACE, SIDE VIEW.

FIG. 154.



INFERIOR MAXILLARY BONE, INNER SURFACE, SIDE VIEW.

Fig. 155.



BASE OF THE SKULL, INNER OR CEREBRAL SURFACE.

FIG. 156.



BASE OF THE SKULL, EXTERNAL SURFACE.

Fig. 157.



Anterior Region of the Skull.

Fig. 158.



Hyoid Bone, anterior surface.

FIG. 159.



ROOF FLOOR, AND OUTER WALL OF LEFT NASAL FOSSA.

FIG. 160.



LEFT CLAVICLE, ANTERIOR SURFACE.

FIG. 161.



LEFT CLAVICLE, INFERIOR SURFACE.

FIG. 157.



ANTERIOR REGION OF THE SKULL.

FIG. 158.



HYOID BONE, ANTERIOR SURFACE.

Fig. 163.


Fig. 194.
Fig. 195.
1st
Shortest
has a single artic. facet
Surfaces horizontal
Tubercle
Scalenus anticus
Groove for Subclavian Vein
Scalenus medius
Groove for Subclavian Artery
no Angle
Serratus magnus
double origin of 1st Serratus
2nd
Angle
Slightly marked & close to tuberosity
Rough muscular eminence



Peculiar Ribs.

Fig. 164.



Left Scapula, anterior surface or venter.

FIG. 165.



LEFT SCAPULA, POSTERIOR SURFACE OR DORSUM.

Fig. 166.



Left Humerus, anterior view.

Fig. 167.



Left Humerus, posterior surface.

Fig. 168.



Bones of the Left Forearm, anterior surface.

FIG. 166.



LEFT HUMERUS, ANTERIOR VIEW.

FIG. 167.



LEFT HUMERUS,
POSTERIOR SURFACE.

FIG. 168.



BONES OF THE LEFT FOREARM, ANTERIOR SURFACE.

FIG. 169.



BONES OF THE LEFT FOREARM, POSTERIOR SURFACE.

FIG. 170.



BONES OF THE LEFT HAND, DORSAL SURFACE.

Fig. 171.


Semi lunar
Scaphoid
Carpus
Cuneiform
Pisiform
Unciform
Os magnum
Trapezoid
Trapezium
FLEXOR CARPI ULNARIS
FLEXOR BREVIS MINIMI DIGITI
FLEXOR OSSIS METACARPI MINIMI DIGITI
Groove for tendon of FLEXOR CARPI RADIALIS
FLEXOR OSSIS METACARPI POLLICIS
EXTENSOR OSSIS METACARPI POLLICIS
FLEX. OSS. MET. POLLICIS
Metacarpus
3rd PALMAR INTEROSSEOUS
2nd PALMAR INTEROSSEOUS
1st PALMAR INTEROSSEOUS
ADDUCTOR POLLICIS
Sesamoid bones
ABDUCTOR POLLICIS
FLEXOR BREVIS POLLICIS
FLEX. BREVIS & ABDUCTOR MINIMI DIGITI
Phalanges
FLEXOR LONGUS POLLICIS
FLEXOR DIGITORUM SUBLIMIS
FLEXOR DIGITORUM PROFUNDUS
FLEXOR DIGITORUM SUBLIMIS
FLEXOR DIGITORUM PROFUNDUS

BONES OF THE LEFT HAND, PALMER SURFACE.

FIG. 172.



RIGHT OS INNOMINATUM, EXTERNAL SURFACE.

FIG. 173.



RIGHT OS INNOMINATUM, INTERNAL SURFACE.

FIG. 174.


EXTENSOR CRURIS
Subcutaneous & covered by a Bursa.
Outer Facet for Artic. with External Condyle of Femur.

RIGHT PATELLA,
ANTERIOR SURFACE. POSTERIOR SURFACE.

FIG. 175.



RIGHT FEMUR,
ANTERIOR SURFACE.

FIG. 176.



RIGHT FEMUR,
POSTERIOR SURFACE.

FIG. 177.



BONES OF THE RIGHT LEG,
ANTERIOR SURFACE.

FIG. 178.



BONES OF THE RIGHT LEG,
POSTERIOR SURFACE.

FIG. 179.



BONES OF THE RIGHT FOOT, DORSAL SURFACE.

FIG. 180.



BONES OF THE RIGHT FOOT, PLANTAR SURFACE.

INDEX.

ABDOMEN, 243
  muscles of, 129
Abdominal cavity, 243
  regions, 245
Absorbents, 184
Acervulus cerebri, 196
Acetabulum, 68
Alveoli of lower jaw, 50
  of upper jaw, 43
  of the lungs, 258
Amphiarthrosis, 92
Anastomoses, 157, 175
Anatomy, general, 17
  descriptive, 17
Angle of the jaw, 51
  of subscapular fossa, 76
Annulus ovalis, 154
Antihelix, 310
Antitragus, 310
Antrum of Highmore, 40
Anus, 232
Aorta, 158, 167, 168
Appendices epiploicæ, 232
Appendix auriculæ, 154, 155
  ensiform, 65
  vermiformis, 231
Aponeuroses, 115
  pharyngeal, 225
Apophysis, 21
Aqueduct of Sylvius (iter e tertio, etc.), 194, 195
Aquæductus cochleæ, 36
  Fallopii, 36, 203
  vestibuli, 319
Arachnoid membrane of brain, 186
  of spinal cord, 199
Arbor vitæ, 198, 282
Arch, crural, deep, 330
  palmar, 166
  plantar, 175
Areola of mammæ, 286
Arm, 78
Arnold's ganglion, 202
Arteriæ propriæ renales, 263
Arterial anastomoses, 175
Arteries, helicine, 274
  hepatic, 168
  lingual, 161
  pharyngeal, 161
  sublingual, 161
Arteries of the—
  ankle-joint, 112
  auditory canal, 311
  auricle, 311
  bone, 19
  choroid, 298
  elbow-joint, 104
  Eustachian tube, 317
  eye, 304
  eye-lids, 304
  hip-joint, 108
  iris, 296
  kidney, 263, 264
  knee-joint, 110
  labyrinth, 325
  larynx, 254
  lung, 259
  mammæ, 287
  membrana tympani, 312
  nasal fossæ, 291
  nose, 291
  œsophagus, 226
  ovaries, 286
  pancreas, 239
  penis, 274
  pharynx, 226
  retina, 299
  shoulder-joint, 103
  spleen, 240
  stomach, 000
  supra-renal capsules, 243
  testicle, 277
  thymus gland, 242
  thyroid gland, 241
  tympanum, 316
  uterus, 284
  wrist-joint, 105
Arteriolæ rectæ, 264
Artery—
  anastomotica magna, 173

Artery—
  aorta, 158, 167, 168
  arteria receptaculi, 162
  auricular, 161
  axillary, 165
  basilar, 164
  brachial, 165
  carotid (external), 161
    (internal), 162
  cerebral, anterior, 163
    middle, 163
    posterior, 163
  choroid, anterior, 166
  communicating, anterior, 163
    posterior, 163
  coronary, 159
  digital, 167, 175
  dorsalis hallucis, 174
    pedis, 174
  epigastric, deep, 171
    superficial, 171
  facial, 161
  femoral, 171
  gastric, 168
  gluteal, 170
  hemorrhoidal, 170
  hepatic, 168
  iliac, circumflex, 171
    external, 171
    internal, 169
  ilio-lumbar, 170
  infra-orbital, 162
  innominate, 159
  intercostal, 167
  lingual, 161
  lumbar, 169
  mammary, internal, 164
  maxillary, internal, 161
  meningeal, anterior, 162
  mesenteric, superior, 168
    inferior, 168
  obturator, 170
  occipital, 161
  ophthalmic, 162
  palmar arches, 166
  peroneal, 174
  pharyngeal, ascending, 161
  phrenic, 168
  plantar, external, 174
    internal, 174
  popliteal, 173
  profunda femoris, 173
    inferior, 165
    superior, 165

Artery—
  pudic, deep external, 173
    superficial external, 173
      internal, 170
  pulmonary, 175
  radial, 166
  renal, 169
  sacral, lateral, 170
    middle, 169
  sciatic, 170
  spermatic, 169
  splenic, 168
  subclavian, 163
  subscapular, 165
  suprarenal, 169
  suprascapular, 164
  temporal, 161, 162
  thoracic, acromial, 165
    superior, 165
  thoracica alaris, 165
    longa, 165
  thyroid, superior, 161
    inferior, 164
  thyroidea ima, 159
  tibial, anterior, 173
    posterior, 174
  transversalis colli, 164
  tympanic, 162
  ulnar, 166
  uterine, 170
  vaginal, 170
  vertebral, 164
  vesical, 169, 170
Arthrodial joints, 93
Articulations, 92
  atlo-axoidean, 95
  carpal, 105
  carpo-metacarpal, 106
  costo-sternal, 98
  costo-transverse, 98
  costo-vertebral, 97
  metacarpo-metacarpal, 106
  metatarso-metatarsal, 113
  occipito-atloid, 94
  occipito-axoidean, 96
  phalangeal, of foot, 114
    of hand, 107
  radio-ulnar, 104
  sacro-coccygeal, 100
  sacro-iliac, 99
  sacro-ischiatic, 100
  sacro-vertebral, 99
  scapulo-clavicular, 101
  scapulo-humeral, 102

- Artery—
  - sterno-clavicular, 101
  - tarsal, 112
  - tarso-metatarsal, 113
  - temporo-maxillary, 96
  - tibio-fibular, 110
  - vertebral, 94
- Arytenoid cartilages, 250
- Atlas, 63
- Auricle, 310
- Axis, 63
  - cœliac, 168
  - optic, 292
  - thyroid, 164
  - visual, 292
- Azygos veins, 180
  - uvulæ, 122

- BARTHOLIN, duct of, 221
  - glands of, 280
- Bauhin, valve of, 231
- Bell, external nerve of, 206
  - internal nerve of, 205
- Bladder, 266
- Bodies, Malpighian, 263
  - Pacchionian, 186
- Body, ciliary, 296
  - olivary, 196
  - perineal, 335
  - pituitary, 191
  - restiform, 197
- Bone, astragalus, 90
  - calcaneum, 90
  - clavicle, 75
  - coccyx, 72
  - cuboid, 90
  - cuneiform, 91
  - ethmoid, 38
  - femur, 85
  - fibula, 89
  - frontal, 23
  - humerus, 78
  - hyoid, 65
  - ilium, 69
  - incus, 314
  - innominate, 68
  - ischium, 70
  - lachrymal, 44
  - malar, 44
  - malleus, 314
  - maxillary, inferior, 49
    - superior, 40
  - nasal, 40
- Bone—
  - occipital, 27
  - os calcis, 90
  - os magnum, 84
  - palate, 45
  - parietal, 25
  - pectineal, 70
  - peroneal, 89
  - phalanges of foot, 92
    - of hand, 85
  - pisiform, 84
  - pubic, 70
  - sacral, 71
  - scaphoid, 83
  - scapula, 76
  - semilunar, 83
  - sphenoid, 30
  - stapes, 315
  - sternum, 65
  - structure, microscopic, 18
  - temporal, 34
  - tibia, 88
  - trapezium, 84
  - trapezoid, 84
  - turbinated, inferior, 47
    - middle, 39
    - superior, 39
  - ulna, 80
  - unciform, 84
  - vomer, 48
- Bones of the body, 20
  - of the carpus, 83
  - of the foot, 90
  - of the hand, 82
  - of the leg, 88
  - of the lower extremity, 85
  - of the metacarpus, 84
  - of the metatarsus, 91
  - of the tarsus, 90
  - of the trunk, 61
  - of the upper extremity, 74
  - Wormian, 20
- Bonnet, capsule of, 293
- Bowman's membrane, 295
- Brain, commissures of, 192
  - divisions of, 187
  - points on base, 190
    - on surface, 187–190
  - ventricles, 192
  - weights, average, 187
- Bronchi, 255
- Brunner's gland, 230
- Bulb of corpus cavernosum, 273
  - of corpus spongiosum, 273

Bulbi vestibuli, 280
Bulbs, olfactory, 191
Burns, ligament of, 329
Bursæ of ankle, 149
    of hip, 108
    of knee, 110
    of shoulder, 102
    of wrist, 139

CÆCUM, 231
Calamus scriptorius, 197
Calices of kidney, 261
Canal, alimentary, 224
    auditory, external, 311
        internal, 36
    carotid, 36
    central, of the cord, 200
    crural, 329
    dental, 41, 50
    femoral, 328
    Hunter's, 144, 171
    hyaloid, 301
    infraorbital, 41
    inguinal, 326
    of the modiolus, 320
    nutrient, 19
    palatine, 46
    sacral, 72
    spiral, of the cochlea, 320
    for tensor tympani, 37
    Vidian, 33
Canal of—
    Huguier, 35
    Nuck, 383
    Petit, 302
    Schlemm, 297
    Stilling, 301
Canaliculi, 308
Canalis reuniens, 322
Canals, semicircular, 319
    membranous, 322
Canthus, 306
Capillaries, 158
Capsule, internal, 194
    of Bonnet, 293
    of Glisson, 237
    of the lens, 301
    of Tenon, 293
Capsules, suprarenal, 242
Caput cæci, 231
    gallinaginis, 269
Carpus, 83
Cartilages of the larynx, 249
Cartilages of Santorini, 250
    of tarsal, 306
    of Wrisberg, 251
Cartilago triticea, 251
Caruncula lachrymalis, 307
    myrtiformes, 280
Cauda equina, 211
Cavity, cotyloid, 68
    glenoid, of scapula, 78
    of the omentum, 246
    orbital, 55
    of reserve, 220
    sigmoid, greater, 80
        lesser, 80
        of radius, 82
Cells, auditory, 324
    ethmoid, 39
    hepatic, 237
    mastoid, 35, 314
    olfactory, of Schultze, 291
Cement, 218
Cerebellum, 197
Cerebro-spinal system, 185
Cerebrum, 187
    convolutions of, 189
    fissures of, 187
    lobes of, 188
Cervix of penis, 273
    of uterus, 282
Chordæ tendineæ, 155
Choroid tunic, 295
    plexus, 193
Cilia, 307
Circle of Willis, 163
Circulus arteriosus iridis major, 296
        iridis minor, 296
    venosus, 287
Clavicle, 75
Clinoid processes, 30, 33
Clitoris, 280
Coccyx, 72
Cochlea, membranous, 322
    osseous, 320
Cœliac axis, 168
Collar-bone, 75
Colon, 231
    flexures of, 231
Columnæ carneæ, 155, 157
    papillares, 155
Columns of Bertin, 261
    of spinal cord, 199
Commissure, optic, 200
Concha, 310

Condyles, of femur, 86
Condyles of humerus, 79
Coni-vasculosi, 276
Conjunctiva, 307
Conus arteriosus, 154
Convolutions of cerebrum, 189
Coracoid process, 77
Corium, 288
Cornea, 294
Cornicula laryngis, 250
Cornua of thyroid cartilage, 250
    of the ventricles, 192
Corona glandis, 273
    radiata, 195
Coronoid process, 49, 81
Corpora albicantia, 191
    cavernosa, 273
    quadrigemina, 192, 195
    striata, 192
Corpus Arantii, 155
    callosum, 193
    dentatum of cerebellum, 198
        of olivary body, 196
    fimbriatum, 193
    highmorianum, 275
    luteum, 285
    spongiosum, 273
Corpuscles, Malpighian, 240
    tactile, 288
Corti, organ of, 323
Cortical substance of kidney, 261
Cotyloid cavity, 68
Coverings of the testicle, 275
    of hernia, 328, 330
Cowper's glands, 272
Crest of the ilium, 69
    lachrymal, 44
    nasal, 43, 47
    of the pubes, 71
    of the tibia, 88
    turbinated, 42, 46
Cricoid cartilage, 250
Crista galli, 38
Crura cerebri, 191
    of clitoris, 280
    of penis, 273
Crusta petrosa, 218
Crypts of Lieberkühn, 230
Cuneiform cartilages, 251
Cuneus, 190
Cupola of cochlea, 320
Cuticle, 287, 288

DARTOS, 276
    Dentine, 218
Derma, 288
Descemet, membrane of, 295
Diaphragm, 133
Diarthrosis, 93
Digestion, organs of, 216
Disk, optic, 298
Discus proligerus, 285
Dorsum of scapula, 76
Douglas's cul-de-sac, 281, 283
Duct, cystic, 238
    hepatic, 237
    lymphatic, 184
    nasal, 309
    pancreatic, 238
    of Bartholin, 221
    of Gærtner, 286
    of Müller, 277
    of Rivinus, 221
    of Steno, 221
    of Wharton, 221
    of Wirsung, 238
    thoracic, 184
Ducts, biliary, 237
    ejaculatory, 278
Ductus cochlearis, 322
    communis choledochus, 238
Duodenum, 229
Dura mater of cerebrum, 186
        of spinal cord, 199

EAR, 310
    internal, 318
Earstones, 321
Eminence, hypothenar, 141
    thenar, 141
Eminentia collateralis (pes accessorius), 194
Enamel, 218
Enarthrosis, 93
Endocardium, 157
Endolymph, 318
Endosteum, 19
Epidermis, 288
Epididymis, 276
Epigastric region, 245
Epiglottis, 251
Epiphysis, 22
Equator of eyeball, 292
Eustachian tube, 317
Eye, 292
    its appendages, 306

Eyebrows, 306
Eyelashes, 307
Eyelids, 306

FALLOPIAN tubes, 284
  Fallopius, aqueduct of, 313
    hiatus of, 36
Falx cerebelli, 186
  cerebri, 186
Fascia, anal, 333
  Buck's, 334
  cremasteric, 327
  cribriform, 329
  dentata, 194
  intercolumnar, 326
  ischio-rectal, 333
  lata, 142
  lumbar, 131
  obturator, 333
  palmar, 138
  perineal, 333
  plantar, 150
  recto-vesical, 333
Fasciæ, 114
  of the perineum, 334
Fauces, 221
Femur, 85
Fenestra ovalis, 313
  rotunda, 313
Ferrein, pyramids of, 263
Fibro-cartilage, 94, 97, 102, 107, 110, 307, 310
Fibula, 89
Fimbriæ, 284
Fimbriated extremity, 284
Fissure, auricular, 37
  calcarine, 188
  calloso-marginal, 188
  collateral, 188
  dentate, 189
  Glaserian, 35
  palpebral, 306
  parieto-occipital, 187
  pterygo-maxillary, 60
  of Rolando, 187
  of Sylvius, 187
  precentral, 187
  spheno-maxillary, 60
  sphenoidal, 53
  transverse, 189
Fissures of cerebellum, 198
  of cerebrum, 187
  of liver, 234
Fissures of spinal cord, 199
Folds, palpebral, 307
Fontana, spaces of, 297
Fontanelles, 52
Foot, 90
Foramen for Arnold's nerve, 37
  cæcum, 24
  condyloid, 28
  ethmoidal, 25
  for Jacobson's nerve, 37
  infra-orbital, 41
  inter-vertebral, 62
  lacerum anterius, 33, 53
  magnum, 27, 54
  mastoid, 35
  medium, 54
  mental, 49
  nutrient, 19
  obturator, 68
  of Monro, 193
  of Scarpa, 43
  of Stenson, 43
  of Winslow, 247
  olfactory, 38
  ovale, 31
  optic, 33
  orbital, 32
  palatine, 43
  petrosal, 36
  posterius, 54
  pterygo-palatine, 47
  rotundum, 31
  sacral, 71
  spheno-palatine, 46
  spinal, 63
  spinosum, 32
  stylo-mastoid, 37
  supra-orbital, 24
  supra-scapular, 77
  thyroid, 68
  Vesalii, 32
Foramina at the base of the skull, 54
  olfactory, 38
  orbital, 32
  spheno-maxillary, 60
  Thebesii, 154
Forearm, 80
Fornix, 193
Fossa, canine, 40
  coronoid, 80
  digastric, 35
  digital, 86
  glenoid, 78
  hyaloid, 301

Fossa, incisive, 40
  infra-spinous, 77
  ischio-rectal, 330
  jugular, 29
  lachrymal, 25
  navicularis of vulva, 281
    of penis, 270
  olecranon, 80
  ovalis, 154
  pterygoid, 33
  scaphoid, 33
  spheno-maxillary, 61
  sublingual, 50
  submaxillary, 50
  subscapular, 76
  supraspinous, 77
  temporal, 59
  zygomatic, 60
Fossæ, nasal, 58
  of the skull, 53
Fourchette, 281
Fovea centralis, 298
  hemispherica, 319
  semi-elliptica, 319
Frænulum, 196
Frænum preputii, 274

GÆRTNER, duct of, 286
Gall-bladder, 238
Ganglia, basic, 191
  cervical, 213
  of fifth nerve, 202
  of the sympathetic, 211–214
Ganglion, Arnold's, 211, 212
  Gasserian, 202
  impar, 211
  Meckel's, 212
  of ribs, 211
  spheno-palatine, 212
  spirale, 320, 325
Gasser, ganglion of, 202
Gimbernat's ligament, 326
Ginglymus, 93
Gladiolus, 65
Gland, lachrymal, 308
  mammary, 286
  pineal, 196
  prostate, 271
  thymus, 242
  thyroid, 241
Glands, Brunner's, 230
  Cowper's, 272
  ductless, 239
Glands, gastric, 228
  intestinal, 230
  lachrymal, 308
  lymphatic, 184
  Meibomian, 307
  of Bartholin, 280
  of Tyson, 274
  sebaceous, 289
  solitary, 230
  sweat, 289
Glans clitoridis, 280
  penis, 273
Glenoid, 78
Glisson's capsule, 235
Globus major, 276
  minor, 276
Glottis, 252
Gomphosis, 93
Graafian vesicle, 285
Groove, bicipital, 79
  infraorbital, 41
  musculo-spiral, 79
  mylo-hyoid, 51
  naso-palatine, 49
  optic, 30
  pterygo-palatine, 47
Gubernaculum testis, 278
Gums, 217
Gyrus, angular, 189
  fornicatus, 189

HAIRS, 288
Hamstrings, 146
Hamular process, 33
Hand, 82
Haversian canals, 18
  system, 18
Head of the humerus, 79
  of the ulna, 81
Heart, 153
Helicotrema, 320
Helix, 310
Henle, tubes of, 261, 263
Hernia, femoral, 328
  inguinal, 325
Hey, ligament of, 329
Hiatus Fallopii, 36
Hilum of kidney, 261
Hippocampus major, 193
  minor, 193
Horner's muscle, 117
Houston, folds of, 232
Huguier, canal of, 35

Humerus, 78
Humor, aqueous, 300
  vitreous, 301
Humors of the eyeball, 300
Hunter's canal, 144
Hydatids of Morgagni, 277
Hymen, 280
Hypochondrium, 244, 246
Hypogastrium, 246

ILEUM, 69, 230
  Ilium, 69
Incisura intertragica, 310
Infundibula of kidney, 261
Infundibulum of brain, 191
Inguinal regions, 245, 246
Intestine, large, 231
  small, 229
Internal capsule, 194
Intumescentia ganglioniformis Scarpæ of seventh nerve, 325
Iris, 296
Ischium, 70
Iter chordæ anterius, 313
    posterius, 313
  e tertio ad quartum ventriculum, 194
Ivory, 218

JACOB'S membrane, 299
  Jacobson's nerve, 203
Jejunum, 229
Joint, ankle-, 111
  elbow-, 103
  hip-, 107
    rotators of the, 144–146
  knee-, 108
  shoulder-, 102
  wrist-, 105
Joints, motion in, 93
  structures of, 93

KIDNEYS, 260

LABIA majora, 279
  minora, 279
Labium tympanicum, 323
Labium vestibulare, 323
Labyrinth, membranous, 321
  osseous, 318
Lacteals, 184
Lacunus magnus, 270
Lacunæ, 18
Lacus lachrymalis, 306
Lamellæ, 18
Lamina cinerea, 191
  cribrosa (of sclerotic), 293
  fusca, 294
  reticularis, 324
  spiralis, 320
Laminæ of cornea, 295
Lancisi, nerves of, 193
Larynx, 249
Layer, dermoid, 288
  ganglionic, 299
  molecular, 299, 300
Leg, 88
Lens, crystalline, 301
Lieberkühn, crypts or follicles of, 230
Ligament, annular, of foot, 149
  annular of hand, 139
  conoid, 102
  coronary, 110
  costo-xiphoid, 98
  cotyloid, 107
  crucial, of knee-joint, 110
  deltoid, 112
  falciform, 329
  Gimbernat's, 326
  glenoid, 102
  ilio-femoral or Y, 107
  of Burns, 329
  of Hey, 326
  of Zinn, 301
  orbicular, 104
  Poupart's, 129, 326
  rhomboid, 101
  round, of the liver, 234
  round, of the uterus, 283
  sacro-sciatic, 100
  stellate, 97
  stylo-maxillary, 97
  suspensory, of lens, 301
    of liver, 234
    of penis, 273
    of spleen, 240
  transverse, of hip-joint, 107
  trapezoid, 102
  triangular, of perineum, 334
Ligaments of the bladder, 267
  broad, of uterus, 283

Ligaments, check, 96
  of the knee-joint, 108
  of the larynx, 251
  of the liver, 234
  of the ossicles, 315
  of the ovaries, 284
  of the scapula, 102
  of the sternum, 99
  of the uterus, 283
  peritoneal, 248
  tarsal, 307
Ligamentum latum pulmonis, 259
  mucosum, 110
  nuchæ, 124
  patellæ, 108
  pectinatum iridis, 297
  suspensorium, 96.
  teres, 107
  Winslowii, 109
Limbus laminæ spiralis, 323
Line, inter-trochanteric, 86
Linea aspera, 86
  ilio-pectinea, 70
  quadrati, 86
Liquor Morgagni, 301
Lithotomy, structures affected, 334
Liver, fissures of, 234
  lobes of, 235
  structure of, 237
  vessels of, 235
Lobe of the ear, 310
Lobes of the cerebellum, 197
  of the cerebrum, 187
  of the liver, 235
  optic, 192
Lobule of the ear, 310
Lobules of the liver, 237
  of the lung, 258
Lobulus caudatus, 235
  quadratus, 235
  Spigelii, 235
Lower, tubercle of, 154
Lumbar regions, 245, 246
Lung, broad ligament of, 259
Lungs, 256
Lunula, 288
Lymphatics, 184
  hepatic, 237
  of lung, 259
  of penis, 274
  of thymus gland, 242
  of trachea and bronchi, 256
  of uterus, 284
Lymph-spaces of eyeball, 304
Lyra, 193

MACULA acoustica, 322
  lutea, 298
Maculæ cribrosæ, 325
Malleolus, 88, 89
Malpighi, bodies of, 240
  pyramids of, 261
Malpighian corpuscles, 263
  tuft, 263
Mammæ, 286
Manubrium of the malleus, 314
  of the sternum, 65
Marrow of bone, 19
Masses, lateral, of ethmoid, 39
Mastoid process, 85
Meatus auditorius externus, 36
    internus, 36
  nasi, 59
  urinarius, 273, 280
Meckel's ganglion, 212
Mediastinum, 260
  testis, 275
Medulla oblongata, 196
Medullary substance, 261
Membrana basilaris, 323
  granulosa, 285
  pupillaris, 298
  Ruyschiana, 296
  sacciformis, 104
  tectoria, 323
  tympani, 311
    secundaria, 321
Membrane, hyaloid, 301
  interosseous, 104, 111
  Jacob's, 300
  limiting, 299, 300
  of Bowman, 295
  of Descemet, 295
  of Reissner, 323
  pituitary, 290
  Schneiderian, 290
  Shrapnell's, 312
  vitelline, 285
Membranes of brain, 186
  of the spinal cord, 199
  synovial, 93
  tarsal synovial, 114
  wrist synovial, 105
Meridians of eyeball, 292
Mesenteries, 248
Meso-cephalon (pons Varolii), 195

Metacarpus, 84
Metatarsus, 91
Modiolus, 320
Monro, foramen of, 193
Mons veneris, 279
Morgagni, hydatids of, 277
Mouth, 220
Müller, duct of, 277
  muscle of, 296
Muscle of Bowman (ciliary), 296
  ciliary, 296
  cremaster, 327
  dilator pupillæ, 298
  Horner's, 117
  kerato-cricoideus, 253
  levator glandulæ thyroideæ, 241
  of Müller, 296
  sphincter pupillæ, 297
  triticeo-glossus, 253
  of the arm, 134
  of the auricle, 310
  of the back, 123
Muscles, double-bellied, 116
  of the abdomen, 129
  of the ear (external), 116
  of the epiglottis, 253
  of the Eustachian tube, 317
  of the eyeball, 302
  of the face, 117, 118, 119
  of the forearm, 136
  of the foot, 150
  of the hand, 140
  of the head, 111
  of the hip, 144, 145
  of the larynx, 253
  of the leg, 147
  of the neck, 120
  of the palate, 122
  of the perineum, 331
  of the pharynx, 225
  of the shoulder, 134
  of the stomach, 227
  of the thigh, 142
  of the thorax, 132
  of the tympanum, 315
Muscular fibre, 114
Musculi pectinati, 154

NABOTH, ovules of, 284
Nails, 288
Nates of cerebrum, 196
Neck of humerus, 79
Nerve tissue, 185
Nerve, abducens, 202
  Arnold's, 37, 204
  auditory, 203
  Bell's external respiratory, 206
    internal respiratory, 205
  chorda tympani, 202, 313, 316
  circumflex, 206
  cochlear, 203, 320
  crural anterior, 209
  facial, 202
  genito-crural, 208
  glosso-pharyngeal, 203
  hypoglossal, 204
  ilio-hypogastric, 208
  ilio-inguinal, 208
  intercosto-humeral, 206, 208
  interosseous, 208
  Jacobson's, 37, 203, 316
  laryngeal, 204, 254
  maxillary, inferior, 202
    superior, 202
  median, 206
  motor oculi, 201
  musculo-spiral, 207
  nonus or ninth pair, 204
  obturator, 209
  of Wrisberg, 206
  olfactory, 200
  ophthalmic division of fifth, 202
  optic, 200
  par vagum, 204
  patheticus, 201
  phrenic, 205
  pneumogastric, 204
  portio dura, 202
  portio mollis, 203
  pudic, 209
  radial, 207
  spinal accessory, 204
  sympathetic, 211
  thoracic, 206
  trigeminus, 201
  tympanic, 203
  ulnar, 207
  vestibular, 203, 324
  Vidian, 213
Nerve-fibres, optic, 200, 305
Nerves of ankle-joint, 112
  of auditory canal, 311
  of auricle, 311
  of bladder, 269
  of choroid, 296

Nerves, cranial, 200
  hepatic, 237
  of Eustachian tube, 318
  of elbow-joint, 104
  of eye, 304
  of eyelids, 304
  of eye-muscles, 303
  of hip-joint, 107
  of iris, 298
  of kidney, 265
  of knee-joint, 110
  of Lancisi, 193
  of larynx, 204, 254
  of lung, 259
  of membrana tympani, 312
  of nasal fossæ, 291
  of nose, 291
  of œsophagus, 227
  of ovaries, 286
  of pancreas, 239
  of penis, 274
  of pharynx, 226
  of shoulder-joint, 102
  of spleen, 241
  of stomach, 229
  of supra-renal capsules, 243
  of taste, 222, 290
  of testicles, 278
  of thymus gland, 242
  of tongue, 222
  of trachea and bronchi, 256
  of tympanum, 316
  of uterus, 284
  of wrist-joint, 105
  petrosal, 213
  popliteal, 210
  sciatic, 209
  spinal, 205
  splanchnics, 214
  thyroid, 241
Nervous system, 185
Neurilemma, 185
Nipple, 286
Nose, 290
Notch, cotyloid, 68
  inter-clavicular, 65
  inter-condyloid, 87
  popliteal, 88
  sacro-sciatic, 70
  sigmoid, 49
  supra-orbital, 24
  supra-scapular, 77
Nuck, canal of, 283
Nymphæ, 279

ODONTOBLASTS, 220
Odontoid ligaments, 95, 96
    process, 63
Œsophagus, 226
Omentum, 247
Opening, saphenous, 329
Opercula, 220
Optic thalami, 192
Ora serrata, 298
Orbital cavities, 55
Organ of Corti, 323
  of hearing, 310
  of Rosenmüller, 286
  of sight, 292
  of smell, 290
Organs of generation, female, 279
    male, 271
  of sense, 287
  of taste, 289
  of voice and respiration, 249
  urinary, 260
Os calcis, 90
  orbiculare, 314
  planum (orbital of ethmoid), 39
  tincæ, 282
  uteri, 282
Ossa innominata, 68
  triquetra, 20
Osteology, 17
Ossification, 20
Ossicles of tympanum, 314
Otoliths, 321
Ovaries, 284
Oviducts, 284
Ovisacs, 285
Ovules of Naboth, 284
Ovum, 285

PACCHIONIAN bodies, 186
Palate, 221
Palpebræ, 306
Pampiniform plexus, 278, 286
Pancreas, 238
  lesser, 239
Papilla, lachrymal, 306
  optic, 298
Papillæ of skin, 288
Parovarium, 286
Pars ciliaris retinæ, 299
    intermedia, of vulva, 281
Peduncles of cerebellum, 195, 196, 197
Pelvis, 73

- Pelvis of the kidney, 261
- Penis, 272
- Pericardium, 152
- Perilymph, 318
- Perineum, 331
  - female, 335
- Periosteum, 19
- Peritoneum, 246
- Peronei muscles, 147, 149
- Pes accessorius, 194
  - hippocampi, 194
- Petit, canal of, 302
- Peyer's patches, 230
- Phalanges of ear, 324
  - of foot, 92
  - of hand, 85
- Pharynx, 224
- Pia mater of brain, 187
  - of spinal cord, 199
  - of the testes, 275
- Pinna, or auricle, 310
- Plane, equatorial, 292
  - meridional, 292
- Plantar arch, 175
- Plate, cribriform, 38
  - orbital, 39, 41, 44
  - perpendicular, 38
- Pleuræ, 259
- Plexus, brachial, 206
  - carotid, 214
  - cavernous, 214
  - cervical, 205
  - choroid, 193
  - gastric, 215
  - lumbar, 208
  - pampiniform, 278, 286
  - sacral, 209
  - solar, 215
  - splenic, 215
  - tympanic, 316
- Plexuses of the sympathetic, 214, 215, 216
- Pli courbe (angular gyrus), 252
- Plica semilunaris, 307
- Point, nodal, 292
- Poles of the eyeball, 292
- Pomum Adami, 249
- Pons Varolii, 195
- Portal system, 178, 235
- Porus opticus, 293
- Pouches of membrana tympani, 312
- Poupart's ligament, 129, 326
- Prepuce of clitoris, 280
  - of penis, 273
- Process, acromion, 77
  - alveolar, 43, 50
  - articular, 21
  - auditory, 36
  - basilar, 28
  - clinoid, 30, 33
  - condyloid, 28, 49, 79, 85, 86
  - coracoid, 77
  - coronoid, 50, 81
  - ethmoidal, 48
  - falciform, 329
  - hamular, of cochlea, 320
    - of pterygoid plate of sphenoid, 33
  - lachrymal, 48
  - malar, 42
  - mastoid, 35
  - maxillary, 48
  - mental, 49
  - nasal, 42
  - odontoid, 63
  - olecranon, 80
  - olivary, 30
  - orbital, 39, 41, 45, 46
  - palatine, 43
  - pterygoid, 33
  - sphenoidal, 46
  - spinous, 32, 43, 47, 63, 70, 71
  - styloid, 37
  - transverse, 63
  - turbinated, 39
  - unciform, 39
  - vaginal, 37
  - vermiform, 197
  - zygomatic, 34
- Processes, ciliary, 296
  - clinoid, 30, 33
  - of Ingrassias, 32
- Processus cochleariformis, 37
  - cerebello ad medullam (inferior peduncles), 197
    - ad pontem (middle peduncles), 195
    - ad testes, 196
  - gracilis, 314
- Promontory of sacrum, 71
  - of tympanum, 313
- Prostate gland, 271
- Prostatic urethra, 269
  - sinus, 269
- Pterygoid processes, 33
- Pubes, 70
- Punctum lachrymale, 306
- Pupil, 297

Pylorus, 227
Pyramid, anterior, 196
  posterior, 197
  of tympanum, 314
Pyramids of Ferrein, 263
  of Malpighi, 261

RADIUS, 81
Rami of inferior maxillary, 50
Ramus of ischium, 70
  of the pubes, 70
Receptaculum chyli, 184
Rectum, 232
Regions of abdomen, 244
Reil, island of, 188
Reissner, membrane of, 323
Renal vessels, 263
Rete mucosum, 288
Rete testis, 276
Retina, 298
Ribes, ganglion of, 211
Ribs, 66
Ridge, mylo-hyoid, 51
  pterygoid, 32
Rima glottidis, 252
Ring, abdominal, 326
  crural, 328
  femoral, 328
Rings and phalanges, 324
Rivinus, duct of, 221, 222
  notch of, 311
Rods and cones (Jacob's), 300
    of Corti, 323
Rolando, fissure of, 187
Rosenmüller, organ of, 286
Rostrum of the sphenoid, 31

SAC, lachrymal, 309
Saccule, 322
Sacculus laryngis, 252
Sacs of the peritoneum, 246
Sacrum, 71
Santorini, cartilages of (cornicula laryngis), 250
Scala media, 322
  tympani, 321
  vestibuli, 320
Scapula, 76
Scarpa, foramen of, 43
  intumescentia ganglioniformis, 325
Schindelysis, 93
Schlemm, canal of, 297
Schultze, olfactory cells of, 291
Schwann, substance of, 185
Sclerotic, 293
Scrotum, 275
Sella turcica, 30
Septum crurale, 329
  lucidum, 193
  nasi, 58
  pectiniforme, 273
Sheath, crural, 329
Shin, 88
Shoulder-joint, 102
  -blade, 76
Shrapnell's membrane, 312
Sight, organs of, 292
Sigmoid flexure, 232
Sylvius, aqueduct of, 36, 203, 313
  fissure of, 187
Sinus aortici or sinus of Valsalva, 156
  circularis iridis, 297
  of the kidney, 261
  of Morgagni (sacculus laryngis), 252
  pocularis, 270
Sinuses, 23
  sphenoidal, 31
  of Valsalva, 155, 156
Skin, 287
Slit, nasal, 38, 58
Smell, organ of, 290
Space, anterior perforated, 191
  posterior perforated, 191
Spaces, Fontana's, 297
Spermatic cord, 278
Sphincter vesicæ, 268
Spinal cord, 198
Spine, ethmoidal, 30
  of the ischium, 70
  nasal anterior, 43, 58
    posterior, 47
  of the pubes, 71
  of the scapula, 76
  of the sphenoid, 32
  of the tibia, 88
Spines of the ilium, 69
Spinous processes, 21
Spleen, 239
Spot, blind, 298
  germinal, 285
  light, 312
  yellow, 298
Squamous, 22

Stars of Verheyen, 264
Steno's ducts, 221
Stenson's foramina, 43
Sternum, 65
Stilling, canal of, 301
Styloid, 22
Stomach, 227
Substance of Schwann, 185
Sulci, 189
Sulcus spiralis, 323
  tympanicus, 311
Surface, auricular, 70
  trochlear, 79
Sustentaculum tali, 90
Sutura, 92
Suture, baso-sphenoidal, 52
  coronal, 52
  lachrymo-ethmo-frontal, 52
  lambdoidal, 52
  masto-occipital, 52
  masto-parietal, 52
  petro-occipital, 52
  petro-sphenoidal, 52
  sagittal, 52
  spheno-parietal, 52
  squamo-parietal. 52
  squamo-sphenoidal, 52
Sutures, facial, 52
  of the orbit, 56, 57
  of the skull, 51
Sympathetic system, 211
Symphysis, 92
  of inferior maxillary, 49
Synarthrosis, 92
Synchondrosis, 92
Synovial membranes, 93

TARSUS, 90
Taste, nerves of, 290
Taste-goblets. 289
Tænia hippocampi, 193
  semicircularis, 193
Teeth, 217
Tendo-oculi, 307
Tendons, 115
Tenon, capsule of, 292
Tentorium cerebelli, 186
Testes, 274
  of the cerebrum, 196
  descent of, 278
Thebesius, foramina of, 154
Thigh, 85
Thorax, 64
Thyroid axis, 164
  cartilage, 249
Tongue, 222
Tonsils, 221
Trachea, 254
Tract, lateral, 197
  optic, 305
  uveal, 297
Tracts, olfactory, 200
Tractus spiralis foraminulentus, 325
Tragus, 310
Trigonum vesicæ, 268
Trochanters of femur, 85
Trochlea, 22
Tube, Eustachian, 317
Tuber annulare (pons Varolii), 195
  cinereum, 191
Tubercle, 21
  genial, 50
  of Lower, 154
Tuberosities, 21
  of femur, 87
  of humerus, 79
  of ischium, 70
  maxillary, 49
Tubes of Henle (convoluted tubes), 261
  spiral, of Schachowa, 262
Tubuli lactiferi, 286
  seminiferi, 276
  uriniferi, 262
Tuft, Malpighian, 263
Tunica albuginea, 275
  vaginalis, 275
  vasculosa, 275
Tunics of eyeball, 293
Tutamina oculi, 306
Tympanum. 313
Tyson, glands of, 274

ULNA, 80
Umbilical region, 245
Umbilicus, 244
Umbo, 312
Urachus, 267
Ureters, 265
Urethra, 271
  bulb of, 270
Uterus, 281
  masculinus, 270
Utricle, 321

Uvea, 297
Uvula, 221
    vesicæ, 269

VAGINA, 281
    Vaginal, 22
Valley of cerebellum, 197
Valsalva, sinuses of, 155, 156
Valve, Eustachian, 154
    ilio-cæcal, 231
    mitral, 156
    tricuspid, 155
    of Bauhin, 231
    of Vieussens, 196
Valves of Kirkring (valvulæ conniventes), 230
    semilunar, 155
Valvulæ conniventes, 230
Vas aberrens, 276
    deferens, 277
Vasa efferentia, 276
    recta, 276
    vasorum, 158
Vein, axillary, 179
    basilic, 179
    cephalic, 179
    jugular, 178
    femoral, 182
    iliac, 182
    median, 179
    popliteal, 182
    portal, 183
    radial, 179
    splenic, 183
    subclavian, 179
    supra-renal, 183
    ulnar, 179
    vena cava, 180, 183
    vertebral, 181
Veins, 177
    azygos, 180
    cardiac, 183
    choroid, 296
    hepatic, 183
    iliac, 182
    innominate, 180
    interlobular, 264
    jugular, 178
    of bone, 19
    of head and neck, 178
    of lower extremities, 182
    of œsophagus, 226
    of supra-renal capsules, 243
Veins of thymus, 242
    of trachea and bronchi, 256
    of the eye, 296, 298
    of the kidney, 265
    of the larynx, 254
    of the lung, 259
    of the pancreas, 239
    of the penis, 274
    of the stomach, 229
    of the thorax, 181
    pulmonary, 183
    saphenous, 182
    spinal, 181
    thyroid, 241
    uterine, 284
    without valves, 178
Velum interpositum, 187, 193, 194
Venæ comites, 177
    propriæ renales, 265
    rectæ, 264
    vorticosæ, 296
Venter of the ilium, 70
    of the scapula, 76
Ventricle of the cord, 200
    fifth, 194
    fourth, 195
    of the larynx, 252
    third, 194
Ventricles, lateral, 192
Verheyen, stars of, 264
Vertebra prominens, 64
Vertebræ, 61
Verumontanum, 269
Vesaluis, foramen of, 32
Vesicle, germinal, 285
    Graafian, 285
Vesiculæ seminales, 278
Vestibule, 280
    of the ear, 318
Vidian canal, 33
    nerve, 213
    Vieussens, valve of, 196
Villi, 230
Viscus, 216
Viscera, covered by peritoneum, 216
        partially, 248
        wholly, 248
Vitellus, 285
Vocal cords, 252
Vomer, 48
Vortex of heart, 157
Vulva, 279

WALLS of the tympanum, 313
Wharton's ducts, 221
Whorl of heart, 157
Willis, circle of, 163
Wings of the sphenoid, 31, 32
Winslow, foramen of, 247
    ligament of, 109
Wirsung, duct of, 238
Wormian bones, 20
Wrisberg, cartilages of, 251
    nerve of, 206

YELK, 285

ZINN, zonula of, 301
Zona pellucida, 285
Zone, anterior vascular, 294
Zonula of Zinn, 301
Zygoma, 34, 60

Standard Medical and Surgical Works

PUBLISHED BY

W. B. SAUNDERS, 925 Walnut Street, Philadelphia, Pa.

	PAGE
*American Text-Book of Applied Therapeutics	8
*American Text-Book of Diseases of Children	3
*American Text-Book of Gynecology	4
*American Text-Book of Nursing	8
*American Text-Book of Obstetrics	8
*American Text-Book of Physiology	8
*American Text-Book of Practice	2
*American Text-Book of Surgery	1
Ashton's Obstetrics	23
Ball's Bacteriology	27
Bastin's Laboratory Exercises in Botany	18
Beck's Surgical Asepsis	12
Brockway's Physics	27
Burr's Nervous Diseases	12
Cerna's Notes on the Newer Remedies	18
Chapman's Medical Jurisprudence and Toxicology	14
Cohen and Eshner's Diagnosis	26
Cragin's Gynæcology	24
DaCosta's Manual of Surgery	13
*De Schweinitz's Diseases of the Eye	5
Dorland's Obstetrics	13
Frothingham's Guide to Bacteriological Laboratory	14
Garrigues' Diseases of Women	10
Gleason's Diseases of the Ear	28
Griffin's Materia Medica and Therapeutics	12
*Gross's Autobiography	7
Hare's Physiology	22
Hampton's Nursing: its Principles and Practice	15
Hyde's Syphilis and Venereal Diseases	12
Jackson and Gleason's Diseases of the Eye, Nose, and Throat	25
Jewett's Outlines of Obstetrics	18
*Keating's Pronouncing Dictionary of Medicine	7
Keating's How to Examine for Life Insurance	20
Keen's Operation Blanks	16
Kyle's Diseases of Nose and Throat	12
Laine's Temperature Charts	9
Lockwood's Practice of Medicine	12
Long's Syllabus of Gynecology	9
Martin's Surgery	22
Martin's Minor Surgery, Bandaging, and Venereal Diseases	25
Morris' Materia Medica and Therapeutics	23
Morris' Practice of Medicine	24
Morton's Nurses' Dictionary	9
Nancrede's Anatomy and Manual of Dissection	16
Nancrede's Anatomy	22
Norris' Syllabus of Obstetrical Lectures	17
Powell's Diseases of Children	26
Raymond's Physiology	13
Saunders' Pocket Medical Formulary	19
Saunders' Pocket Medical Lexicon	19
Saunders' New Aid Series of Manuals	11, 12
Saunders' Series of Question Compends	21
Sayre's Practice of Pharmacy	26
Semple's Pathology and Morbid Anatomy	23
Semple's Legal Medicine, Toxicology, and Hygiene	25
Senn's Syllabus of Lectures on Surgery	17
Shaw's Nervous Diseases and Insanity	27
Stelwagon's Diseases of the Skin	24
Stevens' Materia Medica and Therapeutics	20
Stevens' Practice of Medicine	17
Stewart and Lawrance's Medical Electricity	28
Thornton's Dose-Book and Manual of Prescription-Writing	14
*Vierordt and Stuart's Medical Diagnosis	6
Warren's Surgical Pathology	10
Wilson's Orthopædic Surgery	15
Wolff's Chemistry	23
Wolff's Examination of Urine	26

MR. SAUNDERS, in presenting to the profession the following list of his publications, begs to state that the aim has been to make them worthy of the confidence of medical book-buyers by the high standard of *authorship* and by the excellence of *typography, paper, printing,* and *binding*.

The works indicated thus (*) are sold by SUBSCRIPTION (*not by booksellers*), usually through travelling solicitors, but they can be obtained *direct* from the office of publication (charges of shipment prepaid) by remitting the quoted prices. Full *descriptive cirulars* of such works will be sent to any address upon application.

All the other BOOKS advertised in this catalogue are commonly for sale by *booksellers* in all parts of the United States; but any book will be sent by the publisher to any address (post-paid) on receipt of the price herein given.

(*For Announcement of Forthcoming Publications see next page.*)

Announcement of Forthcoming Publications.

AN AMERICAN TEXT-BOOK OF OBSTETRICS. By American Teachers. (See page 8.)

AN AMERICAN TEXT-BOOK OF PHYSIOLOGY. By American Teachers. (See page 8.)

AN AMERICAN TEXT-BOOK OF APPLIED THERAPEUTICS. By American Teachers.

AN AMERICAN TEXT-BOOK OF NURSING. By American Teachers.

SURGICAL PATHOLOGY AND THERAPEUTICS. By J. Collins Warren, M. D., Professor of Surgery, Harvard Medical School, etc. (See page 10.)

A SYLLABUS OF GYNÆCOLOGY, arranged in conformity with The American Text-Book of Gynecology. By J. W. Long, M. D., Professor of Diseases of Women and Children, Medical College of Virginia, etc. (See page 9.)

TEMPERATURE CHART. Prepared by D. T. Lainé, M. D. (See page 9.)

LABORATORY EXERCISES IN BOTANY. By Edson S. Bastin, M. A., Professor of Materia Medica and Botany in the Philadelphia College of Pharmacy. (See page 18.)

A GUIDE TO THE BACTERIOLOGICAL LABORATORY. By Langdon Frothingham, M. D (See page 14.)

SAUNDERS' NEW AID SERIES OF MANUALS.

New volumes in active preparation. See pages 11, 12.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF SURGERY. Edited by WILLIAM W. KEEN, M. D., LL.D., and J. WILLIAM WHITE, M. D., PH. D. Forming one handsome royal-octavo volume of over 1200 pages (10×7 inches), with nearly 500 wood-cuts in text, and 37 colored and half-tone plates, many of them engraved from original photographs and drawings furnished by the authors. Prices: Cloth, $7.00 net; Sheep, $8.00 net; Half Russia, $9.00 net.

The want of a text-book which could be used by the practitioner and at the same time be recommended to the medical student has been deeply felt, especially by teachers of surgery; hence, when it was suggested to a number of these that it would be well to unite in preparing a text-book of this description, great unanimity of opinion was found to exist, and the gentlemen below named gladly consented to join in its production. While there is no distinctive American Surgery, yet America has contributed very largely to the progress of modern surgery, and among the foremost of those who have aided in developing this art and science will be found the authors of the present volume. All of them are teachers of surgery in leading medical schools and hospitals in the United States and Canada.

Especial prominence has been given to Surgical Bacteriology, a feature which is believed to be unique in a surgical text-book in the English language. Asepsis and Antisepsis have received particular attention. The text is brought well up to date in such important branches as cerebral, spinal, intestinal, and pelvic surgery, the most important and newest operations in these departments being described and illustrated.

The text of the entire book has been submitted to all the authors for their mutual criticism and revision—an idea in book-making that is entirely new and original. The book as a whole, therefore, expresses on all the important surgical topics of the day the consensus of opinion of the eminent surgeons who have joined in its preparation.

One of the most attractive features of the book is its illustrations. Very many of them are original and faithful reproductions of photographs taken directly from patients or from specimens, and the modern improvements in the art of engraving have enabled the publisher to produce illustrations which it is believed are superior to those in any similar work.

CONTRIBUTORS:

Dr. Charles H. Burnett, Philadelphia.
Phineas S. Conner, Cincinnati.
Frederic S. Dennis, New York.
William W. Keen, Philadelphia.
Charles B. Nancrede, Ann Arbor, Mich.
Roswell Park, Buffalo, N. Y.
Lewis S. Pilcher, New York.

Dr. Nicholas Senn, Chicago.
Francis J. Shepherd, Montreal, Canada.
Lewis A. Stimson, New York.
William Thomson, Philadelphia.
J. Collins Warren, Boston.
J. William White, Philadelphia.

"If this text-book is a fair reflex of the present position of American surgery, we must admit it is of a very high order of merit, and that English surgeons will have to look very carefully to their laurels if they are to preserve a position in the van of surgical practice."—*London Lancet.*

"The soundness of the teachings contained in this work needs no stronger guarantee than is afforded by the names of its authors."—*Medical News,* Philadelphia.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK ON THE THEORY AND PRACTICE OF MEDICINE. By American Teachers. Edited by WILLIAM PEPPER, M. D., LL.D., Provost and Professor of the Theory and Practice of Medicine and of Clinical Medicine in the University of Pennsylvania. Complete in two handsome royal-octavo volumes of about 1000 pages each, with illustrations to elucidate the text wherever necessary. Price per Volume: Cloth, $5.00 net; Sheep, $6.00 net; Half Russia, $7.00 net.

VOLUME I. CONTAINS:

Hygiene.—Fevers (Ephemeral, Simple Continued, Typhus, Typhoid, Epidemic Cerebrospinal Meningitis, and Relapsing).—Scarlatina, Measles, Rötheln, Variola, Varioloid, Vaccinia, Varicella, Mumps, Whooping-cough, Anthrax, Hydrophobia, Trichinosis, Actinomycosis, Glanders, and Tetanus.—Tuberculosis, Scrofula, Syphilis, Diphtheria, Erysipelas, Malaria, Cholera, and Yellow Fever.—Nervous, Muscular, and Mental Diseases etc.

VOLUME II. CONTAINS:

Urine (Chemistry and Microscopy).—Kidney and Lungs.—Air-passages (Larynx and Bronchi) and Pleura.—Pharynx, Œsophagus, Stomach and Intestines (including Intestinal Parasites), Heart, Aorta, Arteries and Veins.—Peritoneum, Liver, and Pancreas.—Diathetic Diseases (Rheumatism, Rheumatoid Arthritis, Gout, Lithæmia, and Diabetes.)—Blood and Spleen.—Inflammation, Embolism, Thrombosis, Fever, and Bacteriology.

The articles are not written as though addressed to students in lectures, but are exhaustive descriptions of diseases, with the newest facts as regards Causation, Symptomatology, Diagnosis, Prognosis, and Treatment, including a large number of approved formulæ. The recent advances made in the study of the bacterial origin of various diseases are fully described, as well as the bearing of the knowledge so gained upon prevention and cure. The subjects of Bacteriology as a whole and of Immunity are fully considered in a separate section.

Methods of diagnosis are given the most minute and careful attention, thus enabling the reader to learn the very latest methods of investigation without consulting works specially devoted to the subject.

CONTRIBUTORS:

Dr. J. S. Billings, Philadelphia.
Francis Delafield, New York.
Reginald H. Fitz, Boston.
James W. Holland, Philadelphia.
Henry M. Lyman, Chicago.
William Osler, Baltimore.
Dr. William Pepper, Philadelphia.
W. Gilman Thompson, New York.
W. H. Welch, Baltimore.
James T. Whittaker, Cincinnati.
James C. Wilson, Philadelphia.
Horatio C. Wood, Philadelphia.

"We reviewed the first volume of this work, and said: 'It is undoubtedly one of the best text-books on the practice of medicine which we possess.' A consideration of the second and last volume leads us to modify that verdict and to say that the completed work is, in our opinion, THE BEST of its kind it has ever been our fortune to see. It is complete, thorough, accurate, and clear. It is well written, well arranged, well printed, well illustrated, and well bound. It is a model of what the modern text-book should be."—*New York Medical Journal.*

"A library upon modern medical art. The work must promote the wider diffusion of sound knowledge."—*American Lancet.*

"A trusty counsellor for the practitioner or senior student, on which he may implicitly rely."—*Edinburgh Medical Journal.*

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF THE DISEASES OF CHILDREN. By American Teachers. Edited by LOUIS STARR, M. D., assisted by THOMPSON S. WESTCOTT, M. D. In one handsome royal-8vo volume of 1190 pages, profusely illustrated with wood-cuts, half-tone and colored plates. Net Prices: Cloth, $7.00; Sheep, $8.00; Half Russia, $9.00.

The plan of this work embraces a series of original articles written by some sixty well-known pædiatrists, representing collectively the teachings of the most prominent medical schools and colleges of America. The work is intended to be a PRACTICAL book, suitable for constant and handy reference by the practitioner and the advanced student.

One decided innovation is the large number of authors, nearly every article being contributed by a specialist in the line on which he writes. This, while entailing considerable labor upon the editors, has resulted in the publication of a work THOROUGHLY NEW AND ABREAST OF THE TIMES.

Especial attention has been given to the latest accepted teachings upon the etiology, symptoms, pathology, diagnosis, and treatment of the disorders of children, with the introduction of many special formulæ and therapeutic procedures.

Special chapters embrace at unusual length the Diseases of the Eye, Ear, Nose and Throat, and the Skin; while the introductory chapters cover fully the important subjects of Diet, Hygiene, Exercise, Bathing, and the Chemistry of Food. Tracheotomy, Intubation, Circumcision, and such minor surgical procedures coming within the province of the medical practitioner are carefully considered.

CONTRIBUTORS:

Dr. S. S. Adams, Washington.
John Ashhurst, Jr., Philadelphia.
A. D. Blackader, Montreal, Canada.
Dillon Brown, New York.
Edward M. Buckingham, Boston.
Charles W. Burr, Philadelphia.
W. E. Casselberry, Chicago.
Henry Dwight Chapin, New York.
W. S. Christopher, Chicago.
Archibald Church, Chicago
Floyd M. Crandall, New York.
Andrew F. Currier, New York.
Roland G. Curtin, Philadelphia
J. M. DaCosta, Philadelphia.
I. N. Danforth, Chicago.
Edward P. Davis, Philadelphia.
John B. Deaver, Philadelphia.
G. E. de Schweinitz, Philadelphia.
John Dorning, New York.
Charles Warrington Earle, Chicago.
Wm. A. Edwards, San Diego, Cal.
F. Forchheimer, Cincinnati.
J. Henry Fruitnight, New York.
Landon Carter Gray, New York.
J. P. Crozer Griffith, Philadelphia.
W. A. Hardaway, St. Louis.
M. P Hatfield, Chicago.
Barton Cooke Hirst, Philadelphia.
H. Illoway, Cincinnati.
Henry Jackson, Boston.
Charles G. Jennings, Detroit.
Henry Koplik, New York.

Dr. Thomas S. Latimer, Baltimore.
Albert R. Leeds, Hoboken, N. J.
J. Hendrie Lloyd, Philadelphia.
George Roe Lockwood, New York.
Henry M. Lyman, Chicago.
Francis T. Miles, Baltimore.
Charles K. Mills, Philadelphia.
John H. Musser, Philadelphia.
Thomas R. Neilson, Philadelphia.
W. P. Northrup, New York.
William Osler, Baltimore.
Frederick A. Packard, Philadelphia.
William Pepper, Philadelphia.
Frederick Peterson, New York.
W. T. Plant, Syracuse, New York.
William M. Powell, Atlantic City.
B. Alexander Randall, Philadelphia.
Edward O. Shakespeare, Philadelphia.
F. C. Shattuck, Boston.
J. Lewis Smith, New York.
Louis Starr, Philadelphia.
M. Allen Starr, New York.
J. Madison Taylor, Philadelphia.
Charles W. Townsend, Boston.
James Tyson, Philadelphia.
W. S. Thayer, Baltimore.
Victor C. Vaughan, Ann Arbor, Mich.
Thompson S. Westcott, Philadelphia.
Henry R. Wharton, Philadelphia.
J. William White, Philadelphia.
J. C. Wilson, Philadelphia.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF GYNECOLOGY, MEDICAL AND SURGICAL, for the use of Students and Practitioners. Edited by J. M. BALDY, M. D. Forming a handsome royal-octavo volume, with 360 illustrations in text and 37 colored and half-tone plates. Prices: Cloth, $6.00 net; Sheep, $7.00 net; Half Russia, $8.00 net.

In this volume all anatomical descriptions, excepting those essential to a clear understanding of the text, have been omitted, the illustrations being largely depended upon to elucidate the anatomy of the parts. This work, which is thoroughly practical in its teachings, is intended, as its title implies, to be a working text-book for physicians and students. A clear line of treatment has been laid down in every case, and although no attempt has been made to discuss mooted points, still the most important of these have been noted and explained. The operations recommended are fully illustrated, so that the reader, having a picture of the procedure described in the text under his eye, cannot fail to grasp the idea. All extraneous matter and discussions have been carefully excluded, the attempt being made to allow no unnecessary details to cumber the text. The subject-matter is brought up to date at every point, and the work is as nearly as possible the combined opinions of the ten specialists who figure as the authors.

The work is well illustrated throughout with wood-cuts, half-tone and colored plates, mostly selected from the authors' private collections.

CONTRIBUTORS:

Dr. Henry T. Byford.
John M. Baldy.
Edwin Cragin.
J. H. Etheridge.
William Goodell.
Dr. Howard A. Kelly.
Florian Krug.
E. E. Montgomery.
William R. Pryor.
George M. Tuttle.

"The most notable contribution to gynecological literature since 1887, ... and the most complete exponent of gynecology which we have. No subject seems to have been neglected, ... and the gynecologist and surgeon, and the general practitioner who has any desire to practise diseases of women, will find it of practical value. In the matter of illustrations and plates the book surpasses anything we have seen."—*Boston Medical and Surgical Journal.*

"A valuable addition to the literature of Gynecology. The writers are progressive, aggressive, and earnest in their convictions."—*Medical News*, Philadelphia.

"A thoroughly modern text-book, and gives reliable and well-tempered advice and instruction."—*Edinburgh Medical Journal.*

"The harmony of its conclusions and the homogeneity of its style give it an individuality which suggests a single rather than a multiple authorship."—*Annals of Surgery.*

"It must command attention and respect as a worthy representation of our advanced clinical teaching."—*American Journal of Medical Sciences.*

For Sale by Subscription.

DISEASES OF THE EYE. A Handbook of Ophthalmic Practice. By G. E. DE SCHWEINITZ, M. D., Professor of Diseases of the Eye, Philadelphia Polyclinic; Professor of Clinical Ophthalmology, Jefferson Medical College, Philadelphia, etc. Forming a handsome royal-octavo volume of more than 600 pages, with over 200 fine wood-cuts, many of which are original, and 2 chromo-lithographic plates. Prices: Cloth, $4.00 net; Sheep, $5.00 net; Half Russia, $5.50 net.

The object of this work is to present to the student and practitioner who is beginning work in the fields of ophthalmology a plain description of the optical defects and diseases of the eye. To this end special attention has been paid to the clinical side of the question; and the method of examination, the symptomatology leading to a diagnosis, and the treatment of the various ocular defects have been brought into special prominence. The general plan of the book is eminently practical. Attention is called to the large number of illustrations (nearly one-third of which are new), which will materially facilitate the thorough understanding of the subject.

"For the student and practitioner it is the best single volume at present published."—*Medical News*, Philadelphia.

"A most complete and sterling presentation of the present status of modern knowledge concerning diseases of the eye."—*Medical Age.*

"Pre-eminently a book for those wishing a clear yet comprehensive and full knowledge of the fundamental truths which underlie and govern the practice of ophthalmology."—*Medical and Surgical Reporter.*

"At once comprehensive and thoroughly up to date."—*Hospital Gazette* (London).

PROFESSIONAL OPINIONS.

"A work that will meet the requirements not only of the specialist, but of the general practitioner in a rare degree. I am satisfied that unusual success awaits it."

WILLIAM PEPPER, M. D.,
Provost and Professor of Theory and Practice of Medicine and Clinical Medicine in the University of Pennsylvania.

"Contains in concise and reliable form the accepted views of Ophthalmic Science."

WILLIAM THOMSON, M. D.,
Professor of Ophthalmology, Jefferson Medical College, Philadelphia, Pa.

"Contains in the most attractive and easily understood form just the sort of knowledge which is necessary to the intelligent practice of general medicine and surgery."

J. WILLIAM WHITE, M. D.,
Professor of Clinical Surgery in the University of Pennsylvania.

"A very reliable guide to the study of eye diseases, presenting the latest facts and newest ideas."

SWAN M. BURNETT, M. D.,
Professor of Ophthalmology and Otology, Medical Department Univ. of Georgetown, Washington, D. C.

For Sale by Subscription.

MEDICAL DIAGNOSIS. By Dr. OSWALD VIERORDT, Professor of Medicine at the University of Heidelberg. Translated, with additions, from the Second Enlarged German Edition, with the author's permission, by FRANCIS H. STUART, A. M., M. D. Third and Revised Edition. In one handsome royal-octavo volume of 700 pages, 178 fine wood-cuts in text, many of which are in colors Prices: Cloth, $4.00 net; Sheep, $5.00 net; Half Russia, $5.50 net.

In this work, as in no other hitherto published, are given full and accurate explanations of the phenomena observed at the bedside. It is distinctly a clinical work by a master teacher, characterized by thoroughness, fulness, and accuracy. It is a mine of information upon the points that are so often passed over without explanation. Especial attention has been given to the germ-theory as a factor in the origin of disease.

This valuable work is now published in German, English, Russian, and Italian. The issue of a third American edition within two years indicates the favor with which it has been received by the profession.

"Rarely is a book published with which a reviewer can find so little fault as with the volume before us. All the chapters are full, and leave little to be desired by the reader. Each particular item in the consideration of an organ or apparatus, which is necessary to determine a diagnosis of any disease of that organ, is mentioned; nothing seems forgotten. The chapters on diseases of the circulatory and digestive apparatus and nervous system are especially full and valuable. Notwithstanding a few minor errors in translating, which are of small importance to the accuracy of the rest of the volume, the reviewer would repeat that the book is one of the best—probably *the best*—which has fallen into his hands. An excellent and comprehensive index of nearly one hundred pages closes the volume."—*University Medical Magazine,* Philadelphia.

"Thorough and exact. . . . . The author has rendered no mean service to medicine in having prepared a work which proves as useful to the teacher as to the student and practitioner."—*The Lancet* (London).

PROFESSIONAL OPINIONS.

"One of the most valuable and useful works in medical literature."

ALEXANDER J. C. SKENE, M. D.,
Dean of the Long Island College Hospital, and Professor of the Medical and Surgical Diseases of Women.

"Indispensable to both 'students and practitioners.'"

F. MINOT, M. D.,
Hersey Professor of Theory and Practice of Medicine, Harvard University.

"It is very well arranged and very complete, and contains valuable features not usually found in the ordinary books."

J. H. MUSSER, M. D.,
Assistant Professor Clinical Medicine, University of Pennsylvania.

"One of the most valuable works now before the profession, both for study and reference."

N. S. DAVIS, M. D.,
Professor of Principles and Practice of Medicine and Clinical Medicine, Chicago Medical College.

For Sale by Subscription.

A NEW PRONOUNCING DICTIONARY OF MEDICINE, with Phonetic Pronunciation, Accentuation, Etymology, etc. By JOHN M. KEATING, M. D., LL.D., Fellow of the College of Physicians of Philadelphia; Vice-President of the American Pædiatric Society; Ex-President of the Association of Life Insurance Medical Directors; Editor "Cyclopædia of the Diseases of Children," etc.; and HENRY HAMILTON, author of "A New Translation of Virgil's Æneid into English Rhyme;" co-author of "Saunders' Medical Lexicon," etc.; with the Collaboration of J. CHALMERS DACOSTA, M. D., and FREDERICK A. PACKARD, M. D. With an Appendix containing important Tables of Bacilli, Micrococci, Leucomaïnes, Ptomaïnes, Drugs and Materials used in Antiseptic Surgery, Poisons and their Antidotes, Weights and Measures, Thermometric Scales, New Official and Unofficial Drugs, etc. Forming one very attractive volume of over 800 pages. Second Revised Edition. Prices: Cloth, $5.00 net; Sheep, $6.00 net; Half Russia, $6.50 net. With Denison's Patent Index for Ready Reference.

PROFESSIONAL OPINIONS.

"I am much pleased with Keating's Dictionary, and shall take pleasure in recommending it to my classes."

HENRY M. LYMAN, M. D.,
Professor of Principles and Practice of Medicine, Rush Medical College, Chicago, Ill.

"I am convinced that it will be a very valuable adjunct to my study-table, convenient in size and sufficiently full for ordinary use."

C. A. LINDSLEY, M. D.,
Professor of Theory and Practice of Medicine, Medical Dept. Yale University;
Secretary Connecticut State Board of Health, New Haven, Conn.

"I will point out to my classes the many good features of this book as compared with others, which will, I am sure, make it very popular with students."

JOHN CRONYN, M. D., LL.D.,
Professor of Principles and Practice of Medicine and Clinical Medicine;
President of the Faculty, Medical Dept. Niagara University, Buffalo, N. Y.

AUTOBIOGRAPHY OF SAMUEL D. GROSS, M. D., Emeritus Professor of Surgery in the Jefferson Medical College of Philadelphia, with Reminiscences of His Times and Contemporaries. Edited by his sons, SAMUEL W. GROSS, M. D., LL.D., late Professor of Principles of Surgery and of Clinical Surgery in the Jefferson Medical College, and A. HALLER GROSS, A. M., of the Philadelphia Bar. Preceded by a Memoir of Dr. Gross, by the late Austin Flint, M. D., LL.D. In two handsome volumes, each containing over 400 pages, demy 8vo, extra cloth, gilt tops, with fine Frontispiece engraved on steel. Price, $5.00 net.

This autobiography, which was continued by the late eminent surgeon until within three months of his death, contains a full and accurate history of his early struggles, trials, and subsequent successes, told in a singularly interesting and charming manner, and embraces short and graphic pen-portraits of many of the most distinguished men—surgeons, physicians, divines, lawyers, statesmen, scientists, etc.—with whom he was brought in contact in America and in Europe; the whole forming a retrospect of more than three-quarters of a century.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF OBSTETRICS. By American Teachers. By Richard C. Norris, A. M., M. D.; James H. Etheridge, M. D.; Chauncey D. Palmer, M. D.; Howard A. Kelly, M. D.; Charles Jewett, M. D.; Henry J. Garrigues, M. D.; Barton Cooke Hirst, M. D.; Theophilus Parvin, M. D.; George A. Piersol, M. D.; Edward P. Davis, M. D.; Charles Warrington Earle, M. D.; Robert L. Dickinson, M. D.; Edward Reynolds, M. D.; Henry Schwarz, M. D.; and James C. Cameron, M. D. In one very handsome imperial-octavo volume, with a large number of original illustrations, including full-page plates, and uniform with "The American Text-Book of Gynecology." (In active preparation.)

Such an array of well-known teachers is a sufficient guarantee of the high character of the work, and it gives the assurance that this work will have the same measure of success awarded it as has attended the recent publication of its companion volume, "The American Text-Book of Gynecology." The illustrations will receive the most minute attention; the cuts interspersed throughout the text, and the full-page plates, which will reflect the highest attainments of the artist and engraver, will appeal at once to the eye as well as to the mind of the student and practitioner.

AN AMERICAN TEXT-BOOK OF PHYSIOLOGY. By American Teachers. Edited by WILLIAM H. HOWELL, PH. D., M. D., Professor of Physiology, Johns Hopkins University. With the collaboration of such eminent specialists as Henry P. Bowditch, M. D.; John G. Curtis, M. D.; Henry H. Donaldson, M. D.; Frederick S. Lee, M. D.; Warren P. Lombard, A. B., M. D.; Graham Lusk, PH. D.; Henry Sewall, M. D.; Edward T. Reichert, M. D.; Joseph W. Warren, M. D. In one imperial-octavo volume (with a large number of original illustrations), uniform with The American Text-Books of "Surgery," "Practice," "Gynecology," etc. (In preparation for early publication.)

This will be the most notable attempt yet made in this country to combine in one volume the entire subject of Human Physiology by well-known teachers who have given especial study to that part of the subject upon which they will write. The completed work will represent the present status of the science of Physiology, and in particular from the standpoint of the student of medicine and the medical practitioner. Illustrations largely drawn from original sources will be used freely throughout the text.

AN AMERICAN TEXT-BOOK OF APPLIED THERAPEUTICS. By American Teachers. (In preparation.)

AN AMERICAN TEXT-BOOK OF NURSING. By American Teachers. (In preparation.)

A SYLLABUS OF GYNÆCOLOGY, arranged in conformity with The American Text-Book of Gynecology. By J. W. LONG, M. D., Professor of Diseases of Women and Children, Medical College of Virginia, etc. (Preparing.)

Based upon the teaching and methods laid down in the larger work, this will not only be useful as a supplementary volume, but to those who do not already possess the text-book it will also have an independent value as an aid to the practitioner in gynecological work, and to the student as a guide in the lecture-room, as the subject is presented in a manner at once systematic, clear, succinct, and practical.

TEMPERATURE CHART. Prepared by D. T. LAINÉ, M. D. Size 8 x 13½ inches. Price, per pad of 25 charts, 50 cents.

A conveniently arranged chart for recording Temperature, with columns for daily amounts of Urinary and Fecal Excretions, Food, Remarks, etc. On the back of each chart is given in full the method of Brand in the treatment of Typhoid Fever.

THE NURSE'S DICTIONARY of Medical Terms and Nursing Treatment, containing Definitions of the Principal Medical and Nursing Terms, Abbreviations, and Physiological Names, and Descriptions of the Instruments, Drugs, Diseases, Accidents, Treatments, Operations, Foods, Appliances, etc. encountered in the ward or in the sick-room. Compiled for the use of nurses. By HONNOR MORTEN, author of "How to Become a Nurse," "Sketches of Hospital Life," etc. Second and enlarged edition. 16mo, 140 pages. Price, Cloth, $1.00.

This little volume is intended for use merely as a small reference-book which can be consulted at the bedside or in the ward. It gives sufficient explanation to the nurse to enable her to comprehend a case until she has leisure to look up larger and fuller works on the subject.

"Should be at the disposal of every nurse."—*Birmingham Medical Review.*

"Maintains its reputation for brevity and simplicity."—*Hahnemannian Monthly.*

"Though ostensibly for professional nurses, contains in a compact form just such information as almost every intelligent man would like to have at hand in these days when the interest in all matters of sanitation and medicine has become so great."—*Medical Examiner.*

"A book which every progressive nurse must have."—*Medical World.*

"This little volume is almost indispensable in the training school and in the library of the nurse."—*New York Medical Times.*

SURGICAL PATHOLOGY AND THERAPEUTICS. By J. Collins Warren, M. D., Professor of Surgery, Harvard Medical School, etc. In one very handsome octavo volume of over 800 pages, with 135 illustrations, 33 of which are chromo-lithographs, and all of which are drawn from original specimens. (Passing through the press.)

Covering as it does the entire field of Surgical Pathology and Surgical Therapeutics by an acknowledged authority, the publisher is confident that the work will rank as a standard authority on the subject of which it treats. Particular attention has been paid to Bacteriology and Surgical Bacteria from the standpoint of recent investigations, and the chromo-lithographic plates in their fidelity to nature and in scientific accuracy have hitherto been unapproached.

DISEASES OF WOMEN. By Henry J. Garrigues, A. M., M. D., Professor of Obstetrics in the New York Post-Graduate Medical School and Hospital; Gynecologist to St. Mark's Hospital and to the German Dispensary, etc., New York City. In one very handsome octavo volume of about 700 pages, illustrated by numerous wood-cuts and colored plates. Prices: Cloth, $4.00 net; Sheep, $5.00 net.

A practical work on gynecology for the use of students and practitioners, written in a terse and concise manner. The importance of a thorough knowledge of the anatomy of the female pelvic organs has been fully recognized by the author, and considerable space has been devoted to the subject. The chapters on Operations and on Treatment are thoroughly modern, and are based upon the large hospital and private practice of the author. The text is elucidated by a large number of illustrations and colored plates, many of them being original, and forming a complete atlas for studying *embryology* and the *anatomy* of the *female genitalia*, besides exemplifying, whenever needed, morbid conditions, instruments, apparatus, and operations.

EXCERPT OF CONTENTS.

Development of the Female Genitals.—Anatomy of the Female Pelvic Organs.—Physiology.—Puberty.—Menstruation and Ovulation.—Copulation.—Fecundation.—The Climacteric.—Etiology in General.—Examinations in General.—Treatment in General—Abnormal Menstruation and Metrorrhagia.—Leucorrhea.—Diseases of the Vulva.—Diseases of the Perineum.—Diseases of the Vagina.—Diseases of the Uterus.—Diseases of the Fallopian Tubes.—Diseases of the Ovaries.—Diseases of the Pelvis.—Sterility.

The reception accorded to this work has been most flattering. In the short period which has elapsed since its issue it has been adopted and recommended as a text-book by more than 60 of the Medical Schools and Universities of the United States and Canada.

"One of the best text-books for students and practitioners which has been published in the English language; it is condensed, clear, and comprehensive. The profound learning and great clinical experience of the distinguished author find expression in this book in a most attractive and instructive form. Young practitioners, to whom experienced consultants may not be available, will find in this book invaluable counsel and help."

Thad. A. Reamy, M. D., LL.D.,
Professor of Clinical Gynecology, Medical College of Ohio; Gynecologist to the Good Samaritan and Cincinnati Hospitals.

Practical, Exhaustive, Authoritative.

SAUNDERS'

NEW AID SERIES OF MANUALS

FOR

Students and Practitioners.

MR. SAUNDERS is pleased to announce as in *active preparation* his **NEW AID SERIES OF MANUALS for Students and Practitioners.** As publisher of the STANDARD SERIES OF QUESTION COMPENDS, and through intimate relations with leading members of the medical profession, Mr. Saunders has been enabled to study progressively the essential *desiderata* in practical "self-helps" for students and physicians.

This study has manifested that, while the published "Question Compends" earn the highest appreciation of students, whom they serve in reviewing their studies preparatory to examination, there is special need of thoroughly reliable handbooks on the leading branches of Medicine and Surgery, each subject being compactly and authoritatively written, and exhaustive in detail, without the introduction of *cases* and foreign subject-matter which so largely expand ordinary text-books.

The Saunders Aid Series will not merely be condensations from present literature, but will be ably written by well-known authors and practitioners, most of them being teachers in representative American Colleges. This *new series*, therefore, will form an admirable collection of advanced lectures, which will be invaluable aids to students in reading and in comprehending the contents of "recommended" works.

Each Manual will further be distinguished by the beauty of the *new* type; by the quality of the paper and printing; by the copious use of illustrations; by the attractive binding in cloth; **and by the extremely low price, which will uniformly be $1.25 per volume.**

SAUNDERS' NEW AID SERIES OF MANUALS.

VOLUMES NOW READY.

PHYSIOLOGY. By Joseph Howard Raymond, A. M., M. D., Professor of Physiology and Hygiene and Lecturer on Gynecology in the Long Island College Hospital, etc. Price, $1.25 net.

SURGERY, General and Operative. By John Chalmers DaCosta, M. D,, Demonstrator of Surgery, Jefferson Medical College, Philadelphia, etc. Double number. Price, $2.50 net.

DOSE-BOOK AND MANUAL OF PRESCRIPTION-WRITING. By E. Q. Thornton, M. D., Demonstrator of Therapeutics, Jefferson Medical College, Philadelphia. Price, $1.25 net.

MEDICAL JURISPRUDENCE. By Henry C. Chapman, M. D., Professor of Institutes of Medicine and Medical Jurisprudence in the Jefferson Medical College of Philadelphia, etc Price, $1.25 net.

SURGICAL ASEPSIS. By Carl Beck, M. D., Surgeon to St. Mark's Hospital and to the German Poliklinik; Instructor in Surgery, New York Post-Graduate Medical School, etc. Price, $1.25 net.

VOLUMES IN PREPARATION FOR EARLY PUBLICATION.

OBSTETRICS. By W. A. Newman Dorland, M. D., Demonstrator of Obstetrics, University of Pennsylvania; Chief of Gynecological Dispensary, Pennsylvania Hospital; Member of Philadelphia Obstetrical Society, etc. Price, $1.25 net.

MATERIA MEDICA AND THERAPEUTICS. By Henry A. Griffin, A. B., M. D., Assistant Physician to the Roosevelt Hospital, Out-patient Department, New York City. Price, $1.25 net.

SYPHILIS AND THE VENEREAL DISEASES. By James Nevins Hyde, M. D., Professor of Skin and Venereal Diseases in Rush Medical College, Chicago. Double number. Price, $2.50 net.

NERVOUS DISEASES. By Charles W. Burr, M. D., Clinical Professor of Nervous Diseases, Medico-Chirurgical College, Philadelphia, etc. Price, $1.25 net.

PRACTICE OF MEDICINE. By George Roe Lockwood, M. D., Professor of Practice in the Woman's Medical College and in the New York Infirmary, etc. Double number. Price, $2.50 net.

NOSE AND THROAT. By D. Braden Kyle, M. D., Chief Laryngologist to St. Agnes' Hospital, Philadelphia; Instructor in Clinical Microscopy and Assistant Demonstrator of Pathology in the Jefferson Medical College, etc. Price, $1.25 net.

⁂ There will be published in the same series, at close intervals, carefully-prepared works on the subjects of Anatomy, Gynecology, Pathology, Hygiene, etc., by prominent specialists.

Saunders' New Aid Series of Manuals.

A MANUAL OF PHYSIOLOGY. By JOSEPH H. RAYMOND, A. M., M.D., Professor of Physiology and Hygiene and Lecturer on Gynecology in the Long Island College Hospital; Director of Physiology in the Hoagland Laboratory; formerly Lecturer on Physiology and Hygiene in the Brooklyn Normal School for Physical Education; Ex-Vice-President of the American Public Health Association; Ex-Health Commissioner City of Brooklyn, etc. Illustrated. Price, Cloth, $1.25 net. (Just ready.)

In this manual the author has endeavored to put into a concrete and available form the results of twenty years' experience as a teacher of Physiology to medical students, and has produced a work for the student and practitioner, representing in a concise form the existing state of Physiology and its methods of investigation, based upon Comparative and Pathological Anatomy, Clinical Medicine, Physics, and Chemistry, as well as upon experimental research.

A MANUAL OF SURGERY, General and Operative. By JOHN CHALMERS DACOSTA, M. D., Demonstrator of Surgery, Jefferson Medical College, Philadelphia; Chief Assistant Surgeon, Jefferson Medical College Hospital; Surgical Registrar, Philadelphia Hospital, etc. One very handsome volume of over 700 pages, with a large number of illustrations. (Double number.) Price, Cloth, $2.50 net.

A new manual of the Principles and Practice of Surgery, intended to meet the demands of students and working practitioners for a medium-sized work which will embody all the newer methods of procedure detailed in the larger text-books. The work has been written in a concise, practical manner, and especial attention has been given to the most recent methods of treatment. Illustrations are freely used to elucidate the text.

A MANUAL OF OBSTETRICS. By W. A. NEWMAN DORLAND, M. D., Demonstrator of Obstetrics, University of Pennsylvania; Chief of Gynecological Dispensary, Pennsylvania Hospital; Member of Philadelphia Obstetrical Society, etc. Profusely illustrated. Price, Cloth, $1.25 net. (Preparing.)

This work, which is thoroughly practical in its teachings, is intended, as its title implies, to be a working text-book for the student and of value to the practitioner as a convenient handbook of reference. Although concisely written, nothing of importance is omitted that will give a clear and succinct knowledge of the subject as it stands to-day. Illustrations are freely used throughout the text.

Saunders' New Aid Series of Manuals.

DOSE-BOOK AND MANUAL OF PRESCRIPTION-WRITING.

By E. Q. THORNTON, M. D., Demonstrator of Therapeutics, Jefferson Medical College, Philadelphia. Illustrated. Price, Cloth, $1.25 net.

But little attention is generally given, in works on Materia Medica and Therapeutics, to the methods of combining remedies in the form of prescriptions, and this manual has been written especially for students in the hope that it may serve to give a thorough and comprehensive knowledge of the subject.

The work, which is based upon the last (1890) edition of the *Pharmacopœia*, fully covers the subjects of Weights and Measures, Prescriptions (form of writing, general directions to pharmacist, grammatical construction, etc.), Dosage, Incompatibles, Poisons, etc.

MEDICAL JURISPRUDENCE AND TOXICOLOGY.

By HENRY C. CHAPMAN, M. D., Professor of Institutes of Medicine and Medical Jurisprudence in the Jefferson Medical College of Philadelphia; Member of the College of Physicians of Philadelphia, of the Academy of Natural Sciences of Philadelphia, of the American Philosophical Society, and of the Zoological Society of Philadelphia. 232 pages, with 36 illustrations, some of which are in colors. Price, $1.25 net.

For many years there has been a demand from members of the medical and legal professions for a medium-sized work on this most important branch of medicine. The necessarily proscribed limits of the work permit the consideration only of those parts of this extensive subject which the experience of the author as coroner's physician of the city of Philadelphia for a period of six years leads him to regard as the most material for practical purposes.

Particular attention is drawn to the illustrations, many being produced in colors, thus conveying to the layman a far clearer idea of the more intricate cases.

"The salient points are clearly defined, and ascertained facts are laid down with a clearness that is unequivocal."—*St. Louis Medical and Surgical Journal.*

"The presentation is always thorough, the text is liberally interspersed with illustrations, and the style of the author is at once pleasing and interesting."—*Therapeutic Gazette.*

"One that is not overloaded with an unnecessary detail of a large amount of literature on the subject, requiring hours of research for the essential points in the decision of a question; that contains the most lucid symptomatology of questionable conditions, tests of poisons, and the readiest means of making them—such is the new book before us."—*The Sanitarian.*

A GUIDE TO THE BACTERIOLOGICAL LABORATORY.

By LANGDON FROTHINGHAM, M. D. Illustrated. (In preparation.)

The technical methods involved in bacteria-culture, methods of staining, and microscopical study are fully described and arranged as simply and concisely as possible. The book is especially intended for use in laboratory work.

NURSING: ITS PRINCIPLES AND PRACTICE. By ISABEL ADAMS HAMPTON, Graduate of the New York Training School for Nurses attached to Bellevue Hospital; Superintendent of Nurses and Principal of the Training School for Nurses, Johns Hopkins Hospital, Baltimore, Md.; late Superintendent of Nurses, Illinois Training School for Nurses, Chicago, Ill. In one very handsome 12mo volume of 484 pages, profusely illustrated. Price, Cloth, $2.00 net.

This entirely new work on the important subject of nursing is at once comprehensive and systematic. It is written in a clear, accurate, and readable style, suitable alike to the student and the lay reader. Such a work has long been a desideratum with those intrusted with the management of hospitals and the instruction of nurses in training schools. It is also of especial value to the graduated nurse who desires to acquire a practical working knowledge of the care of the sick and the hygiene of the sick-room.

The author, who has had considerable experience as superintendent of training schools for nurses and hospital management, brings to her task a mind thoroughly equipped to make the subject attractive as well as scientific and instructive.

Thoroughly attested and approved processes in practical nursing only have been given, particularly in antiseptic surgery, and the minutest details regarding the nurse's technique have been explained.

Illustrations to elucidate the text have been used freely throughout the book, and they will be found of material help in showing the forms of modern appliances for the hospital ward and sick-room, the registration of temperature, daily records, etc.

METHODS OF PREVENTING AND CORRECTING DEFORMITIES OF THE BONES AND JOINTS: A Handbook of Practical Orthopedic Surgery. By H. AUGUSTUS WILSON, M. D, Professor of General and Orthopedic Surgery, Philadelphia Polyclinic; Clinical Professor of Orthopedic Surgery, Jefferson Medical College, Philadelphia, etc. (In preparation.)

The aim of the author is to provide a book of moderate size, containing comprehensive details that will enable general practitioners to understand thoroughly the mechanical features of the many forms of congenital and acquired deformities of the bones and joints.

The mechanical functions that are impaired will be considered first as to prevention as of primary importance, and following this will be described the methods of correction that have been proved practical by the author. Operative procedures will be considered from a mechanical as well as a surgical standpoint. Prominence will be given to the mechanical requirements for braces and artificial limbs, etc., with description of the methods for constructing the simplest forms, whether made of plaster of Paris, felt, leather, paper, steel, or other materials, together with the methods of readjustment to suit the changes occurring during the progress of the case. A very large number of original illustrations will be used.

AN OPERATION BLANK, with Lists of Instruments, etc. required in Various Operations. Prepared by W. W. KEEN, M. D., LL.D., Professor of Principles of Surgery in the Jefferson Medical College, Philadelphia. Price per Pad, containing Blanks for fifty operations, 50 cents net.

A convenient blank, suitable for all operations, giving complete instructions regarding necessary preparation of patient, etc., with a full list of dressings and medicines to be employed.

At the back of pad is a list of instruments used—viz. general instruments, etc., required for all operations; and special instruments for surgery of the brain and spine, mouth and throat, abdomen, rectum, male and female genito-urinary organs, the bones, etc.

The whole forming a neat pad, arranged for hanging on the wall of a surgeon's office or in the hospital operating-room.

"Will serve a useful purpose for the surgeon in reminding him of the details of preparation for the patient and the room as well as for the instruments, dressings, and antiseptics needed"—*New York Medical Record*

"Covers about all that can be needed in any operation."—*American Lancet.*

"The plan is a capital one."—*Boston Medical and Surgical Journal.*

ESSENTIALS OF ANATOMY AND MANUAL OF PRACTICAL DISSECTION, containing "Hints on Dissection." By CHARLES B. NANCREDE, M. D., Professor of Surgery and Clinical Surgery in the University of Michigan, Ann Arbor; Corresponding Member of the Royal Academy of Medicine, Rome, Italy; late Surgeon Jefferson Medical College, etc. Fourth and revised edition. Post 8vo, over 500 pages, with handsome full-page lithographic plates in colors, and over 200 illustrations. Price: Extra Cloth or Oilcloth for the dissection-room, $2.00 net.

Neither pains nor expense has been spared to make this work the most exhaustive yet concise Student's Manual of Anatomy and Dissection ever published, either in America or in Europe.

The colored plates are designed to aid the student in dissecting the muscles, arteries, veins, and nerves. The wood-cuts have all been specially drawn and engraved, and an Appendix added containing 60 illustrations representing the structure of the entire human skeleton, the whole being based on the eleventh edition of Gray's *Anatomy*, and forming a handsome post 8vo volume of over 500 pages.

"The plates are of more than ordinary excellence, and are of especial value to students in their work in the dissecting-room."—*Journal of American Medical Association.*

"Should be in the hands of every medical student."—*Cleveland Medical Gazette.*

"A concise and judicious work."—*Buffalo Medical and Surgical Journal.*

A MANUAL OF PRACTICE OF MEDICINE. By A. A. STEVENS, A. M., M. D., Instructor of Physical Diagnosis in the University of Pennsylvania, and Demonstrator of Pathology in the Woman's Medical College of Philadelphia. Specially intended for students preparing for graduation and hospital examinations, and includes the following sections: General Diseases, Diseases of the Digestive Organs, Diseases of the Respiratory System, Diseases of the Circulatory System, Diseases of the Nervous System, Diseases of the Blood, Diseases of the Kidneys, and Diseases of the Skin. Each section is prefaced by a chapter on General Symptomatology. Third edition. Post 8vo, 502 pages. Numerous illustrations and selected formulæ. Price, $2.50.

Contributions to the science of medicine have poured in so rapidly during the last quarter of a century that it is well-nigh impossible for the student, with the limited time at his disposal, to master elaborate treatises or to cull from them that knowledge which is absolutely essential. From an extended experience in teaching, the author has been enabled, by classification, to group allied symptoms, and by the judicious elimination of theories and redundant explanations to bring within a comparatively small compass a complete outline of the practice of medicine.

A SYLLABUS OF LECTURES ON THE PRACTICE OF SURGERY, arranged in conformity with The American Text-Book of Surgery. By NICHOLAS SENN, M. D., PH. D., Professor of Surgery in Rush Medical College, Chicago, and in the Chicago Polyclinic. Price, $2.00.

This, the latest work of its eminent author, himself one of the contributors to the "American Text-Book of Surgery," will prove of exceptional value to the advanced student who has adopted that work as his text-book. It is not only the syllabus of an unrivalled course of surgical practice, but it is also an epitome or supplement to the larger work.

SYLLABUS OF OBSTETRICAL LECTURES in the Medical Department, University of Pennsylvania. By RICHARD C. NORRIS, A. M., M. D., Demonstrator of Obstetrics in the University of Pennsylvania. Third edition, thoroughly revised and enlarged. Crown 8vo. Price, Cloth, interleaved for notes, $2.00 net.

"This work is so far superior to others on the same subject that we take pleasure in calling attention briefly to its excellent features. It covers the subject thoroughly, and will prove invaluable both to the student and the practitioner. The author has introduced a number of valuable hints which would only occur to one who was himself an experienced teacher of obstetrics. The subject-matter is clear, forcible, and modern. We are especially pleased with the portion devoted to the practical duties of the accoucheur, care of the child, etc. The paragraphs on antiseptics are admirable; there is no doubtful tone in the directions given. No details are regarded as unimportant; no minor matters omitted. We venture to say that even the old practitioner will find useful hints in this direction which he cannot afford to despise."—*New York Medical Record.*

OUTLINES OF OBSTETRICS: A Syllabus of Lectures Delivered at Long Island College Hospital. By CHARLES JEWETT, A. M., M. D., Professor of Obstetrics and Pediatrics in the College, and Obstetrician to the Hospital. Edited by HAROLD F. JEWETT, M. D. Post 8vo, 264 pages. Price, $2.00.

This book treats only of the general facts and principles of obstetrics: these are stated in concise terms and in a systematic and natural order of sequence, theoretical discussion being as far as possible avoided; the subject is thus presented in a form most easily grasped and remembered by the student. Special attention has been devoted to practical questions of diagnosis and treatment, and in general particular prominence is given to facts which the student most needs to know. The condensed form of statement and the orderly arrangement of topics adapt it to the wants of the busy practitioner as a means of refreshing his knowledge of the subject and as a handy manual for daily reference.

NOTES ON THE NEWER REMEDIES: their Therapeutic Applications and Modes of Administration. By DAVID CERNA, M. D., PH. D., Demonstrator of and Lecturer on Experimental Therapeutics in the University of Pennsylvania. Post-octavo, 175 pages. Price, $1.25.

The work takes up in alphabetical order all the newer remedies, giving their physical properties, solubility, therapeutic applications, administration, and chemical formula.

It thus forms a very valuable addition to the various works on therapeutics now in existence.

Chemists are so multiplying compounds, that, if each compound is to be thoroughly studied, investigations must be carried far enough to determine the practical importance of the new agents.

"Especially valuable because of its completeness, its accuracy, its systematic consideration of the properties and therapy of many remedies of which doctors generally know but little, expressed in a brief yet terse manner."—*Chicago Clinical Review.*

"A timely and needful book . . . . which physicians who avail themselves of the use of the newer remedies cannot afford to do without."—*The Sanitarian.*

LABORATORY EXERCISES IN BOTANY. By EDSON S. BASTIN, M. A., Professor of Materia Medica and Botany in the Philadelphia College of Pharmacy. With over 75 plates. (In preparation.)

This work is intended for the beginner and the advanced student, and it fully covers the structure of flowering plants, roots, ordinary stems, rhizomes, tubers, bulbs, leaves, flowers, fruits, and seeds. Particular attention is given to the gross and microscopical structure of plants, and to those used in medicine. Illustrations have freely been used to elucidate the text, and a complete index to facilitate reference has been added.

The folding charts which supplement the subjects will be found useful in connection with the study of the text.

SAUNDERS' POCKET MEDICAL LEXICON; or, Dictionary of Terms and Words used in Medicine and Surgery. By JOHN M. KEATING, M. D., editor of "Cyclopædia of Diseases of Children," etc.; author of the "New Pronouncing Dictionary of Medicine; and HENRY HAMILTON, author of "A New Translation of Virgil's Æneid into English Verse;" co-author of a "New Pronouncing Dictionary of Medicine." A new and revised edition. 32mo, 282 pages. Prices: Cloth, 75 cents; Leather Tucks, $1.00.

This new and comprehensive work of reference is the outcome of a demand for a more modern handbook of its class than those at present on the market, which, dating as they do from 1855 to 1884, are of but trifling use to the student by their not containing the hundreds of new words now used in current literature, especially those relating to Electricity and Bacteriology.

"Remarkably accurate in terminology, accentuation, and definition."—*Journal of American Medical Association.*

"Brief, yet complete . . . . it contains the very latest nomenclature in even the newest departments of medicine."—*New York Medical Record.*

SAUNDERS' POCKET MEDICAL FORMULARY. By WILLIAM M. POWELL, M. D., Attending Physician to the Mercer House for Invalid Women at Atlantic City. Containing 1750 Formulæ, selected from several hundred of the best-known authorities. Forming a handsome and convenient pocket companion of nearly 300 printed pages, with blank leaves for Additions; with an Appendix containing Posological Table, Formulæ and Doses for Hypodermatic Medication, Poisons and their Antidotes, Diameters of the Pemale Pelvis and Fœtal Head, Obstetrical Table, Diet List for Various Diseases, Materials and Drugs used in Antiseptic Surgery, Treatment of Asphyxia from Drowning, Surgical Remembrancer, Tables of Incompatibles, Eruptive Fevers, Weights and Measures, etc. Third edition, revised and greatly enlarged. Handsomely bound in morocco, with side index, wallet, and flap. Price, $1.75 net.

A concise, clear, and correct record of the many hundreds of famous formulæ which are found scattered through the works of the *most eminent physicians and surgeons* of the world. The work is helpful to the student and practitioner alike, as through it they become acquainted with numerous formulæ which are not found in text-books, but have been collected from among *the rising generation of the profession, college professors, and hospital physicians and surgeons.*

"This little book, that can be conveniently carried in the pocket, contains an immense amount of material. It is very useful, and as the name of the author of each prescription is given is unusually reliable."—*New York Medical Record.*

"Designed to be of immense help to the general practitioner in the exercise of his daily calling."—*Boston Medical and Surgical Journal.*

HOW TO EXAMINE FOR LIFE INSURANCE. By JOHN M. KEATING, M. D., Fellow of the College of Physicians and Surgeons of Philadelphia; Vice-President of the American Pædiatric Society; Ex-President of the Association of Life Insurance Medical Directors. Royal 8vo, 211 pages, with two large phototype illustrations, and a plate prepared by Dr. McClellan from special dissections; also, numerous cuts to elucidate the text. Second edition. Price, in Cloth, $2.00 net.

PART I., which has been carefully prepared from the best works on Physical Diagnosis, is a short and succinct account of the methods used to make examinations; a description of the normal condition and of the earliest evidences of disease.

PART II. contains the Instructions of twenty-four Life Insurance Companies to their medical examiners.

"This is by far the most useful book which has yet appeared on insurance examination, a subject of growing interest and importance. Not the least valuable portion of the volume is Part II., which consists of instructions issued to their examining physicians by twenty-four representative companies of this country. As the proofs of these instructions were corrected by the directors of the companies, they form the latest instructions obtainable. If for these alone, the book should be at the right hand of every physician interested in this special branch of medical science."—*The Medical News*, Philadelphia.

MANUAL OF MATERIA MEDICA AND THERAPEUTICS. By A. A. STEVENS, A. M., M. D., Instructor of Physical Diagnosis in the University of Pennsylvania, and Demonstrator of Pathology in the Woman's Medical College of Philadelphia. 435 pages. Price, Cloth, $2.25.

This wholly new volume, which is based on the 1890 edition of the *Pharmacopœia*, comprehends the following sections: Physiological Action of Drugs; Drugs; Remedial Measures other than Drugs; Applied Therapeutics; Incompatibility in Prescriptions; Table of Doses; Index of Drugs; and Index of Diseases; the treatment being elucidated by more than two hundred formulæ.

"The author is to be congratulated upon having presented the medical student with as accurate a manual of therapeutics as it is possible to prepare."—*Therapeutic Gazette.*

"Far superior to most of its class; in fact, it is very good. Moreover, the book is reliable and accurate."—*New York Medical Journal.*

"The author has faithfully presented modern therapeutics in a comprehensive work, . . . and it will be found a reliable guide."—*University Medical Magazine.*

"Will be of immense service to the busy practitioner."—*Medical Reporter* (Calcutta).

"Reliable and timely."—*North American Practitioner.*

"Concise, up to date, and withal comprehensive."—*Pacific Medical Journal.*

SAUNDERS' QUESTION COMPENDS.

Arranged in Question and Answer Form.

THE LATEST, CHEAPEST, and BEST ILLUSTRATED SERIES OF COMPENDS EVER ISSUED.

Now the Standard Authorities in Medical Literature

WITH

Students and Practitioners in every City of the United States and Canada.

THE REASON WHY.

They are the advance guard of "Student's Helps"—that DO HELP; they are the leaders in their special line, *well and authoritatively written by able men, who, as teachers in the large colleges, know exactly what is wanted by a student preparing for his examinations.* The judgment exercised in the selection of authors is fully demonstrated by their professional elevation. Chosen from the ranks of Demonstrators, Quiz-masters, and Assistants, most of them have become Professors and Lecturers in their respective colleges.

Each book is of convenient size (5 x 7 inches), containing on an average 250 pages, profusely illustrated, and elegantly printed in clear, readable type, on fine paper.

The entire series, numbering twenty-four subjects, has been kept thoroughly revised and enlarged when necessary, many of them being in their fourth and fifth editions.

TO SUM UP.

Although there are numerous other Quizzes, Manuals, Aids, etc. in the market, none of them approach the "Blue Series of Question Compends;" and the claim is made for the following points of excellence:

1. Professional distinction and reputation of authors.
2. Conciseness, clearness, and soundness of treatment.
3. Size of type and quality of paper and binding.

⁂ Any of these Compends will be mailed on receipt of price.

1. **ESSENTIALS OF PHYSIOLOGY.** By H. A. HARE, M. D., Professor of Therapeutics and Materia Medica in the Jefferson Medical College of Philadelphia; Physician to St. Agnes' Hospital and to the Medical Dispensary of the Children's Hospital; Laureate of the Royal Academy of Medicine in Belgium, of the Medical Society of London, etc. Third edition, revised and enlarged by the addition of a series of handsome plate illustrations taken from the celebrated "Icones Nervorum Capitis" of Arnold. Crown 8vo, 230 pages, numerous illustrations. Price, Cloth, $1.00 net; interleaved for notes, $1.25 net.

"An exceedingly useful little compend. The author has done his work thoroughly and well. The plates of the cranial nerves from Arnold are superb."—*Journal of American Medical Association.*

2. **ESSENTIALS OF SURGERY,** containing also Venereal Diseases, Surgical Landmarks, Minor and Operative Surgery, and a Complete Description, together with full Illustrations, of the Handkerchief and Roller Bandages. By EDWARD MARTIN, A. M., M. D., Clinical Professor of Genito-Urinary Diseases, Instructor in Operative Surgery, and Lecturer on Minor Surgery, University of Pennsylvania; Surgeon to the Howard Hospital; Assistant Surgeon to the University Hospital, etc. Fifth edition. Crown 8vo, 334 pages, profusely illustrated. Considerably enlarged by an Appendix containing full directions and prescriptions for the preparation of the various materials used in Antiseptic Surgery; also several hundred recipes covering the medical treatment of surgical affections. Price, Cloth, $1.00; interleaved for notes, $1.25.

"Written to assist the student, it will be of undoubted value to the practitioner, containing as it does the essence of surgical work."—*Boston Medical and Surgical Journal.*

"Cleverly combines all the merits of condensation, while avoiding the errors of superficiality and inaccuracy."—*University Medical Magazine.*

3. **ESSENTIALS OF ANATOMY, including the Anatomy of the Viscera.** By CHARLES B. NANCREDE, M. D., Professor of Surgery and of Clinical Surgery in the University of Michigan, Ann Arbor; Corresponding Member of the Royal Academy of Medicine, Rome, Italy; late Surgeon to the Jefferson Medical College, etc. Fifth edition. Crown 8vo, 380 pages, 180 illustrations. Enlarged by an Appendix containing over sixty illustrations of the Osteology of the Human Body. The whole based upon the last (eleventh) edition of Gray's *Anatomy.* Price, Cloth, $1.00; interleaved for notes, $1.25.

"Truly such a book as no student can afford to be without."—*American Practitioner and News.*

"The questions have been wisely selected and the answers accurately and concisely given."—*University Medical Magazine.*

4. **ESSENTIALS OF MEDICAL CHEMISTRY, ORGANIC AND INORGANIC,** containing also Questions on Medical Physics, Chemical Physiology, Analytical Processes, Urinalysis, and Toxicology. By LAWRENCE WOLFF, M D., Demonstrator of Chemistry, Jefferson Medical College; Visiting Physician to the German Hospital of Philadelphia; Member of Philadelphia College of Pharmacy, etc. Fourth and revised edition, with an Appendix. Crown 8vo, 212 pages. Price, Cloth, $1.00; interleaved for notes, $1.25.

"The scope of this work is certainly equal to that of the best course of lectures on Medical Chemistry."—*Pharmaceutical Era.*

"We could wish that more books like this would be written, in order that medical students might thus early become interested in what is often a difficult and uninteresting branch of medical study."—*Medical and Surgical Reporter.*

5. **ESSENTIALS OF OBSTETRICS.** By W. EASTERLY ASHTON, M. D., Professor of Gynecology in the Medico-Chirurgical College of Philadelphia; Obstetrician to the Philadelphia Hospital. Third edition, thoroughly revised and enlarged. Crown 8vo, 244 pages, 75 illustrations. Price, Cloth, $1.00; interleaved for notes, $1.25.

"An excellent little volume containing correct and practical knowledge. An admirable compend, and the best condensation we have seen."—*Southern Practitioner.*

"Of extreme value to students, and an excellent little book to freshen up the memory of the practitioner."—*Chicago Medical Times.*

6. **ESSENTIALS OF PATHOLOGY AND MORBID ANATOMY.** By C. E. ARMAND SEMPLE, B. A., M. B., Cantab. L. S. A., M. R. C. P. Lond., Physician to the Northeastern Hospital for Children, Hackney; Professor of Vocal and Aural Physiology and Examiner in Acoustics at Trinity College, London, etc. Crown 8vo, 174 pages, illustrated. Sixth thousand. Price, Cloth, $1.00; interleaved for notes, $1.25.

"A valuable little volume—truly a *multum in parvo.*"—*Cincinnati Medical News.*

"The volume is very comprehensive, covering the entire field of pathology."—*St. Joseph Medical Herald.*

7. **ESSENTIALS OF MATERIA MEDICA, THERAPEUTICS, AND PRESCRIPTION-WRITING.** By HENRY MORRIS, M. D., late Demonstrator, Jefferson Medical College; Fellow of the College of Physicians, Philadelphia; co-editor Biddle's Materia Medica; Visiting Physician to St. Joseph's Hospital, etc. Fourth edition. Crown 8vo, 250 pages. Price, Cloth, $1.00; interleaved for notes, $1.25.

"One of the best compends in this series. Concise, pithy, and clear, well suited to the purpose for which it is prepared."—*Medical and Surgical Reporter.*

"The subjects are treated in such a unique and attractive manner that they cannot fail to impress the mind and instruct in a lasting manner."—*Buffalo Medical and Surgical Journal.*

8, 9. **ESSENTIALS OF PRACTICE OF MEDICINE.** By HENRY MORRIS, M. D., author of "Essentials of Materia Medica," etc., with an Appendix on the Clinical and Microscopical Examination of Urine, by LAWRENCE WOLFF, M. D., author of "Essentials of Medical Chemistry," etc. Colored (Vogel) urine scale and numerous fine illustrations. Third edition, enlarged by some three hundred essential formulæ, selected from the writings of the most eminent authorities of the medical profession, collected and arranged by WILLIAM M. POWELL, M. D., author of "Essentials of Diseases of Children." Crown 8vo, 460 pages. Price, Cloth, $2.00.

"The teaching is sound, the presentation graphic, matter as full as might be desired, and the style attractive."—*American Practitioner and News.*

"A first-class practice of medicine boiled down, and giving the real essentials in as few words as is consistent with a thorough understanding of the subject."—*Medical Brief.*

"Especially full, and an excellent illustration of what the best of the compends can be made to be."—*Gaillard's Medical Journal.*

10. **ESSENTIALS OF GYNÆCOLOGY.** By EDWIN B. CRAGIN, M. D., Attending Gynæcologist, Roosevelt Hospital, Out-Patients' Department; Assistant Surgeon, New York Cancer Hospital, etc. Fourth edition, revised. Crown 8vo, 198 pages, 62 fine illustrations. Price, Cloth, $1.00; interleaved for notes, $1.25.

"This is a most excellent addition to this series of question compends. The style is concise, and at the same time the sentences are well rounded. This renders the book far more easy to read than most compends, and adds distinctly to its value."—*Medical and Surgical Reporter.*

"Useful not only to the student who is barely at the threshold of professional life, but to the busy practitioner as well."—*New York Medical Journal.*

11. **ESSENTIALS OF DISEASES OF THE SKIN.** By HENRY W. STELWAGON, M. D., Clinical Lecturer on Dermatology in the Jefferson Medical College, Philadelphia; Physician to the Skin Service of the Northern Dispensary; Dermatologist to Philadelphia Hospital; Physician to Skin Department of the Howard Hospital; Clinical Professor of Dermatology in the Woman's Medical College, Philadelphia, etc. Third edition. Crown 8vo, 270 pages, 86 illustrations, many of which are original. Price, Cloth, $1.00; interleaved for notes, $1.25 net.

"An immense amount of literature has been gone over and judiciously condensed by the writer's skill and experience."—*New York Medical Record.*

"The book admirably answers the purpose for which it is written. The experience of the reviewer has taught him that just such a book is needed."—*New York Medical Journal.*

12. **ESSENTIALS OF MINOR SURGERY, BANDAGING, AND VENEREAL DISEASES.** By EDWARD MARTIN, A. M., M. D., author of "Essentials of Surgery," etc. Second edition. Crown 8vo, thoroughly revised and enlarged, 78 illustrations. Price, Cloth, $1.00; interleaved for notes, $1.25.

"Characterized by the same literary excellence that has distinguished previous numbers of this series of compends."—*American Practitioner and News.*

"The best condensation of the subjects of which it treats yet placed before the profession."—*Medical News,* Philadelphia.

"A capital little book. The illustrations are remarkably clear and intelligible."—*Australian Medical Gazette.*

"We have nothing but praise for the subject-matter of this book."—*Bristol Medico-Chirurgical Journal.*

13. **ESSENTIALS OF LEGAL MEDICINE, TOXICOLOGY, AND HYGIENE.** By C. E. ARMAND SEMPLE, M. D., author of "Essentials of Pathology and Morbid Anatomy." Crown 8vo, 212 pages, 130 illustrations. Price, Cloth, $1.00; interleaved for notes, $1.25.

"The leading points, the essentials of this too much neglected portion of medical science, are here summed up systematically and clearly."—*Southern Practitioner.*

"But for the author's judicious condensation of facts, the information it contains would be sufficient to fill an ordinary octavo volume."—*College and Clinical Record.*

14. **ESSENTIALS OF REFRACTION AND DISEASES OF THE EYE.** By EDWARD JACKSON, A. M., M. D., Professor of Diseases of the Eye in the Philadelphia Polyclinic and College for Graduates in Medicine; Member of the American Ophthalmological Society; Fellow of the College of Physicians of Philadelphia; Fellow of the American Academy of Medicine, etc.; and **ESSENTIALS OF DISEASES OF THE NOSE AND THROAT.** By E. BALDWIN GLEASON, S. B., M. D., Clinical Professor of Otology, Medico-Chirurgical College, Philadelphia; Surgeon in charge of the Nose, Throat, and Ear Department of the Northern Dispensary of Philadelphia; formerly Assistant in the Nose and Throat Dispensary of the Hospital of the University of Pennsylvania, and Assistant in the Nose and Throat Department of the Union Dispensary, etc. Two volumes in one. Second edition. Crown 8vo, 294 pages, 124 illustrations. Price, Cloth, $1.00; interleaved for notes, $1.25.

"A valuable book to the beginner in these branches, to the student, to the busy practitioner, and as an adjunct to more thorough reading. The authors are capable men, and as successful teachers, know what a student most needs."—*New York Medical Record.*

"Very valuable, since in both sections is given about all that a candidate for examination is required to know."—*Medical Times and Hospital Gazette.*

15. **ESSENTIALS OF DISEASES OF CHILDREN.** By WILLIAM M. POWELL, M. D., Attending Physician to the Mercer House for Invalid Women at Atlantic City, N. J.; late Physician to the Clinic for the Diseases of Children in the Hospital of the University of Pennsylvania and St. Clement's Hospital; Instructor in Physical Diagnosis in the Medical Department of the University of Pennsylvania. Crown 8vo, 216 pages. Price, Cloth, $1.00; interleaved for notes, $1.25.

"This work is gotten up in the clear and attractive style that characterizes the Saunders Series. It contains in appropriate form the gist of all the best works in the department to which it relates."—*American Practitioner and News.*

"The book contains a series of important questions and answers, which the student will find of great utility in the examination of children."—*Annals of Gynæcology.*

16. **ESSENTIALS OF EXAMINATION OF URINE.** By LAWRENCE WOLFF, M. D., author of "Essentials of Medical Chemistry," etc. Colored (Vogel) urine scale and numerous illustrations. Crown 8vo. Price, Cloth, 75 cents.

"A little work of decided value."—*University Medical Magazine.*

"A good manual for students, well written, and answers, categorically, many questions beginners are sure to ask."—*New York Medical Record.*

"The questions have been well chosen, and the answers are clear and brief. The book cannot fail to be useful to students."—*Medical and Surgical Reporter.*

17. **ESSENTIALS OF DIAGNOSIS.** By SOLOMON SOLIS-COHEN, M. D., Professor of Clinical Medicine and Applied Therapeutics in the Philadelphia Polyclinic, and AUGUSTUS A. ESHNER, M. D., Instructor in Clinical Medicine, Jefferson Medical College, Philadelphia. Crown 8vo, 382 pages, 55 illustrations, some of which are colored, and a frontispiece. Price, $1.50 net.

"A good book for the student, properly written from their standpoint, and confines itself well to its text."—*New York Medical Record.*

"Concise in the treatment of the subject, terse in expression of fact. . . . The work is reliable, and represents the accepted views of clinicians of to-day."—*American Journal of Medical Sciences.*

"The subjects are explained in a few well-selected words, and the required ground has been thoroughly gone over."—*International Medical Magazine.*

18. **ESSENTIALS OF PRACTICE OF PHARMACY.** By LUCIUS E. SAYRE, M. D., Professor of Pharmacy and Materia Medica in the University of Kansas. Second edition, revised and enlarged. Crown 8vo, 200 pages. Price, Cloth, $1.00; interleaved for notes, $1.25.

"Covers a great deal of ground in small compass. The matter is well digested and arranged. The research questions are a valuable feature of the book."—*Albany Medical Annals.*

"The best quiz on Pharmacy we have yet examined."—*National Drug Register.*

"The veteran pharmacist can peruse it with pleasure, because it emphasizes his grasp upon knowledge already gleaned."—*Western Drug Record.*

20. **ESSENTIALS OF BACTERIOLOGY: A Concise and Systematic Introduction to the Study of Micro-organisms.** By M. V. BALL, M. D., Assistant in Microscopy, Niagara University, Buffalo, N. Y.; late Resident Physician, German Hospital, Philadelphia, etc. Second edition, revised. Crown 8vo, 200 pages, 81 illustrations, some in colors, and 5 plates. Price, Cloth, $1.00; interleaved for notes, $1.25.

"The amount of material condensed in this little book is so great, and so accurate are the formulæ and methods, that it will be found useful as a laboratory handbook."—*Medical News.*

"Bacteriology is the keynote of future medicine, and every physician who expects success must familiarize himself with a knowledge of germ-life—the agents of disease. This little book, with its beautiful illustrations, will give the students, in brief, the results of years of study and research unaided."—*Pacific Record of Medicine and Surgery.*

"Thoroughly practical, very concise, clear, well-written, and sufficiently illustrated. . . . The best book of the kind in the English language."—*Medical and Surgical Reporter.*

21. **ESSENTIALS OF NERVOUS DISEASES AND INSANITY, their Symptoms and Treatment.** By JOHN C. SHAW, M. D., Clinical Professor of Diseases of the Mind and Nervous System, Long Island College Hospital Medical School; Consulting Neurologist to St. Catherine's Hospital and to the Long Island College Hospital; formerly Medical Superintendent King's County Insane Asylum. Second edition. Crown 8vo, 186 pages, 48 original illustrations, mostly selected from the Author's private practice. Price, Cloth, $1.00; interleaved for notes, $1.25.

"Clearly and intelligently written."—*Boston Medical and Surgical Journal.*

"A valuable addition to this series of compends, and one that cannot fail to be appreciated by all physicians and students."—*Medical Brief.*

"Dr. Shaw's Primer is excellent. The engravings are well executed and very interesting."—*Medical Times and Register.*

"Written with great clearness, devoid of verbosity, it encompasses in a brief space a vast amount of valuable information."—*Pacific Medical Record.*

22. **ESSENTIALS OF PHYSICS.** By FRED J. BROCKWAY, M. D., Assistant Demonstrator of Anatomy in the College of Physicians and Surgeons, New York. Second edition. Crown 8vo, 320 pages, 155 fine illustrations. Price, Cloth, $1.00 net; interleaved for notes, $1.25 net.

"The publisher has again shown himself as fortunate in his editor as he ever has been in the attractive style and make-up of his compends."—*American Practitioner and News.*

"Contains all that one need know of the subject, is well written, and is copiously illustrated."—*New York Medical Record.*

"The author has dealt with the subject in a manner that will make the theme not only comparatively easy, but also of interest."—*Medical News,* Philadelphia.

"Deserving of close investigation at the hands of students and physicians."—*American Gynecological Journal.*

23. **ESSENTIALS OF MEDICAL ELECTRICITY.** By D. D. STEWART, M. D., Demonstrator of Diseases of the Nervous System and Chief of the Neurological Clinic in the Jefferson Medical College; Physician to St. Mary's Hospital and to St. Christopher's Hospital for Children, etc.; and E. S. LAWRANCE, M. D., Chief of the Electrical Clinic, and Assistant Demonstrator of Diseases of the Nervous System in the Jefferson Medical College, etc. Crown 8vo, 148 pages, 65 illustrations. Price, Cloth, $1.00; interleaved for notes, $1.25.

"Clearly written, and affords a safe guide to the beginner in this subject."—*Boston Medical and Surgical Journal.*

"The subject is presented in a lucid and pleasing manner."—*New York Medical Record.*

"A little work on an important subject, which will prove of great value to medical students and trained nurses who wish to study the scientific as well as the practical points of electricity."—*The Hospital,* London.

"The selection and arrangement of material are done in a skilful manner. It gives, in a condensed form, the principles and science of electricity and their application in the practice of medicine."—*Annals of Surgery.*

"The compilation is a good one, and will be found useful both to students and to men in practice."—*New Zealand Medical Journal.*

24. **ESSENTIALS OF DISEASES OF THE EAR.** By E. B. GLEASON, S. B., M. D., Clinical Professor of Otology, Medico-Chirurgical College, Philadelphia; Surgeon in Charge of the Nose, Throat, and Ear Department of the Northern Dispensary of Philadelphia; formerly Assistant in the Nose and Throat Dispensary of the Hospital of the University of Pennsylvania, and Assistant in the Nose and Throat Department of the Union Dispensary. 89 illustrations. Price, Cloth, $1.00; interleaved for notes, $1.25.

This latest addition to the Saunders Compend Series accurately represents the modern aspect of otological science. The effort has been made to state the Essentials of Otology concisely, without sacrificing accuracy to brevity, and the book, while small in compass, is logically and capably written; it comprises upward of 150 pages, with 89 illustrations, most of which are from original sources.

LANE MEDICAL LIBRARY

To avoid fine, this book should be returned on
or before the date last stamped below.

E32 Nancrede, C.B.G.de.
N17 Essentials of anatomy.
1894 65714

NAME	DATE DUE